

0.25 was too high, and there were many peaks in the difference electron-density map which could not be incorporated into the model. A twin model was then proposed such that the crystal is composed of two twin domains, with the reciprocal lattices of these two components being coincident:  $I(hkl) = (1 - \alpha)I(hkl) + \alpha(\bar{k}\bar{h}\bar{l})$ . In *SHELXL93* (Sheldrick, 1993), this is incorporated into the least-squares refinement by a twin matrix of  $(010\bar{1}00/001)$  and refinement of the twin fraction  $\alpha$ . Application of this twin law resulted in a dramatic drop in the *R* factor to 0.076 [isotropic refinement on  $I > 3\sigma(I)$ ] and a clean difference electron-density map. This is an example of a twin by merohedry with the crystal twinned on the (110) plane. A linear decay correction was applied to the data (*TEXSAN*; Molecular Structure Corporation, 1989). Averaging the symmetry equivalent reflections within the two measured octants,  $(+h,+k,+l)$  and  $(-h,+k,+l)$ , resulted in an  $R_{\text{int}}$  value of 0.022. The twin fraction refined to a value of 0.334 (1). Three reflections were omitted from the refinement because of large negative  $F_o$  values, i.e.  $(\bar{2},18,1)$ ,  $(\bar{1},14,2)$  and  $(\bar{1}\bar{1},17,1)$ . H atoms for both structures were included in the model at calculated positions using a riding model, with aromatic C—H = 0.93, secondary C—H = 0.97, tertiary C—H = 0.98 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{bonded C atom})$ . *PLATON* (Spek, 1990) was used to calculate some metric parameters.

For both compounds, data collection: *MSCIAFC Diffractometer Control Software* (Molecular Structure Corporation, 1993); cell refinement: *MSCIAFC Diffractometer Control Software*. Data reduction: *TEXSAN* (Molecular Structure Corporation, 1995) for (3); *TEXSAN* (Molecular Structure Corporation, 1989) for (4). For both compounds, program(s) used to solve structures: *SHELXS86*; program(s) used to refine structures: *SHELXL93*; molecular graphics: *ORTEPII* (Johnson, 1976).

The authors thank the National Science Foundation for financial support. The diffractometer was purchased with a grant from the National Institutes of Health.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FR1123). Services for accessing these data are described at the back of the journal.

## References

- Allegra, G. & Ganis, P. (1962). *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Nat. Rend.* **33**, 438–449.
- Gallucci, J. C., Kozmina, N., Paquette, L. A., Zaegel, F., Meunier, P. & Gautheron, B. (1997). *Acta Cryst.* **C53**, 1416–1420.
- Gowik, P., Klapötke, T. & Pickardt, J. (1990). *J. Organomet. Chem.* **393**, 343–348.
- Howells, E. R., Phillips, D. C. & Rogers, D. (1950). *Acta Cryst.* **3**, 210–214.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Molecular Structure Corporation (1989). *TEXSAN. Single Crystal Structure Analysis Software*. Version 5.0. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Molecular Structure Corporation (1993). *MSCIAFC Diffractometer Control Software*. Version 4.3.0. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Molecular Structure Corporation (1995). *TEXSAN. Single Crystal Structure Analysis Software*. Version 1.7. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.

- Paquette, L. A., Charumilind, P., Kravetz, T. M., Böhm, M. C. & Gleiter, R. (1983). *J. Am. Chem. Soc.* **105**, 3126–3135.
- Paquette, L. A. & Sivik, M. R. (1992). *Organometallics*, **11**, 3503–3505.
- Parthasarathi, V., Beurskens, P. T. & Slot, H. J. B. (1983). *Acta Cryst.* **A39**, 860–864.
- Petersen, J. L. (1980). *Inorg. Chem.* **19**, 181–185.
- Samuel, E., Rogers, R. D. & Atwood, J. L. (1984). *J. Crystallogr. Spectrosc. Res.* **14**, 573–579.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Skapski, A. C. & Troughton, P. G. H. (1970). *Acta Cryst.* **B26**, 716–722.
- Spek, A. L. (1990). *Acta Cryst.* **A46**, C-34.
- Thewalt, U. & Schomburg, D. (1977). *J. Organomet. Chem.* **127**, 169–174.

*Acta Cryst.* (1998). **C54**, 1612–1615

## A Trinuclear Molybdenum Cluster Coordinated by *o*-Nitrobenzoate

Ji-Bo XIA, Yuan-Gen YAO, Ling Wu, Xiao-Ying Huang and Jia-Xi Lu

State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fuzhou, Fujian 350002, People's Republic of China. E-mail: yyg@ms.fjirsm.ac.cn

(Received 28 November 1997; accepted 1 April 1998)


## Abstract

The title compound, tris(diethyl dithiophosphato)- $1\kappa^2S,S';2\kappa^2S,S';3\kappa^2S,S'$ - $\mu$ -*o*-nitrobenzoato- $2\kappa O:3\kappa O'$ -pyridine- $1\kappa N-\mu_3$ -thio-tri- $\mu$ -thio- $1:2\kappa^2S;1:3\kappa^2S;2:3\kappa^2S$ -triangulo-trimolybdenum( $3 Mo-Mo$ ),  $[Mo_3S_4(C_7H_4N-O_4)(C_4H_{10}PS_2)_3(C_5H_5N)]$ , was synthesized in a mixed solvent of acetonitrile and ethanol by a ligand-substitution reaction in which *o*-nitrobenzoic acid replaces the bridging diethyl dithiophosphate (DTP) ligand. The average Mo—Mo bond distance is 2.731 (2) Å. The substitution of *o*-nitrobenzoate for DTP shortens the Mo—Mo bond that it bridges to 2.693 (1) Å.

## Comment

In the course of our study on the incomplete cubane-type Mo—S(O) cluster compounds, one of the authors proposed that there is quasi-aromaticity in the puckered ring of the  $Mo_3S_3$  moiety (Chen *et al.*, 1990). We were interested in studying the electron delocalization

between the Mo cluster core ( $\text{Mo}_3\text{S}_4$ ) and the aromatic ligands; thus, the synthesis of the title compound, (I), was designed. The reaction of  $[\text{Mo}_3\text{S}_4(\text{DTP})_4(\text{H}_2\text{O})]$ , (II) (Lin *et al.*, 1986), with benzoic acid yielded a brown powder which proved difficult to dissolve in any solvent tried. Nevertheless, the reaction with *o*-nitrobenzoic acid and pyridine (py) gave good crystals of the title compound,  $[\text{Mo}_3\text{S}_4(\text{DTP})_3(2\text{-NO}_2\text{C}_6\text{H}_4\text{CO}_2)(\text{py})]$  (DTP is diethyl dithiophosphate), (I).



The main structural feature of compound (I) (Fig. 1) is similar to that of the many incomplete cubane-type cluster compounds with organic ligands (Shibahara, 1991). The dihedral angle calculated between the plane through atoms Mo2, Mo3, O1, O2, C1 and C2, and that through atoms C1–C7 and N1 is  $46.3(6)^\circ$ . There is a dihedral angle of  $31.7(3)^\circ$  between the plane of Mo1, Mo2 and Mo3, and the plane of Mo2, Mo3, O1, O2, C1 and C2, which reveals heavily distorted octahedral

Mo coordination. The substitution of bridging DTP by organic acids shortens the bridged Mo—Mo bond distance. The Mo2—Mo3 bond of the title compound is  $2.693(1)\text{ \AA}$ , which is shorter than the DTP-bridged Mo2—Mo3 bond of compound (II) [ $2.734(1)\text{ \AA}$ ; Lin *et al.*, 1986]. Furthermore, substitution by aromatic acids shortens the bridged Mo—Mo bond more than substitution by aliphatic acids (Huang *et al.*, 1988).

The  $^{31}\text{P}$  NMR spectra of compound (I) were measured in deuterated acetone (Yao *et al.*, 1995). Two peaks at 111.16 (assigned to the two DTPs coordinated to Mo2 and Mo3) and 110.20 p.p.m. (assigned to the DTP coordinated to Mo1) were observed with the intensity ratio 2:1.

Among the calculated intermolecular atomic distances, no  $\text{S}\cdots\text{S}$  interaction between two molecules in the cell was found. This contrasts with the analogous compounds  $[\text{Mo}_3\text{S}_4(\text{DTP})_3(\text{CH}_3\text{C}_6\text{H}_4\text{SO}_3)(\text{C}_2\text{H}_5\text{OH})]$  (Lu *et al.*, 1995) and  $[\text{Mo}_3\text{S}_4(\text{DTP})_3(2\text{-HOC}_6\text{H}_4\text{CO}_2)(\text{CH}_3\text{CN})]$  (Xia *et al.*, 1998), in which several intermolecular  $\text{S}\cdots\text{S}$  interactions ( $\text{S}\cdots\text{S} < 3.60\text{ \AA}$ ) are observed. The loosely coordinated pyridine ligand of each molecule occupies the space between two molecules and hinders  $\text{S}\cdots\text{S}$  interactions.

## Experimental

$[\text{Mo}_3\text{S}_4(\text{DTP})_4(\text{H}_2\text{O})]$  was synthesized according to the literature method of Yao *et al.* (1995); however, HDTP was purified before use by vacuum distillation (Bacon & LeSure, 1954; Olah & McFarland, 1975). In a flask,  $[\text{Mo}_3\text{S}_4(\text{DTP})_4(\text{H}_2\text{O})]$  (0.2 g) was dissolved in ethanol–acetonitrile (40 ml,  $v/v = 1/1$ ); *o*-nitrobenzoic acid (0.03 g) and a few drops of pyridine were then added. After refluxing over an oil bath at 363 K for 1 h, the hot dark-brown solution was filtered into another flask. Rectangular black crystals precipitated over a period of 10 d (yield: 0.12 g, 60%).

### Crystal data

$[\text{Mo}_3\text{S}_4(\text{C}_7\text{H}_4\text{NO}_4)(\text{C}_4\text{H}_{10}\text{PS}_2)_3(\text{C}_5\text{H}_5\text{N})]$

$M_r = 1216.97$

Triclinic

$P\bar{1}$

$a = 10.526(6)\text{ \AA}$

$b = 14.030(4)\text{ \AA}$

$c = 16.884(4)\text{ \AA}$

$\alpha = 93.16(2)^\circ$

$\beta = 103.19(3)^\circ$

$\gamma = 110.99(3)^\circ$

$V = 2241(3)\text{ \AA}^3$

$Z = 2$

$D_x = 1.80\text{ Mg m}^{-3}$

$D_m$  not measured

Mo  $K\alpha$  radiation

$\lambda = 0.71069\text{ \AA}$

Cell parameters from 20 reflections

$\theta = 10.50\text{--}14.95^\circ$

$\mu = 1.41\text{ mm}^{-1}$

$T = 293\text{ K}$

Cuboid

$0.80 \times 0.60 \times 0.40\text{ mm}$

Black

### Data collection

Enraf–Nonius CAD-4 diffractometer

6835 reflections with  $I > 3\sigma(I)$




Fig. 1. The molecular structure of (I) showing 30% probability displacement ellipsoids. H atoms have been omitted for clarity.

$\omega/2\theta$  scans

Absorption correction:

 $\psi$  scan (North *et al.*, 1968) $T_{\min} = 0.440$ ,  $T_{\max} = 0.569$ 

7864 measured reflections

7864 independent reflections

**Refinement**Refinement on  $F$  $R = 0.044$  $wR = 0.055$  $S = 1.80$ 

6835 reflections

470 parameters

H atoms not refined

 $w = 1/\sigma^2(F)$  $\theta_{\max} = 25^\circ$  $h = 0 \rightarrow 12$  $k = -16 \rightarrow 16$  $l = -20 \rightarrow 20$ 

3 standard reflections

every 60 reflections

intensity decay: 0.8%

 $(\Delta/\sigma)_{\max} = 0.006$  $\Delta\rho_{\max} = 1.08 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -1.09 \text{ e } \text{\AA}^{-3}$ 

Extinction correction: none

Scattering factors from *International Tables for X-ray Crystallography* (Vol. IV)

Mo1—Mo2—O1	133.7 (1)	Mo2—S1—Mo3	70.74 (5)
Mo3—Mo2—S1	54.52 (4)	Mo1—S2—Mo2	73.64 (5)
Mo3—Mo2—S2	99.61 (4)	Mo1—S3—Mo3	74.28 (5)
Mo3—Mo2—S4	53.94 (5)	Mo2—S4—Mo3	71.90 (5)
Mo3—Mo2—S7	139.15 (4)	Mo2—O1—C1	122.0 (3)
Mo3—Mo2—S8	138.85 (4)	Mo3—O2—C1	122.2 (2)
Mo3—Mo2—O1	83.68 (9)	O3—N1—O4	125.7 (6)
S1—Mo2—S2	105.75 (5)	O3—N1—C3	117.7 (5)
S1—Mo2—S4	107.88 (6)	O4—N1—C3	116.6 (6)
S1—Mo2—S7	158.95 (5)	Mo1—N2—C8	121.6 (3)
S1—Mo2—S8	85.23 (5)	Mo1—N2—C12	121.2 (4)
S1—Mo2—O1	81.7 (1)	C8—N2—C12	117.1 (4)
S2—Mo2—S4	93.38 (5)	O1—C1—O2	126.9 (4)
S2—Mo2—S7	88.76 (6)	O1—C1—C2	117.8 (4)
S2—Mo2—S8	99.38 (5)	O2—C1—C2	115.2 (4)
S2—Mo2—O1	172.4 (1)	C1—C2—C3	122.4 (4)
S4—Mo2—S7	85.89 (6)	N2—C8—C9	123.5 (5)
S4—Mo2—S8	158.56 (5)	N2—C12—C11	122.5 (6)

The positions of the Mo atoms were located by direct methods and those of the other atoms from successive difference Fourier syntheses. The highest peak on the final difference Fourier map had a height of  $1.076 \text{ e } \text{\AA}^{-3}$  and was located at a position  $1.04 \text{ \AA}$  from Mo1. The next highest peak was  $1.067 \text{ e } \text{\AA}^{-3}$ ,  $1.13 \text{ \AA}$  from Mo3.

Data collection: *CAD-4 Operations Manual* (Enraf-Nonius, 1977). Cell refinement: *CAD-4 Operations Manual*. Data reduction: *PROCESS* in *TEXSAN* (Molecular Structure Corporation, 1987). Program(s) used to solve structure: *MITHRIL* (Gilmore, 1983) in *TEXSAN*. Program(s) used to refine structure: *TEXSAN*. Molecular graphics: *ORTEPII* (Johnson, 1976). Software used to prepare material for publication: *GCIF* (local program).

We are grateful for financial support from the Foundation for National Natural Sciences of China (29303025) and the Foundation of the Chinese State Key Laboratory of Structural Chemistry (970095). The authors also wish to express their appreciation to Dr J. T. Chen and Mr Q. M. Wang for the data collection and structure analysis.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FR1109). Services for accessing these data are described at the back of the journal.

**References**

- Bacon, W. E. & LeSure, W. M. (1954). *J. Am. Chem. Soc.* **76**, 670–676.
- Chen, Z.-D., Li, J., Cheng, W.-D., Huang, J.-Q., Liu, C.-W., Zhang, Q.-E. & Lu, J.-X. (1990). *Chin. Sci. Bull.* **35**, 1698–1704.
- Enraf-Nonius (1977). *CAD-4 Operations Manual*. Enraf-Nonius, Delft, The Netherlands.
- Gilmore G. J. (1983). *MITHRIL. A Computer Program for the Automatic Solution of Crystal Structures from X-ray Data*. University of Glasgow, Scotland.
- Huang, J.-Q., Huang, J.-L., Shang, M.-Y., Lu, S.-F., Lin, X.-T., Lin, Y.-H., Huang, M.-D., Zhuang, H.-H. & Lu, J.-X. (1988). *Pure Appl. Chem.* **60**, 1185–1192.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Lin, X.-T., Lin, Y.-H., Huang, J.-L. & Huang, J.-Q. (1986). *Kexue Tongbao*, pp. 509–513.
- Lu, S.-F., Huang, J.-Q., Huang, X.-Y. & Wu, Q.-J. (1995). *Jiegou Huaxue (Chin. J. Struct. Chem.)*, **14**, 482–487.

Table 1. Selected geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Mo1—Mo2	2.745 (1)	Mo3—S9	2.512 (2)
Mo1—Mo3	2.756 (2)	Mo3—S10	2.558 (2)
Mo1—S1	2.335 (1)	Mo3—O2	2.201 (3)
Mo1—S2	2.291 (2)	O1—C1	1.266 (5)
Mo1—S3	2.283 (1)	O2—C1	1.263 (6)
Mo1—S5	2.568 (2)	O3—N1	1.213 (8)
Mo1—S6	2.539 (2)	O4—N1	1.202 (8)
Mo1—N2	2.368 (4)	N1—C3	1.474 (7)
Mo2—Mo3	2.693 (1)	N2—C8	1.334 (7)
Mo2—S1	2.329 (2)	N2—C12	1.345 (6)
Mo2—S2	2.288 (1)	C1—C2	1.487 (5)
Mo2—S4	2.297 (2)	C2—C3	1.384 (7)
Mo2—S7	2.555 (2)	C2—C7	1.383 (6)
Mo2—S8	2.514 (2)	C3—C4	1.374 (6)
Mo2—O1	2.239 (3)	C4—C5	1.359 (8)
Mo3—S1	2.323 (1)	C5—C6	1.37 (1)
Mo3—S3	2.282 (1)	C6—C7	1.373 (7)
Mo3—S4	2.290 (2)		
Mo2—Mo1—Mo3	58.63 (4)	S4—Mo2—O1	83.02 (9)
Mo2—Mo1—S1	53.85 (4)	S7—Mo2—S8	77.28 (6)
Mo2—Mo1—S2	53.12 (4)	S7—Mo2—O1	84.4 (1)
Mo2—Mo1—S3	98.37 (5)	S8—Mo2—O1	82.19 (9)
Mo2—Mo1—S5	139.77 (5)	Mo1—Mo3—Mo2	60.47 (3)
Mo2—Mo1—S6	140.91 (4)	Mo1—Mo3—S1	53.93 (4)
Mo2—Mo1—N2	91.9 (1)	Mo1—Mo3—S3	52.89 (4)
Mo3—Mo1—S1	53.52 (4)	Mo1—Mo3—S4	96.71 (5)
Mo3—Mo1—S2	97.74 (5)	Mo1—Mo3—S9	102.39 (5)
Mo3—Mo1—S3	52.83 (4)	Mo1—Mo3—S10	144.61 (4)
Mo3—Mo1—S5	140.04 (4)	Mo1—Mo3—O2	133.93 (8)
Mo3—Mo1—S6	102.10 (5)	Mo2—Mo3—S1	54.74 (4)
Mo3—Mo1—N2	136.71 (8)	Mo2—Mo3—S3	99.89 (4)
S1—Mo1—S2	105.44 (5)	Mo2—Mo3—S4	54.17 (4)
S1—Mo1—S3	104.39 (5)	Mo2—Mo3—S9	139.15 (4)
S1—Mo1—S5	161.46 (5)	Mo2—Mo3—S10	138.77 (5)
S1—Mo1—S6	87.10 (5)	Mo2—Mo3—O2	85.17 (9)
S1—Mo1—N2	83.89 (9)	S1—Mo3—S3	104.85 (5)
S2—Mo1—S3	98.58 (6)	S1—Mo3—S4	108.33 (6)
S2—Mo1—S5	86.65 (6)	S1—Mo3—S9	84.81 (5)
S2—Mo1—S6	160.16 (4)	S1—Mo3—S10	156.93 (5)
S2—Mo1—N2	85.4 (1)	S1—Mo3—O2	81.79 (9)
S3—Mo1—S5	87.21 (5)	S3—Mo3—S4	94.54 (6)
S3—Mo1—S6	92.92 (6)	S3—Mo3—S9	95.48 (6)
S3—Mo1—N2	169.36 (9)	S3—Mo3—S10	91.73 (6)
S5—Mo1—S6	77.77 (6)	S3—Mo3—O2	173.18 (8)
S5—Mo1—N2	83.14 (9)	S4—Mo3—S9	160.81 (4)
S6—Mo1—N2	80.7 (1)	S4—Mo3—S10	85.76 (6)
Mo1—Mo2—Mo3	60.90 (3)	S4—Mo3—O2	84.7 (1)
Mo1—Mo2—S1	54.06 (4)	S9—Mo3—S10	77.62 (6)
Mo1—Mo2—S2	53.23 (4)	S9—Mo3—O2	83.4 (1)
Mo1—Mo2—S4	96.86 (5)	S10—Mo3—O2	81.46 (9)
Mo1—Mo2—S7	141.96 (4)	Mo1—S1—Mo2	72.09 (5)
Mo1—Mo2—S8	104.58 (5)	Mo1—S1—Mo3	72.55 (5)

- Molecular Structure Corporation (1987). *TEXSAN. TEXRAY Structure Analysis Package*. Revised. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Olah, G. A. & McFarland, C. W. (1975). *J. Org. Chem.* **40**, 2582–2587.
- Shibahara, T. (1991). *Adv. Inorg. Chem.* **37**, 143–173.
- Xia, J.-B., Wu, L., Wang, Q.-M., Chen, J.-T., Wu, D., Yao, Y.-G. & Lu, J.-X. (1998). *Jiegou Huaxue (Chin. J. Struct. Chem.)*, **17**, 302–308.
- Yao, Y.-G., Akashi, H., Sakane, G., Shibahara, T. & Ohtaki, H. (1995). *Inorg. Chem.* **34**, 42–48.

*Acta Cryst.* (1998). **C54**, 1615–1616

### Hexapotassium $\mu$ -Oxo-bis[(citrate)-dioxomolybdenum] Dihydrate

YONG-HENG XING,<sup>a,b</sup> JI-QING XU,<sup>a,b</sup> HAO-RAN SUN,<sup>a</sup>  
DONG-MEI LI,<sup>a</sup> REN-ZHANG WANG,<sup>a</sup> TIE-GANG WANG,<sup>a</sup>  
WEI-MING BU,<sup>c</sup> LING YE,<sup>c</sup> GUANG-DI YANG<sup>c</sup> AND  
YU-GUO FAN<sup>c</sup>

<sup>a</sup>Department of Chemistry, Jilin University, Changchun, Jilin 130023, People's Republic of China, <sup>b</sup>State Key Laboratory of Coordination Chemistry, Nanjing University, Nanjing 210093, People's Republic of China, and <sup>c</sup>Key Laboratory for Supramolecular Structure and Spectroscopy, Jilin University, Changchun, Jilin 130023, People's Republic of China. E-mail: [people@mail.jlu.edu.cn](mailto:people@mail.jlu.edu.cn)

(Received 5 January 1998; accepted 5 May 1998)


#### Abstract

In the title compound, hexapotassium  $\mu$ -oxo-bis{dioxo[2-hydroxy- $\kappa$ O-1,2,3-propanetricarboxylato(4-)- $\kappa^2$ O<sup>1</sup>,O<sup>2</sup>]molybdenum(VI)} dihydrate,  $K_6\{[MoO_2(C_6H_4O_7)]_2\} \cdot 2H_2O$ , the citrate, as a tridentate ligand, is coordinated to molybdenum through the deprotonated hydroxy group, the  $\alpha$ -carboxylate group and also through one  $\beta$ -carboxylate group. The coordination polyhedron of the Mo atom is distorted octahedral, containing one  $[Mo_2O_5]^{2+}$  core. Principal dimensions are: Mo—O<sub>r</sub> = 1.715 (av), Mo—O<sub>b</sub> = 1.915 (2), Mo—O<sub>hydroxy</sub> = 1.952 (3), Mo—O <sub>$\alpha$ -carboxy</sub> = 2.216 (4), Mo—O <sub>$\beta$ -carboxy</sub> = 2.261 (4) Å and Mo—O<sub>b</sub>—Mo = 137.1 (4)°.

#### Comment

There is at present a growing interest in the chemistry of the coordination compounds of molybdenum,

particularly because of their possible relationship to redox-active molybdoenzymes (Chan *et al.*, 1993; Russ, 1996). However, structural studies on oxomolybdenum complexes containing a citrate ligand are rare. One complex,  $K_2Na_4[(MoO_2)_2O(cit)_2] \cdot 5H_2O$  (cit = citrate), has been structurally characterized by X-ray diffraction (Zhou *et al.*, 1997), and the title complex, (I), was considered to be isomorphous with this and to have the same coordination mode. Structure analysis shows that the dimeric anion of (I) consists of the common oxo-bridged  $[Mo_2O_5]^{2+}$  entity, which is non-centrosymmetric. Each Mo atom is six-coordinate,



with approximately octahedral geometry. The terminal and bridging oxo groups adopt a *fac* stereochemistry. Other coordination positions are occupied by a fully deprotonated tridentate citrate ligand, which forms both five-membered and six-membered chelate rings. The O1—Mo1—O5 and O1—Mo1—O6 bond angles are 73.86 (13) and 79.68 (14)°, respectively. A comparison of the title compound with two related compounds, namely,  $K_2Na_4[(MoO_2)_2O(cit)_2] \cdot 5H_2O$  (Zhou *et al.*, 1997) and  $Na_6[W_2O_5(cit)_2] \cdot 10H_2O$  (Llopis *et al.*, 1993), indicates that both reported structures, as well as the present one, have almost the same structural features. The corresponding bond distances and angles to *M* (*M* = Mo or W) are similar, except for the *M*—O <sub>$\beta$ -carboxy</sub> distance and the *M*—O<sub>b</sub>—*M* angle of the title compound being different to those of  $Na_6[W_2O_5(cit)_2] \cdot 10H_2O$  (Llopis *et al.*, 1993) {*M*—O <sub>$\beta$ -carboxy</sub> = 2.261 (4) Å and *M*—O<sub>b</sub>—*M* = 137.1 (4)° for (I), and 2.289 (2) Å and 180° for  $Na_6[W_2O_5(cit)_2] \cdot 10H_2O$ }.




Fig. 1. A view of the title complex showing displacement ellipsoids drawn at the 50% probability level and the atom-labelling scheme.