

Steiner, T. (1996). *Crystallogr. Rev.* **6**, 1–57.

Xu, X., You, X., Sun, Z., Wang, X. & Liu, H. (1994). *Acta Cryst.* **C50**, 1169–1171.

Acta Cryst. (1999). **C55**, 119–120

2-(2-Pyrrolyl)-1,3-benzothiazole

NAKA DAVIDOVIĆ,^{a†} DUBRAVKA MATKOVIĆ-ČALOGOVIĆ,^{a‡}
ZORA POPOVIĆ^a AND LELJA FIŠER-JAKIĆ^b

^aLaboratory of General and Inorganic Chemistry, Faculty of Science, University of Zagreb, Ul. kralja Zvonimira 8, HR-10000 Zagreb, Croatia, and ^bFaculty of Textile Technology, University of Zagreb, Pierottijeva 6, HR-10000 Zagreb, Croatia. E-mail: zpopovic@zagreb.zoak.pmf.hr


(Received 27 April 1998; accepted 14 September 1998)

Abstract

The title compound, C₁₁H₈N₂S, already known from the literature, was prepared and studied by X-ray diffraction. The five-membered heterocyclic ring is coplanar with its fused benzene ring. However, the molecule itself is non-planar, with the dihedral angle between the planes of the 2-benzothiazolyl and pyrrolyl groups being 17.3 (7)°. The molecules related by the twofold rotation axis are connected in pairs by N—H···N hydrogen bonds of 3.032 (9) Å.

Comment

Organic luminescent materials are widely used in the basic as well as applied research fields and various kinds of luminescent compounds are now known. Among this group of compounds are benzothiazole derivatives, whose importance as substances used as fluorescent brighteners (Allen, 1971) and for fluorometric measurements (Akiyama *et al.*, 1987) has led us to study the spectral characteristics of some aryl- and heteroaryl-substituted benzothiazoles (Tralić-Kulenović *et al.*, 1993). 2-(2-Pyrrolyl)benzothiazole (Brown, 1962) exhibited a relatively large fluorescence quantum efficiency and it was therefore of interest to find out more about the structure of this compound, (I).



† Temporary address: Martin-Luther-Universität Halle-Wittenberg, Institut für Anorganische Chemie (Merseburg), D-06099 Halle, Germany.

‡ Temporary address: Department of Organic Chemistry, University of Padua, Via F. Marzolo 1, I-35131 Padua, Italy.

The bond distances in both five-membered heterocyclic rings are consistent with those usually found in the literature (Potenza & Mastropaolo, 1974; Rudd & Barany, 1984). The benzene ring and its fused thiazolyl ring are coplanar. The bond S—C11 [1.739 (3) Å] is shorter than S—C5 [1.754 (3) Å], due to the fact that C5 is *sp*² hybridized, whereas C11 is part of the aromatic ring (Yeap *et al.*, 1991). Therefore, the molecular structure is not planar; the dihedral angle between the planes of the 2-benzothiazolyl and pyrrolyl groups is 17.3 (7)°. The deviation from coplanarity is significant; the torsion angles N1—C1—C5—N2, N1—C1—C5—S, C4—C1—C5—N2 and C4—C1—C5—S are 11.1 (4), -172.4 (2), -159.6 (3) and 16.9 (5)°, respectively.

The molecules, related by the twofold rotation axis, are connected in pairs by two hydrogen bonds [NH···N 3.032 (9), H···N 2.169 (7) Å and N—H···N 166.0 (7)°].




Fig. 1. View of the title molecule with the atom-labelling scheme for the non-H atoms. Displacement ellipsoids are drawn at the 50% probability level.

Experimental

The title compound was prepared from aminothiophenole and the required heteroaldehyde according to the method of Bogert & Stull (1935). Good quality crystals for X-ray diffraction analysis were obtained by slow evaporation from a dimethylformamide solution.

Crystal data

C₁₁H₈N₂S
M_r = 200.25

Tetragonal
P4₃2₁2

a = 8.0916 (9) Å

c = 30.078 (4) Å

V = 1969.3 (4) Å³

Z = 8

D_x = 1.351 Mg m⁻³

D_m = 1.3 Mg m⁻³

D_m measured by flotation in benzene/CCl₄

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 32 reflections

θ = 7.8–13.1°

μ = 0.285 mm⁻¹

T = 293 (2) K

Bipyramid

0.135 × 0.120 × 0.113 mm

Bronze

Data collection

Philips PW1100 updated by Stoe diffractometer

*R*_{int} = 0.050

θ_{max} = 28.99°

ω -2 θ scans
Absorption correction: none
5310 measured reflections
2791 independent reflections
1857 reflections with
 $I > 2\sigma(I)$

$h = -2 \rightarrow 11$
 $k = 0 \rightarrow 11$
 $l = 0 \rightarrow 41$
4 standard reflections
frequency: 120 min
intensity decay: 3.5%

Refinement

Refinement on F^2
 $R(F) = 0.055$
 $wR(F^2) = 0.087$
 $S = 0.899$
2590 reflections
159 parameters
All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0247P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.131 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.152 \text{ e } \text{Å}^{-3}$
Extinction correction: none
Scattering factors from
International Tables for Crystallography (Vol. C)
Absolute structure: Flack (1983)
Flack parameter = 0.12 (12)

Table 1. Selected geometric parameters (Å, °)

S—C11	1.739 (3)	C2—C3	1.354 (5)
S—C5	1.754 (3)	C3—C4	1.396 (5)
N1—C2	1.346 (4)	C6—C7	1.383 (4)
N1—C1	1.374 (3)	C6—C11	1.390 (4)
N2—C5	1.299 (3)	C7—C8	1.375 (5)
N2—C6	1.398 (3)	C8—C9	1.395 (5)
C1—C4	1.370 (4)	C9—C10	1.366 (5)
C1—C5	1.443 (4)	C10—C11	1.387 (4)
C11—S—C5	89.1 (2)	C1—C5—S	119.5 (2)
C2—N1—C1	109.0 (3)	C7—C6—C11	119.7 (3)
C5—N2—C6	110.6 (3)	C7—C6—N2	124.9 (3)
C4—C1—N1	107.1 (3)	C11—C6—N2	115.3 (3)
C4—C1—C5	131.5 (3)	C8—C7—C6	119.2 (4)
N1—C1—C5	120.9 (3)	C7—C8—C9	120.5 (4)
N1—C2—C3	109.0 (3)	C10—C9—C8	120.8 (4)
C2—C3—C4	107.2 (4)	C9—C10—C11	118.6 (4)
C1—C4—C3	107.7 (4)	C10—C11—C6	121.1 (3)
N2—C5—C1	124.9 (3)	C10—C11—S	129.4 (3)
N2—C5—S	115.5 (2)	C6—C11—S	109.4 (3)
C4—C1—C5—N2	-159.6 (3)	C4—C1—C5—S	16.9 (5)
N1—C1—C5—N2	11.1 (4)	N1—C1—C5—S	-172.4 (2)

Table 2. Hydrogen-bonding geometry (Å, °)

D—H...A	D—H	H...A	D...A	D—H...A
N1—H1...N2 ⁱ	0.88 (2)	2.169 (7)	3.032 (9)	166.0 (7)

Symmetry code: (i) 1 + y, x - 1, 2 - z.

Intensities were corrected for Lorentz and polarization effects. The structure was solved by direct methods. The H atoms were found in the difference Fourier map and were refined isotropically. Final full-matrix least-squares refinement of the coordinates, anisotropic displacement parameters for non-H atoms and isotropic displacement parameters for H atoms reduced R to 0.055. The absolute structure of the title compound was unequivocally established by refining the Flack chirality parameter (Flack, 1983).

Data collection: *STAD14* (Stoe & Cie, 1995). Cell refinement: *STAD14*. Data reduction: *X-RED* (Stoe & Cie, 1995). Program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997a). Program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997b). Molecular graphics: *PLATON97* (Spek, 1990) and *ORTEPII* (Johnson, 1976). Software used to prepare material for publication: *SHELXL97* and *PLATON97*.

The authors thank the Ministry of Science and Technology of the Republic of Croatia for financial support (grant No. 119408).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: KA1289). Services for accessing these data are described at the back of the journal.

References

- Akiyama, S., Ochiai, T., Nakatsuji, S., Nakashima, K. & Ohkura, Y. (1987). *Chem. Pharm. Bull.* **35**, 3029–3032.
- Allen, R. L. M. (1971). *Colour Chemistry*, pp. 278–286. London: Thomas Nelson & Sons Ltd.
- Bogert, M. T. & Stull, H. (1935). *J. Am. Chem. Soc.* **57**, 1529–1533.
- Brown, H. D. (1962). Belg. Patent Appl. 614 237 (22 August 1962); US Patent Appl. (1 March 1961); *Chem. Abstr.* (1963), No. 14666b.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Potenza, J. & Mastropaolo, D. (1974). *Acta Cryst.* **B30**, 2353–2359.
- Rudd, S. & Barany, G. (1984). *Acta Cryst.* **C40**, 2118–2120.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1997a). *SHELXS97. Program for the Solution of Crystal Structures*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997b). *SHELXL97. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Spek, A. L. (1990). *Acta Cryst.* **A46**, C-34.
- Stoe & Cie (1995). *STAD14. Diffractometer Control Program for Windows*. Stoe & Cie, Darmstadt, Germany.
- Stoe & Cie (1995). *X-RED. Data Reduction Program for Windows*. Stoe & Cie, Darmstadt, Germany.
- Tralić-Kulenović, V., Fišer-Jakić, L. & Lazarević, Z. (1993). *Spectrosc. Lett.* **26**, 1771–1784.
- Yeap, G.-Y., Fun, H.-K., Teo, S.-B. & Teoh, S.-G. (1991). *Acta Cryst.* **C47**, 1347–1348.

Acta Cryst. (1999). **C55**, 120–123

Diastereomeric complexes of 1,1'-bi-naphthyl-2,2'-diol and (R,R)-1,2-cyclohexanediamine

SHIGERU FUKUSHIMA,^a HIROYUKI HOSOMI,^a SHIGERU OHBA^a AND MASATOSHI KAWASHIMA^b

^aDepartment of Chemistry, Faculty of Science and Technology, Keio University, Hiyoshi 3-14-1, Kohoku-ku, Yokohama 223-8522, Japan, and ^bKankyo Kagaku Center Co. Ltd, Ookawa 5-1, Kanazawa-ku, Yokohama 236, Japan. E-mail: ohba@chem.keio.ac.jp

(Received 18 June 1997; accepted 15 January 1998)

Abstract

The structures of the molecular complexes (R)-1,1'-bi-naphthyl-2,2'-diol-(R,R)-1,2-cyclohexanediamine-toluene (1/1/1), C₂₀H₁₄O₂·C₆H₁₄N₂·C₇H₈, (I), and (S)-