

***trans*-Chloro(2-nitrobenzenethiolato-*S*)bis(triphenylphosphine-*P*)-palladium(II) monoacetone solvate**John H. Aupers,^a George Ferguson,^{b†} Christopher Glidewell,^{c*} John N. Low^c and James L. Wardell^d

^aRose Cottage, Middle Assendon, Henley-on-Thames, Oxon RG9 6AU, England, ^bSchool of Chemistry, University of St Andrews, St Andrews, Fife KY16 9ST, Scotland, ^cDepartment of Applied Physics and Electronic & Mechanical Engineering, University of Dundee, Nethergate, Dundee DD1 4HN, Scotland, and ^dDepartment of Chemistry, University of Aberdeen, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland
Correspondence e-mail: cg@st-andrews.ac.uk

Received 4 May 2000
Accepted 15 May 2000


Molecules of the title compound, [PdCl(C₆H₄NO₂S)(PPh₃)₂]-C₃H₆O, exhibit a slight distortion from exact planarity at the Pd atom towards tetrahedral, with P–Pd–P and Cl–Pd–S angles of 174.98 (3) and 174.19 (3)°, respectively. The Pd–Cl and Pd–S bonds are, respectively, long [2.3550 (11) Å] and short [2.3020 (12) Å] for their types; the S–C bond is also very short [1.744 (4) Å]. The solvating acetone molecule is linked to one of the phosphine ligands by means of a C–H···O hydrogen bond.

Comment

We have recently reported the structures of members of the 2-O₂NC₆H₄SX system containing the 2-nitrobenzenethiolate group, in which the α -atom of fragment X can be C or N (Low, Storey *et al.*, 2000) or S (Low, Glidewell & Wardell, 2000). In general, at the global energy minimum conformation, both the nitro group and the α -atom of group X are essentially coplanar with the aryl ring. When X is not in the plane of the aryl ring, rotation about the exocyclic C–S bond is associated with a disrotatory twist of the nitro group about the C–N bond; the occurrence of such conformations seems to depend on the presence in the structure of specific intermolecular interactions, such as C–H···O hydrogen bonds. Seeking to investigate the wider generality of this idea, we report here the structure of a further compound of type 2-O₂NC₆H₄SX, namely *trans*-chloro(2-nitrobenzenethiolato-*S*)bis(triphenylphosphine-*P*)palladium(II), in which the α -atom of group X is Pd and which crystallizes from acetone as a 1:1 solvate, (I).

Molecules of the Pd complex (I) lie in general positions (Fig. 1). Although no bond angle at Pd involving a pair of *cis* substituents deviates from 90° by more than 1.7°, nonetheless

there is a slight but significant distortion from planarity towards tetrahedral, and both bond angles at Pd involving *trans* pairs of substituents are less than 175° (Table 1). The deviations (Å) of the ligating atoms from their best plane through Pd are: Cl –0.106 (1), S –0.105 (1), P1 0.105 (1) and



P2 0.106 (1). In this respect, the configuration of the *trans*-PdClP₂S chromophore in (I) differs from those found in two analogous compounds retrieved from the Cambridge Structural Database (Allen & Kennard, 1993) containing *trans*-PdCl₂P₂ and *trans*-PdP₂S₂ chromophores. Molecules of *trans*-[PdCl₂(PPh₃)₂] are centrosymmetric both in the unsolvated compound (BISKIX; Ferguson *et al.*, 1982) and in the bischloroform solvate (RAVYUI; Stark & Whitmire, 1997). Similarly, molecules of *trans*-[Pd{PH(C₆H₁₁)₂}(SPh)₂] (LETNON; Pasquali *et al.*, 1993) lie across inversion centres. Hence, in all these complexes, palladium and its four ligating atoms are strictly coplanar.

The conformation of (I) is close to that expected (Low, Storey *et al.*, 2000) for the global energy minimum. There is a slight rotation around the C–S bond away from coplanarity of the Pd atom with the nitrated aryl ring, and a disrotatory twist of the nitro group around the C–N bond. The dihedral angles between the nitrated aryl ring and the planes defined by Cl–S1–Pd1–Cl1 and C6–NO₂ are 6.2 (2) and 16.3 (2)°, respectively. In centrosymmetric molecules of *trans*-[PdCl₂(PPh₃)₂] (Ferguson *et al.*, 1982; Stark & Whitmire, 1997), the P–C bonds of the two phosphine ligands are necessarily exactly staggered. In contrast, in (I), these P–C bonds are almost eclipsed, with a mean deviation of 5.9 (3)° from a fully eclipsed Ph₃P–Pd–PPh₃ fragment. Presumably, the rotational barriers about the Pd–P bonds are low.

The two independent Pd–P distances in (I) (Table 1) are both very similar to those reported for BISKIX [2.337 (2) Å] and RAVYUI [2.343 (2) Å]. However, the Pd–Cl distance in (I) is very much longer than the corresponding distances in BISKIX [2.291 (2) Å] and RAVYUI [2.293 (2) Å]. On the other hand, the Pd–S distance in (I) is much shorter than those reported for LETNON [2.3393 (16) and 2.3366 (19) Å in two independent molecules]. These observations can all be readily rationalized in terms of the *trans* influence, *t*(X), of a ligand X upon the properties of the metal–ligand bond *trans* to X, which arises from the competition for a common metal *d* orbital of the π type between ligands occupying *trans* sites. Ligands of high *trans*-influence render the metal–ligand bonds *trans* to themselves longer (Appleton *et al.*, 1973), and ligands can thus be ranked in order of *t*(X). This is a static thermodynamic phenomenon which closely parallels the dynamic kinetic *trans* effect describing the influence on the rates of ligand substitution exerted by ligands in *trans* sites (Basolo &

† GF was on leave from the Department of Chemistry and Biochemistry, University of Guelph, Guelph, Ontario, Canada N1G 2W1.

Pearson, 1962). The data for (I) and related compounds show that a Pd–S bond *trans* to Cl is much shorter than a Pd–S bond *trans* to another thiolate; hence, $t(SR^-) \gg t(Cl^-)$. Similarly, Pd–P bonds *trans* to another P are typically much shorter than Pd–P bonds *trans* to chloride. For example, in *cis*-[PdCl₂(Me₂N=CH)PPh₃] (McCrindle *et al.*, 2000), the Pd–P distance is 2.2495 (7) Å, much shorter than the corresponding bonds in (I), BISKIX and RAVYUI. Likewise, in [PdCl₂(Me₂N=CH)PPh₃], the Pd–Cl bond *trans* to P is very much longer [2.3600 (7) Å] than those in BISKIX [2.291 (2) Å] and RAVYUI [2.293 (2) Å], so that $t(PR_3) \gg t(Cl^-)$. The order of PR_3 and SR^- can be established by comparison of analogous *cis*- and *trans*-PdP₂S₂ chromophores, as for example in LETNON and [Pd(Ph₂PCH₂CH₂PPh₂)-(SCH₂Ph)₂] (TERREN; Su *et al.*, 1997). On moving from *trans* to *cis* geometry, the Pd–P bonds become shorter and the Pd–S bonds become longer, hence $t(PR_3) > t(SR^-)$ and overall $t(PR_3) > t(SR^-) > t(Cl^-)$.




Figure 1
The asymmetric unit of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms bonded to C7 have a site-occupation factor of 0.5.

There is evidence of C–C bond fixation in the nitrated aryl ring, where the C–C distances range from 1.358 (6) to 1.414 (5) Å. This range is greater than is usual for compounds of this type (Low, Storey *et al.*, 2000; Low, Glidewell & Wardell 2000), although the C–N distance and the O–N–O angle are both entirely typical of their types, so ruling out any *p*-quinonoid-type bond fixation, as observed in 2-O₂N-C₆H₄SCH=CHPh (Low, Storey *et al.*, 2000). On the other hand, the C–S distance is significantly shorter than those observed in previous examples of 2-O₂NC₆H₄SX compounds. For comparison, the lower quartile value for a C(aryl)–S–C bond distance is 1.765 Å and, indeed, C(aryl)–S bonds as short as that in (I) are generally found only with four-coordinate S^{VI} (Allen *et al.*, 1987). The short C–S bond and the extremely small C–C–C angles *ipso* to S are together consistent with significant electron donation from S into the adjacent ring (Domenicano & Murray-Rust, 1979).

The solvating acetone molecule is linked to a phenyl group in one of the phosphine ligands of (I) by a C–H···O hydrogen bond (Table 2).

Experimental

The title compound was obtained from the reaction of *trans*-[PdCl₂(PPh₃)₂] and 2-O₂NC₆H₄SSnPh₃ in acetone solution. The compound was purified by column chromatography on silica using CHCl₃ as eluent. The compound obtained decomposed on heating, before melting at 478 K. Crystals suitable for single-crystal X-ray diffraction were obtained from an acetone solution.

Crystal data

[PdCl(C ₆ H ₄ NO ₂ S)(C ₁₈ H ₁₅ P) ₂] C ₃ H ₆ O	$D_x = 1.410 \text{ Mg m}^{-3}$
$M_r = 878.63$	Mo $K\alpha$ radiation
Monoclinic, $P2_1/n$	Cell parameters from 11315 reflections
$a = 18.292 (7) \text{ \AA}$	$\theta = 1.31\text{--}30.48^\circ$
$b = 10.969 (3) \text{ \AA}$	$\mu = 0.681 \text{ mm}^{-1}$
$c = 21.363 (8) \text{ \AA}$	$T = 150 (1) \text{ K}$
$\beta = 105.0928 (17)^\circ$	Block, orange
$V = 4139 (2) \text{ \AA}^3$	$0.30 \times 0.10 \times 0.05 \text{ mm}$
$Z = 4$	

Data collection

Enraf–Nonius diffractometer with FAST area detector	11 315 independent reflections
φ and ω scans with κ offsets	5642 reflections with $I > 2\sigma(I)$
Absorption correction: multi-scan (FAST/MADNES; Pflugrath & Messerschmidt, 1989)	$R_{\text{int}} = 0.095$
$T_{\text{min}} = 0.822$, $T_{\text{max}} = 0.967$	$\theta_{\text{max}} = 30.05^\circ$
49 325 measured reflections	$h = -25 \rightarrow 25$
	$k = -14 \rightarrow 15$
	$l = -29 \rightarrow 30$
	Intensity decay: negligible

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.0686P)^2]$
$R[F^2 > 2\sigma(F^2)] = 0.053$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.147$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 0.943$	$\Delta\rho_{\text{max}} = 1.54 \text{ e \AA}^{-3}$
11 315 reflections	$\Delta\rho_{\text{min}} = -1.10 \text{ e \AA}^{-3}$
489 parameters	Extinction correction: SHELXL97
H-atom parameters constrained	Extinction coefficient: 0.0026 (2)

Table 1

Selected geometric parameters (Å, °).

Pd1–S1	2.3020 (11)	S1–C1	1.744 (4)
Pd1–P1	2.3366 (13)	O1–N1	1.228 (4)
Pd1–P2	2.3303 (12)	O2–N1	1.237 (4)
Pd1–Cl1	2.3550 (11)	N1–C6	1.458 (5)
Cl1–Pd1–P1	89.56 (4)	O1–N1–O2	122.4 (4)
Cl1–Pd1–P2	88.59 (4)	C2–C1–C6	113.8 (4)
S1–Pd1–P1	90.63 (4)	C2–C1–S1	121.2 (3)
S1–Pd1–P2	91.69 (4)	C6–C1–S1	125.0 (3)
Cl1–Pd1–S1	174.19 (3)	C1–C6–C5	122.7 (4)
P1–Pd1–P2	174.98 (3)	C1–C6–N1	121.3 (4)
C1–S1–Pd1	105.52 (14)	C5–C6–N1	116.0 (4)
Cl1–Pd1–P1–C111	–165.77 (15)	Cl1–Pd1–P2–C211	51.41 (12)
Cl1–Pd1–P1–C121	–45.54 (13)	Cl1–Pd1–P2–C221	171.16 (14)
Cl1–Pd1–P1–C131	74.17 (13)	Cl1–Pd1–P2–C231	–67.66 (13)

Table 2

Hydrogen-bonding geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C236–H236···O3	0.95	2.51	3.307 (6)	141

Compound (I) crystallized in the monoclinic system as a mono-acetone solvate; space group $P2_1/n$ was assumed from the systematic absences. H atoms were treated as riding atoms with a C–H distance of 0.95 Å for aromatic H atoms and 0.98 Å for methyl H atoms. The acetone molecule was modelled with six half-occupancy H atoms in the methyl group defined by C7; the O and C atoms of the acetone molecule all had large displacement parameters. Examination of the refined structure with *PLATON* (Spek, 2000) showed there were no solvent-accessible voids in the structure, but that there were four symmetry-related voids, each with a volume of *ca* 11 Å³, in the vicinity of the four solvent molecules in the unit cell. These may be associated with the large displacement parameters of the solvent molecule. The sites of maximum and minimum residual density are adjacent to the acetone and Pd atom, respectively.

Data collection: *CAD-4-PC Software* (Enraf–Nonius, 1992) and *FAST/MADNES* (Pflugrath & Messerschmidt, 1989); cell refinement: *CAD-4-PC Software* and *FAST/MADNES*; data reduction: *CAD-4-PC Software* and *FAST/MADNES*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *NRCVAX96* (Gabe *et al.*, 1989) and *SHELXL97* (Sheldrick, 1997); molecular graphics: *NRCVAX96*, *ORTEPII* (Johnson, 1976) and *PLATON* (Spek, 2000); software used to prepare material for publication: *NRCVAX96*, *SHELXL97* and *WordPerfect* macro *PREP8* (Ferguson, 1998).

X-ray data were collected at the EPSRC National X-ray Crystallography Service. We thank Professor M. Hursthouse and his staff for their assistance.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK1393). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H. & Kennard, O. (1993). *Chem. Des. Autom. News*, **8**, 31–37.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Appleton, T. G., Clark, H. C. & Manzer, L. E. (1973). *Coord. Chem. Rev.* **10**, 335–422.
- Basolo, F. & Pearson, R. G. (1962). *Prog. Inorg. Chem.* **4**, 381–453.
- Domenicano, A. & Murray-Rust, P. (1979). *Tetrahedron Lett.* pp. 2283–2286.
- Enraf–Nonius (1992). *CAD-4-PC Software*. Version 1.1. Enraf–Nonius, Delft, The Netherlands.
- Ferguson, G. (1998). *PREP8*. University of Guelph, Canada.
- Ferguson, G., McCrindle, R., McAlees, A. J. & Parvez, M. (1982). *Acta Cryst.* **B38**, 2679–2681.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Low, J. N., Glidewell, C. & Wardell, J. L. (2000). *Acta Cryst.* **C56**, 689–691.
- Low, J. N., Storey, E. J., McCarron, M., Wardell, J. L., Ferguson, G. & Glidewell, C. (2000). *Acta Cryst.* **B56**, 58–67.
- McCrindle, R., McAlees, A. J., Zang, E. & Ferguson, G. (2000). *Acta Cryst.* **C56**, e132–133.
- Pasquali, M., Marchetti, F., Leoni, P., Beringhelli, T. & D'Alfonso, G. (1993). *Gazz. Chim. Ital.* **123**, 659–664.
- Pflugrath, J. W. & Messerschmidt, A. (1989). *MADNES*. Version II. Delft Instruments, The Netherlands.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2000). *PLATON*. Version of March 2000. University of Utrecht, The Netherlands.
- Stark, J. L. & Whitmire, K. H. (1997). *Acta Cryst.* **C53**, IUC9700007.
- Su, W., Hong, M., Cao, R. & Liu, H. (1997). *Acta Cryst.* **C53**, 66–67.