

trans-Di- μ -bromo-bis[bromo(triethylphosphine- κ P)platinum(II)]

Stéphanie M. M. Cornet, Keith. B. Dillon, Andrés E. Goeta and Amber L. Thompson*

Department of Chemistry, University of Durham, South Road, Durham DH1 3LE, England

Correspondence e-mail: a.l.thompson@durham.ac.uk

Received 1 November 2004

Accepted 9 December 2004


Online 15 January 2005

The title compound, $[\text{Pt}_2\text{Br}_4(\text{C}_6\text{H}_{15}\text{P})_2]$, is a centrosymmetric dinuclear platinum(II) complex consisting of two square-planar platinum centres connected by two bridging Br atoms.

Comment

Bridged chloride complexes of the form $[\text{PtCl}_2(\text{PR}_3)]_2$ have been used extensively as starting materials in the synthesis of mononuclear platinum–phosphine complexes, which are formed through cleavage of the bridging Pt–Cl bond (Chatt & Venanzi, 1955; Meidine & Nixon, 1988; Dillon & Goodwin, 1992, 1994). Similar synthetic methodology can also be applied to bromide analogues (Cornet, 2002). However, while the crystal structures of the chloro-bridged complexes $[\text{PtCl}_2(\text{PMe}_3)]_2$, $[\text{PtCl}_2(\text{PEt}_3)]_2$ and $[\text{PtCl}_2(\text{PPr}_3)]_2$ are known (Boag & Ravetz, 1996; Blake *et al.*, 1989; Black *et al.*, 1969), structures of complexes exhibiting the central heavy-atom skeleton $[\text{PtCl}_2\text{P}]_2$ are surprisingly rare, with only a handful known (Simms *et al.*, 1987; Copley *et al.*, 2000).

The title complex, (I), was prepared by the CHCl_2 -mediated reaction of equimolar quantities of PtBr_2 and $[\text{PtBr}_2(\text{PEt}_3)]_2$. Given the rarity of crystal structures containing the $[\text{PtCl}_2\text{P}]_2$ fragment, it is unsurprising that (I) is the first structure to be reported containing the $[\text{PtBr}_2\text{P}]_2$ motif.



The chloride complexes $[\text{PtCl}_2(\text{PR}_3)]_2$ (where $R = \text{CH}_3$, C_2H_5 and C_3H_7) and the title compound are closely related, and all four complexes possess an inversion centre in the middle of the dimer, with the PR_3 ligands in a *trans* geometry (Fig. 1). In addition, all four structures are asymmetric around the bridging halide ligands, but this asymmetry is reduced in (I) with respect to the chloride complexes (Table 1). This degree of asymmetry in (I) is presumably due to the relative positions of the Cl, Br and P atoms in the *trans* influence series and the increased ionic radius of the bromide ligand. These




Figure 1

A view of (I), with selected atoms labelled. Symmetry equivalents related by $(\frac{1}{2} - x, \frac{1}{2} - y, 1 - z)$ are also shown and are indicated by primes. Displacement ellipsoids for the non-H atoms are drawn at the 50% probability level.

factors also appear to reduce the effect bridging has on the bond lengths, since the bonds to the bridging Br atoms are only 0.023 and 0.122 Å longer than those to the terminal Br atoms, compared with averages of 0.033 and 0.146 Å in $[\text{PtCl}_2(\text{PR}_3)]_2$ (where $R = \text{CH}_3$, C_2H_5 and C_3H_7 ; Boag & Ravetz, 1996; Blake *et al.*, 1989; Black *et al.*, 1969, respectively).

Experimental

PtBr_2 (1.48 g, 4.2 mmol) in PhCN (10 ml) was heated to 373 K to give a bright-orange solution and a yellow precipitate on cooling [*cis*- $[\text{PtBr}_2(\text{PhCN})_2]$, yield 81%]. PEt_3 (1.75 g, 2.18 ml, 14.8 mmol) was then added to a solution of $[\text{PtBr}_2(\text{PhCN})_2]$ (1.77 g, 3.15 mmol) in CH_2Cl_2 (15 ml) and the mixture stirred for 3 h. Evaporation of the solvent produced a white solid [*cis*- $[\text{PtBr}_2(\text{PEt}_3)]_2$, yield 83%], some of which (1.45 g, 2.45 mmol) was added to a solution of PtBr_2 (1.03 g, 2.9 mmol) in $(\text{CHCl}_2)_2$ and heated to 423 K for 4 h. The yellow crystals of (I) obtained on cooling were recrystallized from CH_2Cl_2 (yield 79%). Analysis calculated for $\text{C}_{12}\text{H}_{30}\text{Br}_4\text{P}_2\text{Pt}_2$: C 15.23, H 3.20%; found: C 15.27, H 3.23%. ^{31}P NMR (CDCl_3): δ 10.9 (singlet with Pt satellites, $^1J_{\text{P-Pt}} = 3701$ Hz, $^3J_{\text{P-Pt}} = 24.4$ Hz, $^4J_{\text{P-P}} = 1.6$ Hz). The AA'XX' part of the spectrum was insufficiently resolved for $^2J_{\text{Pt-Pt}}$ to be evaluated (Kiffen *et al.*, 1975).

Crystal data

$[\text{Pt}_2\text{Br}_4(\text{C}_6\text{H}_{15}\text{P})_2]$
 $M_r = 946.12$
 Monoclinic, $C2/c$
 $a = 26.522$ (6) Å
 $b = 6.8720$ (13) Å
 $c = 13.811$ (4) Å
 $\beta = 120.930$ (7)°
 $V = 2159.3$ (9) Å³
 $Z = 4$

$D_x = 2.910$ Mg m⁻³
 Mo $K\alpha$ radiation
 Cell parameters from 7378 reflections
 $\theta = 1.9$ – 30.6 °
 $\mu = 20.48$ mm⁻¹
 $T = 120$ (2) K
 Block, clear intense orange
 0.20 × 0.10 × 0.10 mm

Data collection

Bruker SMART CCD 1K area-detector diffractometer
 ω scans
 Absorption correction: multi-scan (SADABS; Bruker, 1998)
 $T_{\text{min}} = 0.058$, $T_{\text{max}} = 0.129$
 10 540 measured reflections

2355 independent reflections
 2158 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\text{max}} = 27.0$ °
 $h = -32 \rightarrow 33$
 $k = -8 \rightarrow 8$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.020$ $wR(F^2) = 0.046$ $S = 1.11$

2355 reflections

94 parameters

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0168P)^2$$

$$+ 10.3265P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.85 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -1.75 \text{ e } \text{\AA}^{-3}$$

graphics: *SHELXTL* (Sheldrick, 1997b); software used to prepare material for publication: *SHELXTL*.

The authors thank the EPSRC for postgraduate studentships (SMMC and ALT) and Johnson–Matthey plc for the loan of platinum compounds.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: HJ1034). Services for accessing these data are described at the back of the journal.

Table 1

Selected geometric parameters (\AA , $^\circ$) for (I) and the related chloro complexes $[\text{PtCl}_2(\text{PMe}_3)_2]$, $[\text{PtCl}_2(\text{PEt}_3)_2]$ and $[\text{PtCl}_2(\text{PPr}_3)_2]$ (Hal = Cl or Br).

	$[\text{PtCl}_2(\text{PMe}_3)_2]$	$[\text{PtCl}_2(\text{PEt}_3)_2]$	$[\text{PtCl}_2(\text{PPr}_3)_2]$	(I)
Pt–P	2.205 (3)	2.212 (3)	2.230 (9)	2.2266 (11)
Pt–Hal _{terminal}	2.281 (3)	2.282 (3)	2.279 (9)	2.4229 (7)
Pt–Hal _{bridging}	2.309 (3)	2.318 (3)	2.315 (8)	2.4455 (7)
	2.423 (3)	2.431 (3)	2.425 (8)	2.5451 (6)
Pt–Hal–Pt	96.19 (10)	96.48 (10)	96.4 (3)	96.17 (2)
Hal–Pt–Hal	83.81 (10)	83.52 (9)	83.6 (2)	83.83 (2)

All H atoms were placed geometrically and refined using a riding model ($\text{C–H} = 0.98$ and 0.99 \AA), with their $U_{\text{iso}}(\text{H})$ values fixed at 1.2 or 1.5 times U_{eq} of the parent C atom. Although there are difference density holes larger than $1 \text{ e } \text{\AA}^{-3}$, they are within 1 \AA of the Pt atom.

Data collection: *SMART-NT* (Bruker, 1998); cell refinement: *SMART-NT*; data reduction: *SAINT-NT* (Bruker, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997a); program(s) used to refine structure: *SHELXS97* (Sheldrick, 1997a); molecular

References

- Black, M., Mais, R. H. B. & Owston, P. G. (1969). *Acta Cryst.* **B25**, 1760–1766.
- Blake, A. J., Gould, R. O., Marr, A. M., Rankin, D. W. H. & Schröder, M. (1989). *Acta Cryst.* **C45**, 1218–1219.
- Boag, N. M. & Ravetz, M. S. (1996). *Acta Cryst.* **C52**, 1942–1943.
- Bruker (1998). *SMART-NT*, *SAINT-NT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chatt, J. & Venanzi, L. M. (1955). *J. Chem. Soc.* pp. 2787–2793.
- Cobley, C. J., Ellis, D. D., Orpen, A. G. & Pringle, P. G. (2000). *J. Chem. Soc. Dalton Trans.* pp. 1101–1107.
- Cornet, S. M. M. (2002). PhD thesis, University of Durham, England.
- Dillon, K. B. & Goodwin, H. P. (1992). *J. Organomet. Chem.* **428**, 169–171.
- Dillon, K. B. & Goodwin, H. P. (1994). *J. Organomet. Chem.* **469**, 125–128.
- Kiffen, A. A., Masters, C. & Visser, J. P. (1975). *J. Chem. Soc. Dalton Trans.* pp. 1311–1315.
- Meidine, M. F. & Nixon, J. F. (1988). Private communication to K. B. Dillon.
- Sheldrick, G. M. (1997a). *SHELXL97* and *SHELXS97*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997b). *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Simms, B. L., Shang, M., Lu, J., Youngs, W. J. & Ibers, J. A. (1987). *Organometallics*, **6**, 1118–1126.