

Hydrogen bonding in C-substituted nitroanilines: simple C(8) chains in 2-bromo-6-chloro-4-nitroaniline

Christopher Glidewell,^{a*} John N. Low,^b Janet M. S. Skakle^b and James L. Wardell^c

^aSchool of Chemistry, University of St Andrews, Fife KY16 9ST, Scotland,

^bDepartment of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland, and ^cInstituto de Química, Departamento de Química Inorgânica, Universidade Federal do Rio de Janeiro, 21945-970 Rio de Janeiro, RJ, Brazil

Correspondence e-mail: cg@st-andrews.ac.uk

Received 29 March 2005

Accepted 30 March 2005

Online 30 April 2005

In the title compound, C₆H₄BrClN₂O₂, the Br and Cl substituents are disordered over two sites, and the intramolecular dimensions provide evidence for charge polarization. The molecules are linked into C(8) chains by a single N—H···O hydrogen bond.


Comment

We report here the structure of the title compound, (I), which we compare with the structures of four other closely related 2,6-substituted 4-nitroanilines, namely 2,6-dichloro-4-nitroaniline, (II) (Hughes & Trotter, 1971), 2,6-dibromo-4-nitroaniline, (III) (Bryant *et al.*, 1998), 2-bromo-6-cyano-4-nitroaniline, (IV) (Glidewell *et al.*, 2002), and 2-iodo-6-methoxy-4-nitroaniline, (V) (Garden *et al.*, 2005).

In molecules of (I) (Fig. 1), the Br and Cl substituents are disordered between the 2- and 6-positions in the aryl ring. Refinement of the site occupancies showed that position 2 is occupied equally by the two substituents, whereas there is slight preponderance of Cl at position 6, corresponding to co-crystallization of (I) with 10% of the dichloro analogue (II); compound (II) is isomorphous with (I) but not strictly isostructural. The sample of (I) originated in an industrial preparation using bromination of 2-chloro-4-nitroaniline and it seems likely that the 2,6-dichloro compound (II) may have been present as an impurity before the bromination step.

The C—C bond distances in (I) show marked bond fixation (Table 1), with the C2—C3 and C5—C6 distances significantly shorter than the rest; correspondingly, the two C—N distances are both short for their types (Allen *et al.*, 1987), while the N—O distances are both long. These observations taken all together point to the importance of the charge-separated form (Ia) (see scheme) as an important contributor to the overall molecular—electronic structure, as commonly found in 4-ni-

troanilines. Consistent with the contribution of form (Ia), the dihedral angle between the nitro group and the aryl ring is only 6.6 (2)°.



The molecules of (I) are linked into simple chains by a single N—H···O hydrogen bond (Table 2). Amine atom N1 in the molecule at (x, y, z) acts as a hydrogen-bond donor, *via* H1A, to nitro atom O41 in the molecule at $(1 - x, -\frac{1}{2} + y, \frac{3}{2} - z)$, so forming a C(8) chain (Bernstein *et al.*, 1995) running parallel to the [010] direction and generated by the 2₁ screw axis along $(\frac{1}{2}, y, \frac{3}{4})$ (Fig. 2). Two antiparallel chains of this type, related to one another by inversion, pass through each unit cell, but there are no direction specific interactions between adjacent chains; in particular, there are no potential acceptors in other chains within hydrogen-bonding distance of H1B.

In (II), the molecules are again linked by a single N—H···O hydrogen bond into C(8) chains virtually identical to those in (I) (Hughes & Trotter, 1971). Hence, the presence of a small proportion of (II) co-crystallized with (I) appears to have no significant influence on the supramolecular structure adopted by (I), which in addition retains the sharp melting point of the pure compound (Körner & Contardi, 1914). In contrast to the very simple aggregation in (I) and (II), the molecules of (III),




Figure 1

A molecule of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. For clarity, only one substituent is drawn bonded to C2 and to C6.




Figure 2

Part of the crystal structure of (I), showing the formation of a hydrogen-bonded C(8) chain along [010]. For clarity, only one substituent is drawn bonded to C2 and to C6, and H atoms bonded to C atoms have been omitted. Atoms marked with an asterisk (*) or a hash (#) are at the symmetry positions $(1 - x, -\frac{1}{2} + y, \frac{3}{2} - z)$ and $(1 - x, \frac{1}{2} + y, \frac{3}{2} - z)$, respectively.

which lie across mirror planes in space group $P2_1/m$, are linked by paired $N-H \cdots O$ hydrogen bonds into $C(8)[R_2^2(6)]$ chains of rings, further linked into sheets by bromo-nitro interactions (Bryant *et al.*, 1998). In compound (IV), the molecules are linked by a combination of one $N-H \cdots O$ and one $N-H \cdots N$ hydrogen bond into sheets of alternating $R_2^2(12)$ and $R_6^6(36)$ rings (Glidewell *et al.*, 2002), while in compound (V), hydrogen-bonded $C(8)C(8)[R_2^2(6)]$ chains of rings are linked into quite complex ribbons by two-centre iodo-nitro interactions (Garden *et al.*, 2005). In each of (III)–(V), the two $N-H$ bonds of the amine group both participate in the hydrogen bonding, in contrast to the situation in (I) and (II), where one of the $N-H$ bonds plays no role in the supramolecular aggregation.

Experimental

The sample of (I) employed was a gift from ICI; crystals suitable for single-crystal X-ray diffraction were grown by slow evaporation of a solution in ethanol [m.p. 450–452 K; literature m.p. 450 K (Körner & Contardi, 1914)].

Crystal data

$C_6H_4Br_{0.90}Cl_{1.10}N_2O_2$
 $M_r = 247.02$
 Monoclinic, $P2_1/c$
 $a = 3.8052$ (3) Å
 $b = 17.9667$ (13) Å
 $c = 12.0417$ (9) Å
 $\beta = 93.224$ (2)°
 $V = 821.95$ (11) Å³
 $Z = 4$

$D_x = 1.996$ Mg m⁻³
 Mo $K\alpha$ radiation
 Cell parameters from 1863 reflections
 $\theta = 2.0$ – 27.5°
 $\mu = 4.83$ mm⁻¹
 $T = 298$ (2) K
 Plate, red
 $0.49 \times 0.16 \times 0.06$ mm

Data collection

Bruker SMART 1000 CCD area detector diffractometer
 φ - ω scans
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{min} = 0.201$, $T_{max} = 0.749$
 5904 measured reflections
 1863 independent reflections

1357 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.037$
 $\theta_{max} = 27.5^\circ$
 $h = -4 \rightarrow 4$
 $k = -23 \rightarrow 16$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.087$
 $S = 1.02$
 1863 reflections
 129 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0477P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{max} = 0.001$
 $\Delta\rho_{max} = 0.50$ e Å⁻³
 $\Delta\rho_{min} = -0.38$ e Å⁻³

Table 1

Selected interatomic distances (Å).

C1–C2	1.408 (4)	C6–C1	1.404 (5)
C2–C3	1.373 (4)	C1–N1	1.345 (4)
C3–C4	1.387 (5)	C4–N4	1.456 (5)
C4–C5	1.387 (5)	N4–O41	1.241 (4)
C5–C6	1.364 (5)	N4–O42	1.228 (4)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N1-H1A \cdots O41^i$	0.86	2.25	3.026 (4)	151

Symmetry code: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$.

The space group $P2_1/c$ was uniquely assigned from the systematic absences. All H atoms were located from difference maps and then treated as riding atoms, with C–H distances of 0.93 Å and N–H distances of 0.86 Å, and with $U_{iso}(H)$ values of $1.2U_{eq}(C,N)$. It was apparent from an early stage that the Cl and Br substituents were disordered between the sites bonded to C2 and C6. Refinement of the site-occupancy factors for the substituents at C2 led to values of 0.503 (6) and 0.497 (6) for Cl and Br, respectively; these were thereafter fixed at 0.50. Refinement of the site-occupancy factors for the substituents at C6 led to values of 0.603 (5) and 0.397 (5) for Cl and Br, respectively; these were thereafter fixed at 0.60 and 0.40, respectively. Refinements with the occupancy factors for all the halogen sites fixed at 0.50 (to force equal populations of Cl and Br) led to significantly higher R values and hence this model was decisively rejected. Because of the fairly close similarity between the C–Cl and C–Br distances, it was necessary to apply DFIX restraints to the four independent C–X ($X = Cl$ and Br) distances.

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: OSCAIL (McArdle, 2003) and SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: OSCAIL and SHELXL97 (Sheldrick, 1997); molecular graphics: PLATON (Spek, 2003); software used to prepare material for publication: SHELXL97 and PRPKAPPA (Ferguson, 1999).

X-ray data were collected at the University of Aberdeen; the authors thank the university for funding the purchase of the diffractometer.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK1832). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (1998). *SMART*. Version 5.0. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2000). *SADABS* (Version 2.03) and *SAINTE* (Version 6.02a). Bruker AXS Inc., Madison, Wisconsin, USA.
- Bryant, R., James, S. C., Norman, N. C. & Orpen, A. G. (1998). *Acta Cryst.* **C54**, 1113–1115.
- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
- Garden, S. J., Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2005). *Acta Cryst.* **C61**, o145–o147.
- Glidewell, C., Low, J. N., McWilliam, S. A., Skakle, J. M. S. & Wardell, J. L. (2002). *Acta Cryst.* **C58**, o100–o102.
- Hughes, D. L. & Trotter, J. (1971). *J. Chem. Soc. A*, pp. 2181–2184.
- Körner, G. & Contardi, A. (1914). *Atti Accad. Lincei*, **22**, I, 823–836; *Chem. Abstr.* (1914), **8**, 73–76.
- McArdle, P. (2003). *OSCAIL for Windows*. Version 10. Crystallography Centre, Chemistry Department, NUI Galway, Ireland.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.