

L-Phenylalanyl-L-tryptophan 0.75-hydrate

Carl Henrik Görbitz

Department of Chemistry, University of Oslo, PO Box 1033 Blindern, N-0315 Oslo, Norway

Correspondence e-mail: c.h.gorbitz@kjemi.uio.no

Received 29 March 2006

Accepted 20 April 2006


Online 16 May 2006

The title compound, $C_{20}H_{21}N_3O_3 \cdot 0.75H_2O$, crystallizes as exceedingly thin fibers. The crystal packing arrangement is related to those of other hydrophobic dipeptides with phenylalanine residues, but the structure has pseudo-tetragonal symmetry in an orthorhombic space group with four peptide molecules and three water molecules in the asymmetric unit.

Comment

In a series of papers, it has been shown that dipeptides with two hydrophobic residues can form two different classes of nanoporous crystal structures (Görbitz, 2005, and references therein). The FF class, named after L-phenylalanyl-L-phenylalanine, includes also L-phenylalanyl-L-leucine (FL), L-leucyl-L-phenylalanine (LF), L-leucyl-L-leucine (LL) (Görbitz, 2001), L-isoleucyl-L-leucine (Görbitz, 2004) and L-tryptophylglycine (WG) (Emge *et al.*, 2000; Birkedal *et al.*, 2002). The common characteristic of this group is aggregation of peptide molecules into hydrophobic tubes with a hydrophilic core that incorporates a central channel filled with solvent molecules.

As part of an investigation focused on the self-assembly of FF, Reches & Gazit (2003) reported that very thin hollow fibers with a diameter of less than 300 nm could be formed by



dilution of a concentrated solution of the dipeptide in 1,1,1,3,3,3-hexafluoropropan-2-ol with water. Furthermore, it




Figure 1

The molecular structure of FW. Displacement ellipsoids are shown at the 50% probability level and H atoms are shown as spheres of arbitrary size. The minor position for the disordered tryptophan side chain of peptide molecule *D* is shown as a stick drawing.




Figure 2

The molecular packing and unit cell of FW viewed along the *a* axis. The four independent peptide molecules in the asymmetric unit have been labeled *A*, *B*, *C* and *D*. For comparison, the structures of FF, FL (Görbitz, 2001) and WG (Emge *et al.*, 2000; Birkedal *et al.*, 2002) are shown on the same scale. Atoms in side chains are shown in a darker tone.

was reported that fibers of similar dimensions could be formed by *L*-phenylalanyl-*L*-tryptophan (FW). Following continued research efforts on the nature of the FF fibers (Görbitz, 2006), we wondered what the nature of the FW fibers could be and decided to test if it was possible to grow them large enough for single-crystal structure determination. This proved to be a challenging task, but eventually needles with diameters of up to 20 μm were grown by diffusion of acetonitrile into a saturated 1,1,1,3,3,3-hexafluoropropan-2-ol solution of FW (Görbitz, 2006). The structure of this peptide is presented in detail here.

The asymmetric unit of FW, shown in Fig. 1, contains four peptide molecules and three water molecules. The main chains of the peptide molecules have rather similar conformations, but the phenylalanine side chain of molecule *A* is in a *gauche*+ orientation, as opposed to the more common *trans* orientation adopted by molecules *B*, *C* and *D*. Furthermore, even if all tryptophan side chains have well defined *gauche*– χ^1 torsion angles (N2–C10–C11–C12), the $\chi^{2,1}$ torsion angles (C10–C11–C12–C13) differ considerably (Table 1).

The crystal packing arrangement of FW is compared in Fig. 2 with the crystal structures of FF, FL (Görbitz, 2001) and WG

(Emge *et al.*, 2000; Birkedal *et al.*, 2002). The hexagonal FF structure and the tetragonal WG structure both have one peptide molecule in the asymmetric unit, while $Z' = 2$ for the orthorhombic structure of FL as well as for the structures of LL, LF (Görbitz, 2001) and IL (Görbitz, 2004) (not shown). It follows that the water-filled channels of FW are the first to be devoid of crystallographic symmetry, and they also have a more irregular appearance than those of the other structures in the family.

In accordance with previous findings, each peptide amino group donates one of its H atoms to a water molecule located in the channel (Table 2). In WG, the side-chain $\text{N}^\epsilon\text{—H}$ donor manages to find a carboxylate acceptor. In the present structure, the equivalent four H atoms are accepted by aromatic groups, two by phenylalanyl side chains and two by the six-membered ring of the tryptophan side chains.

Experimental

The title compound was obtained from Bachem. Extremely thin fibers were grown by diffusion of acetonitrile into a saturated 1,1,1,3,3,3-hexafluoropropan-2-ol solution (50 μl) of the peptide. A 20 \times 18 μm cross-section specimen was used for data collection.

Crystal data

C₂₀H₂₁N₃O₃·0.75H₂O
M_r = 364.91
 Orthorhombic, *P*₂₁₂₁
a = 5.6207 (6) Å
b = 35.556 (4) Å
c = 35.835 (4) Å
V = 7161.5 (15) Å³
Z = 16
D_x = 1.354 Mg m⁻³
 Mo *K*α radiation
μ = 0.10 mm⁻¹
T = 105 (2) K
 Needle, colorless
 0.540 × 0.020 × 0.018 mm

Data collection

Siemens SMART CCD
 diffractometer
ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
T_{min} = 0.876, *T_{max}* = 0.998
 33425 measured reflections
 7087 independent reflections
 3234 reflections with *I* > 2σ(*I*)
R_{int} = 0.170
θ_{max} = 25.0°

Refinement

Refinement on *F*²
R[*F*² > 2σ(*F*²)] = 0.079
wR(*F*²) = 0.182
S = 1.04
 7087 reflections
 726 parameters
 H atoms treated by a mixture of
 independent and constrained
 refinement
w = 1/[σ²(*F_o*²) + (0.064*P*)²]
 where *P* = (*F_o*² + 2*F_c*²)/3
 (Δ/σ)_{max} = 0.004
 Δρ_{max} = 0.45 e Å⁻³
 Δρ_{min} = -0.37 e Å⁻³
 Extinction correction: SHELXL97
 Extinction coefficient: 0.0127 (6)

Table 1

Selected torsion angles (°) in molecules *A–D* of (I).

	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>
N1–C1–C9–N2	144.6 (7)	117.9 (8)	106.6 (8)	108.5 (8)
C1–C9–N2–C10	-179.0 (6)	177.9 (6)	-174.2 (6)	-171.3 (6)
C9–N2–C10–C20	56.0 (10)	51.2 (10)	55.1 (9)	51.2 (9)
N2–C10–C20–O2	29.6 (11)	41.7 (11)	33.3 (10)	39.2 (11)
N1–C1–C2–C3	56.7 (8)	172.8 (7)	-177.0 (6)	179.3 (6)
C1–C2–C3–C4	-97.2 (10)	-117.6 (10)	-116.5 (9)	-113.4 (9)
C1–C2–C3–C8	81.5 (10)	57.7 (12)	62.9 (11)	64.3 (10)
N2–C10–C11–C12	-55.5 (8)	-81.0 (8)	-64.7 (8)	-74.6 (10)
C10–C11–C12–C13	-31.9 (11)	-4.1 (12)	62.5 (11)	72.9 (17)
C10–C11–C12–C15	148.0 (8)	-179.0 (8)	-118.0 (9)	-90.6 (14)

Owing to the combination of a large unit cell and a small crystal, more than 80% of the reflections with 2θ between 40 and 50° were unobserved, resulting in a high value for *R_{int}*. In order not to further impair the rather poor reflection-to-parameter ratio, only O atoms, N atoms and side-chain C atoms that had large *U_{iso}* values in the initial isotropic refinement were refined anisotropically. Other C atoms were refined isotropically. Covalent bond lengths and angles in each peptide molecule were restrained, using SHELXTL (Bruker, 2000) SAME commands, to values fairly similar to those of corresponding geometric parameters in the other three peptide molecules. The tryptophan side chain of peptide molecule *D* is disordered over two nearby positions with occupancies of 0.620 (14) and 0.380 (14), respectively. C- and N-bound H atoms were positioned with idealized geometry and fixed N–H and C–H distances in the range 0.88–1.00 Å. Six water H atoms were positioned by consideration of the local atomic environment, but three of them could also be detected in electron-density maps. The intramolecular water geometries were restrained by tight DFIX commands and the *s.u.* values associated with these atoms are underestimated. *U_{iso}*(H) atoms were set at 1.2*U_{eq}* of the carrier atom or 1.5*U_{eq}* for amine groups and water molecules. In the absence of significant anomalous scattering effects, 5310 Friedel pairs were merged. The absolute configuration was known for the purchased material.

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> –H··· <i>A</i>	<i>D</i> –H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> –H··· <i>A</i>
N1A–H1A···O1W	0.91	2.17	2.917 (11)	139
N1A–H1A···O3W	0.91	2.45	3.135 (10)	132
N1A–H2A···O3D ⁱ	0.91	1.87	2.752 (9)	162
N1A–H3A···O2D	0.91	1.96	2.740 (9)	143
N2A–H4A···O3A ⁱ	0.88	1.99	2.791 (8)	151
N3A–H5A···C19A ⁱⁱ	0.88	2.55	3.365 (10)	155
C1A–H11A···O1A ⁱ	1.00	2.59	3.302 (8)	128
N1B–H1B···O2W ⁱ	0.91	1.92	2.731 (10)	147
N1B–H2B···O2C	0.91	1.88	2.751 (9)	159
N1B–H3B···O3C ⁱ	0.91	1.95	2.811 (8)	158
N2B–H4B···O3B ⁱ	0.88	2.01	2.807 (8)	150
N3B–H5B···C5C ⁱⁱⁱ	0.88	2.66	3.458 (11)	151
C1B–H11B···O1B ⁱ	1.00	2.45	3.383 (9)	155
N1C–H1C···O2W	0.91	1.97	2.842 (9)	160
N1C–H2C···O3A ⁱ	0.91	1.91	2.800 (9)	166
N1C–H3C···O2A	0.91	1.89	2.788 (9)	171
N2C–H4C···O3C ⁱ	0.88	1.99	2.784 (8)	150
N3C–H5C···C6D ^{iv}	0.88	2.77	3.542 (10)	147
C1C–H11C···O1C ⁱ	1.00	2.47	3.333 (8)	144
N1D–H1D···O3W	0.91	2.02	2.809 (9)	144
N1D–H2D···O3B ⁱ	0.91	1.85	2.763 (9)	175
N1D–H3D···O2B	0.91	1.92	2.815 (9)	166
N2D–H4D···O3D ⁱ	0.88	2.00	2.730 (8)	139
N3D–H5D···C17C ^v	0.88	2.39	3.257 (13)	167
C1D–H11D···O1D ⁱ	1.00	2.47	3.278 (8)	138
O1W–H11W···O2A	0.85 (1)	1.95 (2)	2.793 (9)	172 (9)
O1W–H12W···O3W ^{vi}	0.86 (1)	2.37 (4)	3.138 (9)	148 (6)
O2W–H21W···O2C ⁱ	0.85 (1)	1.93 (2)	2.750 (9)	159 (7)
O2W–H22W···O3W	0.85 (1)	1.93 (2)	2.774 (9)	170 (9)
O3W–H31W···O2B ⁱ	0.86 (1)	1.84 (2)	2.684 (8)	167 (9)
O3W–H32W···O2D ⁱ	0.86 (1)	1.84 (3)	2.664 (9)	161 (7)

Symmetry codes: (i) *x* + 1, *y*, *z*; (ii) *x* + ½, -*y* + ½, -*z* + 1; (iii) -*x* + 1, *y* + ½, -*z* + ½; (iv) -*x* + 1, *y* - ½, -*z* + ½; (v) -*x* + ½, -*y* + 1, *z* + ½; (vi) *x* - 1, *y*, *z*.

Data collection: SMART (Bruker, 1998); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXTL (Bruker, 2000); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

The purchase of the diffractometer was made possible through support from the Research Council of Norway (NFR).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FG3015). Services for accessing these data are described at the back of the journal.

References

Birkedal, H., Schwarzenbach, D. & Pattison, P. (2002). *Angew. Chem. Int. Ed.* **41**, 754–756.
 Bruker (1998). SMART. Version 5.054. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2000). SHELXTL. Version 6.10. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2001). SAINT-Plus. Version 6.22. Bruker AXS Inc., Madison, Wisconsin, USA.
 Emge, T. J., Agrawal, A., Dalessio, J. P., Dukovic, G., Inghrim, J. A., Janjua, K., Macaluso, M., Robertson, R. R., Stiglic, T. J., Volovik, Y. & Georgiadis, M. M. (2000). *Acta Cryst.* **C56**, e469–e471.
 Görbitz, C. H. (2001). *Chem. Eur. J.* **7**, 5153–5159.
 Görbitz, C. H. (2004). *Acta Cryst.* **E60**, o626–o628.
 Görbitz, C. H. (2005). *CrystEngComm*, **7**, 670–673.
 Görbitz, C. H. (2006). *Chem. Commun.* In the press.
 Reches, M. & Gazit, E. (2003). *Science*, **300**, 625–627.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.