

The honeycomb network structure of poly[[μ_2 -1,3-bis(1*H*-benzimidazol-2-yl)benzene- κ^2 N³:N^{3'}](μ_2 -terephthalato- κ^2 O:O')zinc(II)] ethanol solvate]

Fa-Yan Meng,^a Xu-Qing Liu^a and Seik Weng Ng^{b*}

^aSchool of Chemistry and Materials Engineering, Huangshi Institute of Technology, Huangshi 435003, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my

Received 27 April 2007


Accepted 5 June 2007

Online 23 June 2007

The terephthalate dianion and the bis(imidazolyl)benzene ligand of the title compound, $\{[\text{Zn}(\text{C}_8\text{H}_4\text{O}_4)(\text{C}_{20}\text{H}_{14}\text{N}_4)] \cdot \text{C}_2\text{H}_6\text{O}\}_n$, each bridges two adjacent zinc centers, resulting in a layer-type coordination polymer; the zinc center shows tetrahedral coordination. The disordered ethanol solvent molecules occupy the spaces between the layers and are hydrogen bonded to the layers. The two symmetry-independent dianions lie on different inversion sites.

Comment

Zinc terephthalate forms a number of 1:1 adducts with bidentate N-heterocyclic ligands; crystallographic studies have focused on rigid heterocycles, such as, for example, 4,4'-bipyridine (Tao *et al.*, 2000) and 1,10-phenanthroline (Sun *et al.*, 2001). As the terephthalate entity itself is rigid, the adducts adopt architectures that reflect such rigidity, *i.e.* the 4,4'-bipyridine adduct is a three-dimensional network and the 1,10-phenanthroline adduct is a linear chain. With a flexible ligand such as 1,3,4-bis(triazol-1-yl)ethane, a two-dimensional rhombic network is formed (Li *et al.*, 2004). 1,3,4-Bis(triazol-1-yl)butane affords a threefold interpenetrating diamondoid network based on a zinc center (Wang *et al.*, 2006).



1,3-Bis(benzimidazol-2-yl)benzene, an N-heterocycle that can be conveniently synthesized by condensing *m*-phenylenediamine and phthalic acid under the influence of micro-

waves (Song *et al.*, 2000), is a ligand that can bind to two metal centers through its two tertiary nitrogen-donor sites. Spectroscopic measurements are available for a small number of adducts with transition metals, and among the zinc complexes that have been characterized spectroscopically are the halide and perchlorate complexes (Chawla & Gill, 1997; Shivakumarraiah & Nanje Gowda, 2003), but to date no example of an adduct has been reported in the literature (CSD, Version 5.28, November 2006 release; Allen, 2002).

The title 1:1 zinc terephthalate–bis(benzimidazolyl)benzene adduct, (I), which exists as an ethanol solvate, is a coordination polymer in which the carboxylate group and benzene ligand both engage in bridging two tetrahedral zinc centers (Fig. 1), this bridging mode giving rise to a layer motif. The two symmetry-independent dianions lie on different inversion sites. The two benzimidazolyl arms are bent with respect to the phenylene ring, one (with the N1 atom) being twisted by 31.6 (1)° and the other (with the N3 atom) being twisted by 15.7 (1)°. The twist angles parallel the deviation of the metal atom from the plane of the arms, the atom being displaced by 0.807 (3) Å from the first arm and by 0.507 (3) Å from the second arm.

The topology of the layer is a (6,3)-honeycomb (Fig. 2) that is somewhat buckled as the two bridging components are of




Figure 1

The tetrahedral geometry of the Zn atom in the layer structure of (I). Displacement ellipsoids are drawn at the 70% probability level. The disordered ethanol solvent molecule is not shown. [Symmetry code: (i) $-x, 1 - y, 1 - z$.]




Figure 2

OLEX (Dolomanov *et al.*, 2003) depiction of the (6,3)-honeycomb topology, shown projected against the unit cell.

different sizes. The atoms comprising the layer framework constitute only about 79% of the volume, as calculated by *PLATON* (Spek, 2003). The layers are consolidated by N—H···O hydrogen bonds (Table 2), but the space between layers is large enough to accommodate ethanol molecules.

Experimental

The N-heterocycle was prepared according to a reported procedure (Chawla & Gill, 1997). Zinc nitrate hexahydrate (0.074 g, 0.25 mmol), benzene-1,4-dicarboxylic acid (0.021 g, 0.125 mmol), 1,3-bis(benzimidazol-2-ylmethyl)benzene (0.039 g, 0.125 mmol), ethanol (2 ml) and water (15 ml) were placed in a 23 ml Teflon-lined stainless steel Parr bomb. (Neither sodium nor potassium hydroxide was added.) The bomb was heated at 433 K for 5 d and cooled to room temperature at a rate of 5 K h⁻¹. Colorless block-shaped crystals were obtained in 20% yield.

Crystal data

[Zn(C ₈ H ₄ O ₄)(C ₂₀ H ₁₄ N ₄)]·C ₂ H ₆ O	$\gamma = 87.931 (1)^\circ$
$M_r = 585.90$	$V = 1331.1 (2) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 9.0660 (8) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.3077 (9) \text{ \AA}$	$\mu = 0.97 \text{ mm}^{-1}$
$c = 15.921 (2) \text{ \AA}$	$T = 291 (2) \text{ K}$
$\alpha = 86.731 (1)^\circ$	$0.28 \times 0.21 \times 0.18 \text{ mm}$
$\beta = 83.140 (1)^\circ$	

Data collection

Bruker APEX area-detector diffractometer	10765 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	5834 independent reflections
$T_{\min} = 0.712$, $T_{\max} = 0.845$	4918 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.098$	$\Delta\rho_{\text{max}} = 0.38 \text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\text{min}} = -0.41 \text{ e \AA}^{-3}$
5834 reflections	
389 parameters	
31 restraints	

Table 1

Selected geometric parameters (\AA , $^\circ$).

Zn1—O1	1.930 (2)	Zn1—N1	2.040 (2)
Zn1—O3	1.966 (2)	Zn1—N3 ⁱ	2.036 (2)
O1—Zn1—O3	106.0 (1)	O3—Zn1—N1	98.9 (1)
O1—Zn1—N3 ⁱ	125.1 (1)	O3—Zn1—N3 ⁱ	104.2 (1)
O1—Zn1—N1	115.7 (1)	N1—Zn1—N3 ⁱ	103.30 (7)

Symmetry code: (i) $-x, -y + 1, -z + 1$.

The ethanol solvent molecule is disordered in the C atoms and was refined with two ethyl components in a 0.72:0.28 ratio. The O—C distance was restrained to 1.45 (1) \AA and the C—C distance to 1.50 (1) \AA . The displacement parameters of the four C atoms were restrained to be nearly isotropic.

Table 2

Hydrogen-bond geometry (\AA , $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2 n ···O4 ⁱⁱ	0.84 (2)	1.94 (2)	2.777 (2)	176 (3)
N4—H4 n ···O2	0.86 (2)	2.11 (2)	2.895 (3)	152 (3)
O5—H5 o ···O1	0.86 (4)	2.14 (4)	2.999 (4)	176 (7)

Symmetry code: (ii) $x, y + 1, z$.

Carbon-bound H atoms were positioned geometrically (C—H = 0.93 and 0.97 \AA), and were included in the refinement in the riding-model approximation, with $U_{\text{iso}}(\text{H})$ values set at 1.2–1.5 $U_{\text{eq}}(\text{C})$. The water and amino H atoms were located in a difference Fourier map, and were refined with a distance restraint of O—H = N—H = 0.85 (1) \AA .

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2007).

The authors thank the Key Laboratory of Non-Ferrous Metal Materials and New Processing Technology, Ministry of Education, China, Huangshi Institute of Technology and the University of Malaya for supporting this work.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BG3037). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2004). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chawla, S. K. & Gill, B. K. (1997). *Polyhedron*, **16**, 1315–1322.
- Dolomanov, O. V., Blake, A. J., Champness, N. R. & Schröder, M. (2003). *J. Appl. Cryst.* **36**, 1283–1284.
- Li, B.-L., Li, B.-Z., Zhu, X., Lu, X.-H. & Zhang, Y. (2004). *J. Coord. Chem.* **16**, 1367–1371.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Shivakumaraiah & Nanje Gowda, N. M. (2003). *Synth. React. Inorg. Met. Org. Chem.* **33**, 1207–1220.
- Song, L.-Q., Xu, X.-L. & Tan, G.-Z. (2000). *Jingxi Huagong (Fine Chemicals)*, **17**, 256–257.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Sun, D.-F., Cao, R., Liang, Y. C., Shi, Q., Su, W.-P. & Hong, M.-C. (2001). *J. Chem. Soc. Dalton Trans.* pp. 2335–2340.
- Tao, J., Tong, M.-L. & Chen, X.-M. (2000). *J. Chem. Soc. Dalton Trans.* pp. 3669–3674.
- Wang, X.-L., Qin, C., Wang, E.-B. & Su, Z.-M. (2006). *Chem. Eur. J.* **12**, 2680–2691.
- Westrip, S. P. (2007). *pubCIF*. In preparation.