

2-Cyano-4-iodoacetanilide: a hydrogen-bonded chain of $R_2^2(12)$ and $R_2^2(14)$ rings

Simon J. Garden,^a Cíntia de A. Custódio,^b James L. Wardell,^a John N. Low^c and Christopher Glidewell^{d*}

^aUniversidade Federal do Rio de Janeiro, Instituto de Química, Cidade Universitária, CT, Bloco A, Ilha do Fundão, Rio de Janeiro, RJ, CEP 21941-909, Brazil, ^bInstituto de Engenharia Nuclear, Via Cinco, Cidade Universitária, Ilha do Fundão, Rio de Janeiro, RJ, CEP 21945-970, Brazil, ^cDepartment of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland, and ^dSchool of Chemistry, University of St Andrews, Fife KY16 9ST, Scotland
Correspondence e-mail: cg@st-andrews.ac.uk

Received 30 May 2007


Accepted 31 May 2007

Online 14 June 2007

The molecules of 2-cyano-4-iodoacetanilide, $C_9H_7IN_2O$, are linked by $N-H\cdots N$ and $C-H\cdots O$ hydrogen bonds into chains of alternating $R_2^2(12)$ and $R_2^2(14)$ rings.

Comment

We report here the molecular and supramolecular structures of 2-cyano-4-iodoacetanilide, (I) (Fig. 1), and compare the structure of (I) with that of the simpler analogue 2-cyanoacetanilide, (II), for which we have re-interpreted the recently published structure (Arslan *et al.*, 2005). The present study is a continuation of our extended programme on the supramolecular structures of substituted iodoaniline derivatives (Ferguson *et al.*, 2005; Garden *et al.*, 2002, 2004, 2005, 2006; Garden, Glidewell *et al.*, 2001; Glidewell *et al.*, 2004; McWilliam *et al.*, 2001).



In (I), the $C6-C1-N1$, $C1-N1-C11$ and $N1-C11-O1$ angles are all significantly larger than 120° (Table 1), suggesting that the short intramolecular $H6\cdots O1$ contact (Table 2) may be repulsive rather than attractive. Despite this, the molecule of (I) is almost planar, as indicated by the leading torsion angles, and it is thus possible that the overall molecular conformation results from a balance between repulsive intramolecular contacts and attractive intermolecular

hydrogen bonds. The remaining bond lengths and angles show no unusual features.

Two intermolecular hydrogen bonds link the molecules of (I) into chains of fused rings. Atoms N1 and C5 in the molecule at (x, y, z) act as hydrogen-bond donors, respectively, to atoms N2 in the molecule at $(2-x, 1-y, 1-z)$ and O1 in the molecule at $(-x, 2-y, 1-z)$. Propagation by inversion of these two interactions generates a chain of rings running parallel to the $[2\bar{1}0]$ direction, with centrosymmetric $R_2^2(12)$ rings (Bernstein *et al.*, 1995) containing $N-H\cdots N$ hydrogen bonds and centred at $(2n+1, \frac{1}{2}-n, \frac{1}{2})$ (where n represents zero or an integer) alternating with centrosymmetric $R_2^2(14)$ rings containing $C-H\cdots O$ hydrogen bonds and centred at $(2n, 1-n, \frac{1}{2})$ (where n represents zero or an integer) (Fig. 2).

The structure of the related compound (II) has recently been reported (Arslan *et al.*, 2005). The structure was described as forming sheets parallel to (100) generated by a combination of $N-H\cdots N$ and $C-H\cdots O$ hydrogen bonds. The formation of a two-dimensional supramolecular structure in the space group $P2_12_12_1$ is somewhat unexpected, and re-analysis of the published structure shows that the hydrogen-bonded structure of (II) is, in fact, three-dimensional. The $N-H\cdots N$ and $C-H\cdots O$ hydrogen bonds, acting individually, form chains running parallel to the [100] and [010] directions, of $C(6)$ and $C(8)$ types, respectively, while in combination they form a $C_2^2(10)$ chain running parallel to the [001] direction, so completing the three-dimensional framework. The original report on (II) (Arslan *et al.*, 2005) showed the structure in projection along [100], and the published packing diagram appears to show the formation of $R_2^2(12)$ rings containing pairs of $N-H\cdots N$ hydrogen bonds, when in fact this hydrogen bond forms $C(6)$ chains along the direction of the projection and thus normal to the plane of the supposed hydrogen-bonded sheet.

Although the iodine substituent in (I) plays no direct role in the supramolecular aggregation, its presence leads to an




Figure 1

The molecule of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.

entirely different supramolecular structure in (I) as compared with (II), even though the structures of (I) and (II) are each built from the combination of one N—H···N hydrogen bond and one C—H···O hydrogen bond.




Figure 2
A stereoview of part of the crystal structure of (I), showing a chain of alternating $R_2^2(12)$ and $R_2^2(14)$ rings running parallel to $[2\bar{1}0]$. For the sake of clarity, H atoms not involved in the motifs shown have been omitted.

Experimental

Compound (I) was prepared by the acetylation of 2-amino-5-iodobenzonitrile, prepared as follows. $K(\text{ICl}_2)$ (Garden, Torres *et al.*, 2001) (3 ml of a 2 mol dm^{-3} solution in water) was added to a methanol solution (15 ml) of 2-aminobenzonitrile (0.510 g, 5.0 mmol). The reaction mixture was stirred at room temperature. After 2 h, the product had precipitated and the benzonitrile substrate had been completely consumed. The reaction mixture was diluted with water (50 ml) and the product was collected by filtration and dried in air (yield 93%). Recrystallization from aqueous ethanol (97:3 *v/v*) gave pale-brown plates [m.p. 356–357 K; literature (Harris *et al.*, 1992) 358 K]. NMR (CDCl_3): δ_{H} 6.20 (2H, s, NH_2), 6.62 (1H, *d*, $J = 8.8$ Hz), 7.52 (1H, *dd*, $J = 8.8$ and 2.0 Hz), 7.65 (1H, *d*, $J = 2.0$ Hz); δ_{C} : 74.9, 96.0, 116.8, 117.8, 139.7, 142.2, 151.2. IR (KBr disk, cm^{-1}): 3456, 3359, 2224, 1643, 1552, 1483, 1402, 1155, 899, 820. The product (0.500 g, 2.0 mmol) was treated with acetic anhydride (1 ml) and acetic acid (2 ml). The reaction mixture was warmed gently for a few minutes then hydrolyzed with water (20 ml). The product, (I), was collected by filtration and recrystallized from aqueous ethanol (97:3 *v/v*) (yield 79%, m.p. 449–451 K). NMR (CDCl_3): δ_{H} 2.27 (3H, s, CH_3), 7.69 (1H, *br*, s, NH), 7.85 (1H, *dd*, $J = 10$ and 1.9 Hz, H-5), 7.87 (1H, *d*, $J = 1.9$ Hz, H-3), 8.19 (1H, *d*, $J = 10$ Hz, H-6); δ_{C} 24.4, 85.5, 103.2, 114.5, 122.4, 139.7, 139.9, 142.7, 168.2. IR (KBr disk, cm^{-1}): 3344, 3054, 2225, 1689, 1577, 1516, 1383, 1369, 1295, 1258, 1182, 1122, 884, 842.

Crystal data

$\text{C}_9\text{H}_7\text{IN}_2\text{O}$	$\gamma = 90.926$ (2) $^\circ$
$M_r = 286.07$	$V = 476.51$ (3) \AA^3
Triclinic, $P\bar{1}$	$Z = 2$
$a = 4.6657$ (2) \AA	Mo $K\alpha$ radiation
$b = 9.4172$ (4) \AA	$\mu = 3.32$ mm^{-1}
$c = 11.6841$ (4) \AA	$T = 120$ (2) K
$\alpha = 109.7520$ (17) $^\circ$	$0.24 \times 0.06 \times 0.02$ mm
$\beta = 98.598$ (2) $^\circ$	

Data collection

Bruker–Nonius KappaCCD diffractometer	7314 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)	2160 independent reflections
$T_{\text{min}} = 0.503$, $T_{\text{max}} = 0.937$	2047 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$	119 parameters
$wR(F^2) = 0.058$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.56$ e \AA^{-3}
2160 reflections	$\Delta\rho_{\text{min}} = -0.49$ e \AA^{-3}

Table 1

Selected bond angles ($^\circ$).

C6—C1—N1	122.5 (3)	N1—C11—O1	122.6 (3)
C1—N1—C11	126.8 (2)		

Table 2

Hydrogen-bond geometry (\AA , $^\circ$).

$D\text{—}H\cdots A$	$D\text{—}H$	$H\cdots A$	$D\cdots A$	$D\text{—}H\cdots A$
N1—H1···N2 ⁱ	0.86	2.54	3.309 (4)	149
C5—H5···O1 ⁱⁱ	0.95	2.38	3.214 (4)	147
C6—H6···O1	0.95	2.24	2.815 (4)	118

Symmetry codes: (i) $-x + 2, -y + 1, -z + 1$; (ii) $-x, -y + 2, -z + 1$.

Crystals of (I) are triclinic; the space group $P\bar{1}$ was selected and confirmed by the subsequent structure analysis. All H atoms were located in difference maps and then treated as riding atoms in geometrically idealized positions, with C—H distances of 0.95 (aromatic) or 0.98 \AA (methyl), N—H distances of 0.86 \AA and $U_{\text{iso}}(\text{H}) = kU_{\text{eq}}(\text{C,N})$, where $k = 1.5$ for the methyl group and $k = 1.2$ for all other H atoms.

Data collection: COLLECT (Nonius, 1999); cell refinement: DENZO (Otwinowski & Minor, 1997) and COLLECT; data reduction: DENZO and COLLECT; program(s) used to solve structure: OSCAIL (McArdle, 2003) and SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: OSCAIL and SHELXL97 (Sheldrick, 1997); molecular graphics: PLATON (Spek, 2003); software used to prepare material for publication: SHELXL97 and PRPKAPPA (Ferguson, 1999).

X-ray data were collected at the EPSRC National Crystallography Service, University of Southampton, England. The authors thank the staff for all their help and advice. SJG thanks CNPq, and JLW thanks CNPq and FAPERJ for financial support.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK3136). Services for accessing these data are described at the back of the journal.

References

- Arslan, B., Kazak, C., Kirilmis, C., Koca, M. & Büyüküngör, O. (2005). *Acta Cryst.* **E61**, o1652–o1653.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.

- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
- Ferguson, G., Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2005). *Acta Cryst.* **C61**, o445–o449.
- Garden, S. J., Fontes, S. P., Wardell, J. L., Skakle, J. M. S., Low, J. N. & Glidewell, C. (2002). *Acta Cryst.* **B58**, 701–709.
- Garden, S. J., Glidewell, C., Low, J. N., McWilliam, S. A., Pinto, A. C., Skakle, J. M. S., Torres, J. C. & Wardell, J. L. (2001). *Acta Cryst.* **C57**, 1212–1214.
- Garden, S. J., Glidewell, C., Low, J. N., Skakle, J. M. S. & Wardell, J. L. (2005). *Acta Cryst.* **C61**, o145–o147.
- Garden, S. J., Torres, J. C., Melo, S. C. D., Lima, A. S., Pinto, A. C. & Lima, E. L. S. (2001). *Tetrahedron Lett.* **42**, 2089–2092.
- Garden, S. J., Wardell, J. L., Low, J. N., Skakle, J. M. S. & Glidewell, C. (2006). *Acta Cryst.* **E62**, o3762–o3764.
- Garden, S. J., Wardell, J. L., Skakle, J. M. S., Low, J. N. & Glidewell, C. (2004). *Acta Cryst.* **C60**, o328–o330.
- Glidewell, C., Low, J. N., Skakle, J. M. S., Wardell, S. M. S. V. & Wardell, J. L. (2004). *Acta Cryst.* **B60**, 472–480.
- Harris, N. V., Smith, C. & Bowden, K. (1992). *Eur. J. Med. Chem.* **27**, 7–18.
- McArdle, P. (2003). *OSCAIL for Windows*. Version 10. Crystallography Centre, Chemistry Department, NUI Galway, Ireland.
- McWilliam, S. A., Skakle, J. M. S., Low, J. N., Wardell, J. L., Garden, S. J., Pinto, A. C., Torres, J. C. & Glidewell, C. (2001). *Acta Cryst.* **C57**, 942–945.
- Nonius (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (2003). *SADABS*. Version 2.10. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.