

A new sulfate acid polymorph of 1,3-dihydrobenzotriazole

Angel Ramos-Organillo^{a*} and Rosalinda Contreras^b


^aFacultad de Ciencias Químicas, Universidad de Colima, km 9 carretera Colima-Coquimatlán, Coquimatlán, Colima, CP 28400, Mexico, and ^bDepartamento de Química, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apartado Postal 14-740, 07000 México, DF, Mexico
Correspondence e-mail: aaramos@uocol.mx

Received 30 April 2007
Accepted 26 June 2007
Online 9 August 2007

A new sulfate acid polymorph of 1,3-dihydrobenzotriazole, viz. 1,3-dihydrobenzotriazolium hydrogensulfate, $C_6H_6N_3^+ \cdot HSO_4^-$, differs from an existing polymorph in that the polymeric interaction between the HSO_4^- anions, together with different classical ($D-H \cdots A$) and nonclassical ($C-H \cdots A$) interactions, changes the space group.

Comment

Heterocyclic compound (I) (see scheme) contains acidic H atoms and N atoms with lone electron pairs. The presence of both acidic and basic characteristics gives the molecule the ability to participate in a wide variety of interactions. Moreover, tautomerism in (I) can change the reactivity depending on the starting material (Jagerovic *et al.*, 2002; Katritzky *et al.*, 1998; Gallinek, 1897; Elderfiel, 1957). This diversity gives rise to the possibility of different structural arrangements or polymorphs. For aliphatic amines this outcome is expected, but in this case the molecule is planar and polymorphs are not common (Blagden & Davey, 2003; Davey, 2003; Mak & Zhou, 1992; Dunitz, 1979).



The title compound, (II), crystallized in the monoclinic space group $P2_1/c$ from a mixture of tetrahydrofuran (THF) and hexane. The structure (Fig. 1) shows the presence of acidic H atoms on atom N3 and on the HSO_4^- counter-ion. The crystal packing (Fig. 2) is structured by classical and nonclassical hydrogen bonds, the most important of which are listed in Table 1. Three of these are classical hydrogen bonds ($N-H \cdots O$), while the others are nonclassical ($C-H \cdots O$). Two particularly strong interactions are $H3 \cdots O4$ and $H1 \cdots O3$. Some of the hydrogen bonds are complex. For

example, the $N1-H1$ group is considered a bifurcated donor because it interacts with atoms N2 and O3. Atom O2 is considered as a trifurcated acceptor because it interacts with the $C6-H6$, $C4-H4$ and $C7-H7$ bonds. All such interactions are important because they contribute to the geometry of the lattice. The HSO_4^- ions are joined together by strong $O-H \cdots O=S$ hydrogen bonds that are nearly ideal in geometry (Steiner, 1998, 2002; Jeffrey, 1997).

As a result of these interactions, the 1,3-dihydrobenzotriazolium cations pack in a herring-bone pattern in the ab plane with the hydrogensulfate anions interspersed (Fig. 3). A polymorphic structure of (II) was reported by Giordano (1980), which crystallizes in the orthorhombic space group $Pbcn$. The phosphoric acid salt of 1,3-dihydrobenzotriazole, (IV), is also known (Emsley *et al.*, 1988) and crystallizes in the triclinic space group $P\bar{1}$ with similar packing to (II).

In Table 2, the bond distances for polymorphs (II) and (III) (Giordano, 1980) and compound (IV) (Emsley *et al.*, 1988) are compared. Polymorphs (II) and (III) show similar bond distances, while in (IV) they are slightly longer. It can also be




Figure 1
A view of the molecule of (II), showing 50% probability displacement ellipsoids and the atom-numbering scheme.






Figure 2
The polymeric arrangements in (II), formed by classical and nonclassical hydrogen bonds (dotted lines). The view is along the b axis. [Symmetry codes: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.]


Figure 3

The herring-bone arrangement formed by classical and nonclassical hydrogen bonds (dotted lines) in the crystalline network of (II). The view is along the *a* axis.


Figure 4


Hydrogen bonds (in Å) between HSO_4^- anions in (II) (upper figure; this work) and (III) (lower figure; Giordano, 1980).

observed that the distances decrease in the crystal system order *triclinic* \rightarrow *monoclinic* \rightarrow *orthorhombic*. This reflects better packing as the symmetry increases. Fig. 4 shows the rearrangement of the HSO_4^- anion corresponding to polymorphs (II) and (III). The $\text{S—O—H}\cdots\text{O}=\text{S}$ hydrogen bond [1.84 (4) and 1.478 Å] is in fact the reason that the HSO_4^- ions stay together in the supramolecular polymeric structure. We can conclude that these interactions are stronger in (III) than in (II). Consequently, the chains in (III) have an almost linear shape, while those in (II) have a zigzag shape, and the distances are shorter, stronger and more efficient in the orthorhombic lattice in (III) compared with the monoclinic lattice in (II).

Experimental

A mixture of (I) (400 mg, 3.36 mmol) and dry THF (10 ml) was cooled to 223 K; H_2SO_4 (0.18 ml, 329 mg, 3.36 mmol) was added and the mixture was stirred for half an hour. The mixture was then filtered and the white powdered product (II) (98% yield, m.p. 437–439 K)

was partially dissolved for crystallization in a THF/hexane mixture. m/z (intensity, %): 207 (1), 133.35 (100), 105 (100). IR (KBr), ν_{max} : 2100–3600 (OH), 3300 (NH), 1723 (N=N), 1612 (C=C). Analysis calculated: C 33.18, H 3.25, N 19.35, S 14.76%; observed: C 33.53, H 3.30, N 19.02, S 13.24%. The structure of (I) (see scheme below) was analyzed by ^1H and ^{13}C NMR spectroscopy, which showed a symmetrical molecule, three signals for ^1H and four for ^{13}C . The N—H chemical shift in (II) was shifted to lower frequency (2.62 p.p.m.) compared with N—H for (I). Double protonation was not observed for (I); in fact, atom N2 is not a basic position, because no different —NH signal was observed in the ^1H NMR spectrum. No significant changes were observed in the ^{13}C NMR spectrum because there were no changes in the aromatic ring.



Crystal data

$\text{C}_6\text{H}_6\text{N}_3^+\cdot\text{HSO}_4^-$
 $M_r = 217.21$
 Monoclinic, $P2_1/c$
 $a = 12.715$ (3) Å
 $b = 5.133$ (1) Å
 $c = 14.406$ (4) Å
 $\beta = 113.63$ (1)°

$V = 861.4$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.37$ mm⁻¹
 $T = 273$ (2) K
 $0.25 \times 0.20 \times 0.20$ mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
 (Blessing, 1995)
 $T_{\text{min}} = 0.913$, $T_{\text{max}} = 0.930$

9336 measured reflections
 1969 independent reflections
 1637 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.095$
 $S = 1.07$
 1969 reflections

156 parameters
 All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1 \cdots O3 ⁱ	0.83 (2)	1.93 (3)	2.695 (2)	153 (2)
O1—H1A \cdots O4 ⁱⁱ	0.82 (4)	1.84 (4)	2.659 (2)	173 (3)
N3—H3 \cdots O4	0.87 (3)	1.93 (3)	2.792 (2)	173 (3)
C4—H4 \cdots O2	0.90 (2)	2.48 (3)	3.269 (3)	146 (2)
C6—H6 \cdots O2 ⁱⁱⁱ	0.96 (3)	2.57 (3)	3.306 (3)	133.7 (19)

Symmetry codes: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

All H atoms were refined freely [$\text{C—H} = 0.90$ (2)– 0.97 (3) Å].

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *SCALEPACK* and *DENZO* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

ARO thanks CONACyT for a scholarship. Financial support by CONACyT and CINVESTAV, Mexico, is acknowledged. We thank Efrén García Baez, M. Antonio

Table 2

Comparison of selected of bond distances (Å) of (II), (III) and (IV).

	(II) ^a	(III) ^b	(IV) ^c
N1–N2	1.315 (2)	1.310	1.317 (3)
N2–N3	1.311 (2)	1.312	1.314 (3)
C4–C5	1.364 (3)	1.365	1.368 (4)
C5–C6	1.406 (3)	1.416	1.414 (4)
C6–C7	1.369 (3)	1.356	1.370 (4)
C7–C8	1.397 (3)	1.391	1.402 (4)
C8–C9	1.389 (2)	1.391	1.390 (3)
C4–C9	1.404 (3)	1.392	1.400 (4)
C9–N3	1.363 (3)	1.367	1.364 (3)
C8–N1	1.358 (3)	1.362	1.365 (3)
O1–S1	1.5576 (17)	1.539	–
O2–S1	1.4179 (18)	1.431	–
O3–S1	1.4453 (17)	1.435	–
O4–S1	1.4696 (14)	1.449	–

Notes: (a) this work, monoclinic; (b) Giordano (1980), orthorhombic; (c) Emsley *et al.* (1988), triclinic.

Leyva Ramirez and A. Corona-Bustamante for their comments for this article.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SQ3078). Services for accessing these data are described at the back of the journal.

References

- Blagden, N. & Davey, R. J. (2003). *Cryst. Growth Des.* **3**, 873–885.
- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Davey, R. J. (2003). *Chem. Commun.* pp. 1463–1467.
- Dunitz, J. D. (1979). *X-ray Analysis and the Structure of Organic Molecules*, pp. 319–322. Ithaca: Cornell University Press.
- Elderfiel, R. C. (1957). *Heterocyclic Compounds*, pp. 267–288. New York: John Wiley and Sons.
- Emsley, J., Prays, N. M., Dawes, H. M., Hursthouse, M. B. & Kuroda, R. (1988). *Phosphorus Silicon*, **35**, 141–149.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gallinek, A. (1897). *Chem. Ber.* **30**, 1909–1913.
- Giordano, F. (1980). *Acta Cryst.* **B36**, 2458–2460.
- Jagerovic, N., Jimeno, M. L., Alkorta, I., Elguero, J. & Claramunt, R. M. (2002). *Tetrahedron*, **58**, 9089–9094.
- Jeffrey, G. A. (1997). In *An Introduction to Hydrogen Bonding*. Oxford University Press.
- Katritzky, A. R., Lan, X., Yang, J. Z. & Denisko, O. V. (1998). *Chem. Rev.* **98**, 409–548.
- Mak, T. C. W. & Zhou, G.-D. (1992). *Crystallography in Modern Chemistry*, pp. 77–79. New York: John Wiley & Sons.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Steiner, T. (1998). *J. Phys. Chem. A*, **102**, 7041–7052.
- Steiner, T. (2002). *Angew. Chem. Int. Ed.* **41**, 48–76.