

7-Methoxy-4,6-dimethyl-3*H*-isobenzofuran-1-one

Jian-Feng Wang,^a Xin Lin,^a
Yao-Jian Huang,^a Wen-Jin Su,^a
Yu-Fen Zhao^b and Seik Weng
Ng^{c*}

^aDepartment of Biology, Xiamen University,
Xiamen 361005, People's Republic of China,

^bDepartment of Chemistry, Xiamen University,
Xiamen 361005, People's Republic of China,
and ^cDepartment of Chemistry, University of
Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my

Key indicators

Single-crystal X-ray study

$T = 298$ K

Mean $\sigma(\text{C}-\text{C}) = 0.006$ Å

R factor = 0.060

wR factor = 0.156


Data-to-parameter ratio = 7.0

For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.

The title compound, $\text{C}_{11}\text{H}_{12}\text{O}_3$, exists as a nearly planar molecule, the dihedral angle between the five- and six-membered rings being $1.9(2)^\circ$.

Comment

7-Methoxy-4,6-dimethyl-3*H*-isobenzofuran-1-one, (I), exists as a nearly planar molecule, the dihedral angle between the five- and six-membered rings being $1.9(2)^\circ$. The compound exhibits moderate cytotoxicity towards the KB cell line (IC_{50} $50 \mu\text{g ml}^{-1}$).



Experimental

The title compound was crystallized from the ethyl acetate extract of the liquid culture of an unidentified marine fungus isolated from the leaves of the sea lotus found in Fujian Province, China.

Crystal data

$\text{C}_{11}\text{H}_{12}\text{O}_3$
 $M_r = 192.21$
Orthorhombic, $Pna2_1$
 $a = 7.5508(7)$ Å
 $b = 12.616(1)$ Å
 $c = 10.1223(9)$ Å
 $V = 964.27(15)$ Å³
 $Z = 4$
 $D_x = 1.324$ Mg m⁻³

Mo $K\alpha$ radiation
Cell parameters from 1595
reflections
 $\theta = 2.6\text{--}21.8^\circ$
 $\mu = 0.10$ mm⁻¹
 $T = 298(2)$ K
Needle, colorless
 $0.45 \times 0.12 \times 0.09$ mm

Data collection

Bruker APEX area-detector
diffractometer
 φ and ω scans
Absorption correction: none
12924 measured reflections
897 independent reflections

840 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.070$
 $\theta_{\text{max}} = 25.0^\circ$
 $h = -8 \rightarrow 8$
 $k = -15 \rightarrow 14$
 $l = -11 \rightarrow 12$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.156$
 $S = 1.13$
897 reflections
129 parameters
H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.1111P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.15$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³


Table 1Selected geometric parameters (\AA , $^\circ$).

O1—C1	1.342 (6)	C3—C4	1.400 (7)
O1—C2	1.434 (7)	C4—C5	1.381 (7)
O2—C1	1.205 (6)	C4—C9	1.489 (7)
O3—C7	1.361 (6)	C5—C6	1.385 (6)
O3—C11	1.418 (7)	C6—C7	1.381 (6)
C1—C8	1.462 (6)	C6—C10	1.515 (6)
C2—C3	1.480 (7)	C7—C8	1.409 (6)
C3—C8	1.377 (7)		
C1—O1—C2	110.3 (4)	C3—C4—C9	121.1 (4)
C7—O3—C11	117.7 (3)	C4—C5—C6	123.7 (4)
O2—C1—O1	120.4 (5)	C7—C6—C5	120.3 (4)
O2—C1—C8	130.7 (5)	C7—C6—C10	119.3 (4)
O1—C1—C8	108.8 (4)	C5—C6—C10	120.4 (4)
O1—C2—C3	105.2 (4)	O3—C7—C6	118.8 (4)
C8—C3—C4	122.4 (4)	O3—C7—C8	123.5 (4)
C8—C3—C2	108.0 (4)	C6—C7—C8	117.5 (4)
C4—C3—C2	129.6 (5)	C3—C8—C7	120.7 (4)
C5—C4—C3	115.3 (4)	C3—C8—C1	107.7 (4)
C5—C4—C9	123.6 (5)	C7—C8—C1	131.5 (4)

The H atoms were positioned geometrically and were included in the refinement in the riding-model approximation. C—H distances were set to 0.93–0.97 \AA , with U_{iso} values for H atoms of 1.2 or 1.5 (methyl H) times U_{eq} of the parent atom. The methyl groups bonded to the aromatic ring were allowed to rotate but not to tip. In the absence of significant anomalous scattering effects, Friedel pairs were merged.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SMART*; data reduction: *SAINTE* (Bruker, 2001); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97*.

The authors thank the Natural Science Foundation of Fujian Province, China (grant No. C0110002), the Key Foundation of Science & Technology Project of Fujian

**Figure 1**

ORTEPII (Johnson, 1976) plot of (I), with displacement ellipsoids drawn at the 50% probability level. H atoms are drawn as spheres of arbitrary radii.

Province, China (grant No. 2002H011), the National High Technology Research & Development Program of China (863 Program, No. 2001AA620401) and the University of Malaya for supporting this work.

References

- Bruker (2001). *SAINTE* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.