

Muhammad S. Khan,^a Birte Ahrens,^b Paul R. Raithby^{b,c,*} and Simon J. Teat^c

^aDepartment of Chemistry, College of Science, Sultan Qaboos University, PO Box 36, Al Khod 123, Sultanate of Oman, ^bDepartment of Chemistry, University of Bath, Bath BA2 7AY, England, and ^cCCLRC Daresbury Laboratory, Daresbury, Warrington WA4 4AD, England

Correspondence e-mail: p.r.raithby@bath.ac.uk

Key indicators

Single-crystal synchrotron study
 $T = 150$ K
 Mean $\sigma(\text{C}-\text{C}) = 0.005$ Å
 Disorder in main residue
 R factor = 0.055
 wR factor = 0.114
 Data-to-parameter ratio = 13.7


For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

2,6-Bis(trimethylsilylethynyl)dithieno[3,2-*b*:2',3'-*d*]thiophene

The title compound, $\text{C}_{18}\text{H}_{20}\text{S}_3\text{Si}_2$, is a precursor in the formation of platinum and gold di-yne complexes and poly-yne polymers. These materials are of interest because of the π -conjugation that extends through the fused oligothiophenyl linker unit along the rigid backbone of the polymer. In the structure of the title compound, the oligothiophenyl group is planar and the trimethylsilylalkyne groups are essentially linear.

Comment

The title compound, 2,6-bis(trimethylsilylethynyl)dithieno[3,2-*b*:2',3'-*d*]thiophene, (I), is a trimethylsilyl-protected di-yne. It is a precursor in the formation of the following series of compounds: the terminal di-yne, $\text{H}-\text{C}\equiv\text{C}-\text{R}-\text{C}\equiv\text{C}-\text{H}$, the dinuclear platinum(II) di-yne, $[(\text{Ph})(\text{PEt}_3)_2\text{Pt}-\text{C}\equiv\text{C}-\text{R}-\text{C}\equiv\text{C}-\text{Pt}(\text{PEt}_3)_2(\text{Ph})]$, and the platinum(II) poly-yne, *trans*- $[(^n\text{Bu}_3\text{P})_2\text{Pt}-\text{C}\equiv\text{C}-\text{R}-\text{C}\equiv\text{C}-]_\infty$ ($\text{R} = \text{thieno}[3,2-*b*]-\text{thiophene-2,5-diy}$ l). Rigid-rod platinum(II) poly-ynes with the general formula *trans*- $[(^n\text{Bu}_3\text{P})_2\text{Pt}-\text{C}\equiv\text{C}-\text{R}-\text{C}\equiv\text{C}-]_\infty$ ($\text{R} = \text{conjugated aromatic/heteroaromatic linker group}$) are considered to be good model systems to study the triplet excited state in polymers and provide important information on the photophysical processes that occur within them (Khan, Al-Mandhary, Al-Suti, Hisham *et al.*, 2002; Khan, Al-Mandhary, Al-Suti, Feeder *et al.*, 2002; Khan *et al.*, 2003). The incorporation of heavy transition metals, such as platinum, at regular intervals along the rigid polymer backbone introduces a large component of spin-orbit coupling that allows emission from the triplet excited state of the system *via* spin cross-over processes (Wittmann *et al.*, 1994; Beljonne *et al.*, 1996; Younus *et al.*, 1998; Chawdhury *et al.*, 1999). The novel photophysics of the platinum(II) poly-ynes leads to materials that are useful for applications in modern opto-electronic devices such as light-emitting diodes (LEDs), lasers, photocells and field-effect transistors (FETs) (Wilson *et al.*, 2000; Wilson, Chawdhury *et al.*, 2001; Wilson, Dhoot *et al.*, 2001).



The title compound, (I), crystallizes in the monoclinic space group $P2_1/n$, with one molecule in the asymmetric unit, so that there is no crystallographically imposed symmetry (Fig. 1). The central dithieno[3,2-*b*:2',3'-*d*]thiophene group is essentially planar and the bond parameters associated with this group are similar to those found in other materials containing

Received 7 June 2004

Accepted 14 June 2004

Online 26 June 2004

this thiophene ring system (Li *et al.*, 1998; Osterod *et al.*, 2001; Frey *et al.*, 2002). The bond parameters for the acetylene groups and the trimethylsilyl ligands are similar to those observed in related compounds (Khan, Ahrens *et al.*, 2002; Khan *et al.*, 2004).

There are a number of significant intermolecular interactions within the crystal structure. All three of the unique S atoms show contacts to symmetry-related S atoms: S1...S2 (related by the symmetry operation $x - 1, y, z$) at a distance of 3.338 (2) Å; S2...S1 (related by $1 + x, y, z$) at a distance of 3.338 Å; S2...S2 (related by $2 + x, -y, -z$) at a distance of 3.545 (2) Å; the S3...S2 (related by $x - 1, y, z$) distance is 3.404 (2) Å, which is the shortest. There is also an indication of the presence of π -stacking between pairs of adjacent dithieno[3,2-*b*:2',3'-*d*]thiophene ring systems, with centroid-centroid separations between the five-membered S1-containing ring and the S3-containing ring, related by the symmetry operation $1 - x, -y, -z$, of 4.328 (5) Å, and the S2-containing ring and its symmetry-related partner (again by $1 - x, -y, -z$) with a distance of 4.539 (5) Å.

Experimental

5,5'-Dibromodithieno[3,2-*b*:2',3'-*d*]thiophene (2.0 g, 5.64 mmol), trimethylsilylethyne (1.46 g, 14.9 mmol) and ¹Pr₂NH–THF (70 ml, 1:1 *v/v*) were mixed with catalytic amounts of CuI (20 mg), Pd(OAc)₂ (20 mg) and PPh₃ (60 mg). The crude product was worked up to yield a dark-brown residue, which was then applied to a silica column in hexane and eluted with the same solvent. The title compound was obtained as a colourless crystalline solid in 78% isolated yield (1.72 g).

Crystal data

C ₁₈ H ₂₀ S ₃ Si ₂	Synchrotron radiation,
<i>M_r</i> = 388.7	$\lambda = 0.6887$ Å
Monoclinic, <i>P</i> ₂ ₁ / <i>n</i>	Cell parameters from 11495
<i>a</i> = 6.191 (2) Å	reflections
<i>b</i> = 28.671 (9) Å	$\theta = 3.2$ – 29.4°
<i>c</i> = 12.066 (4) Å	$\mu = 0.46$ mm ^{−1}
$\beta = 92.51$ (2)°	<i>T</i> = 150 (2) K
<i>V</i> = 2139.7 (13) Å ³	Plate, yellow
<i>Z</i> = 4	0.07 × 0.05 × 0.01 mm
<i>D_x</i> = 1.207 Mg m ^{−3}	

Data collection

Bruker AXS SMART 1K CCD diffractometer	3062 independent reflections
Thin-slice ω scans	2494 reflections with $I > 2\sigma(I)$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	<i>R</i> _{int} = 0.063
<i>T</i> _{min} = 0.969, <i>T</i> _{max} = 0.996	$\theta_{\text{max}} = 22.5^\circ$
11094 measured reflections	<i>h</i> = −6 → 6
	<i>k</i> = −28 → 31
	<i>l</i> = −13 → 13

Refinement

Refinement on <i>F</i> ²	$w = 1/[\sigma^2(F_o^2) + (0.0415P)^2 + 1.5269P]$
$R[F^2 > 2\sigma(F^2)] = 0.055$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.114$	$(\Delta/\sigma)_{\text{max}} < 0.001$
<i>S</i> = 1.17	$\Delta\rho_{\text{max}} = 0.40$ e Å ^{−3}
3062 reflections	$\Delta\rho_{\text{min}} = -0.33$ e Å ^{−3}
224 parameters	
H-atom parameters constrained	

The crystal, a small thin plate, did not diffract significantly beyond 45° in 2 θ and the final refinement θ_{max} was limited to 22.5°. One of the methyl groups associated with Si1 showed positional disorder over




Figure 1
View of (I) (50% probability displacement ellipsoids). The disordered methyl group is not shown for clarity.

two sites, C13 and C13A. These two atoms, and their associated H atoms, were refined with occupancies fixed at 50% each. All the aromatic and methyl H atoms were constrained as riding atoms, fixed to the parent atoms with distances of 0.95 and 0.98 Å, respectively, and *U*(H) = 1.2 (aromatic H atoms) and 1.5 (methyl H atoms) times *U*_{eq}(parent atom).

Data collection: SMART (Bruker, 1998); cell refinement: LSCCELL (Clegg, 1997); data reduction: SAINT (Bruker, 2000); program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

We are grateful to the Sultan Qaboos University, Oman, the Royal Society of Chemistry for a Journals Grant for International Authors (to MSK), and the DAAD (to BA) for funding.

References

- Beljonne, D., Wittmann, H. F., Köhler, A., Graham, S., Younus, M., Lewis, J., Raithby, P. R., Khan, M. S., Friend, R. H. & Bredas, J. L. (1996). *J. Chem. Phys.* **105**, 3868–3877.
- Bruker (1998). SMART. Version 5.054. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2000). SADABS (Version 6.02a) and SAINT (Version 6.02a). Bruker AXS Inc., Madison, Wisconsin, USA.
- Chawdhury, N., Köhler, A., Friend, R. H., Wong, W.-Y., Younus, M., Raithby, P. R., Lewis, J., Corcoran, T. C., Al-Mandhary, M. R. A. & Khan, M. S. (1999). *J. Chem. Phys.* **110**, 4963–4970.
- Clegg, W. (1997). LSCCELL. University of Newcastle upon Tyne, England.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Frey, J., Bond, A. D. & Holmes, A. B. (2002). *Chem. Commun.* pp. 2424–2425.
- Khan, M. S., Ahrens, B., Male, L. & Raithby, P. R. (2002). *Acta Cryst.* **E58**, o1220–o1221.
- Khan, M. S., Ahrens, B., Male, L. & Raithby, P. R. (2004). *Acta Cryst.* **E60**, o915–o916.
- Khan, M. S., Al-Mandhary, M. R. A., Al-Suti, M. K., Feeder, N., Nahar, S., Köhler, A., Friend, R. H., Wilson, P. J. & Raithby, P. R. (2002). *J. Chem. Soc. Dalton Trans.* pp. 2441–2448.
- Khan, M. S., Al-Mandhary, M. R. A., Al-Suti, M. K., Hisham, A. K., Raithby, P. R., Ahrens, B., Mahon, M. F., Male, L., Marseglia, E. A., Tedesco, E., Friend, R. H., Köhler, A., Feeder, N. & Teat, S. J. (2002). *J. Chem. Soc. Dalton Trans.* pp. 1358–1368.
- Khan, M. S., Al-Mandhary, M. R. A., Al-Suti, M. K., Raithby, P. R., Ahrens, B., Male, L., Friend, R. H., Köhler, A. & Wilson, J. S. (2003). *Dalton Trans.* pp. 65–73.
- Li, X.-C., Sirringhaus, H., Garnier, F., Holmes, A. B., Moratti, S. C., Feeder, N., Clegg, W., Teat, S. J. & Friend, R. H. (1998). *J. Am. Chem. Soc.* **120**, 2206–2207.
- Osterod, F., Peters, L., Kraft, A., Sano, T., Morrison, J. J., Feeder, N. & Holmes, A. B. (2001). *J. Mater. Chem.* **11**, 1625–1633.

- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Wilson, J. S., Chawdhury, N., Köhler, A., Friend, R. H., Al-Mandhary, M. R. A., Khan, M. S., Younus, M. & Raithby, P. R. (2001). *J. Am. Chem. Soc.* **123**, 9412–9417.
- Wilson, J. S., Dhoot, A. S., Seeley, A. J. A. B., Khan, M. S., Köhler, A. & Friend, R. H. (2001). *Nature (London)*, **413**, 828–831.
- Wilson, J. S., Köhler, A., Friend, R. H., Al-Suti, M. K., Khan, M. S. & Raithby, P. R. (2000). *J. Chem. Phys.* **113**, 7627–7634.
- Wittmann, H. F., Friend, R. H., Khan, M. S. & Lewis, J. (1994). *J. Chem. Phys.* **101**, 2693–2698.
- Younus, M., Köhler, A., Cron, A., Chawdhury, N., Al-Mandhary, M. R. A., Khan, M. S., Lewis, J., Long, N. J., Friend, R. H. & Raithby, P. R. (1998). *Angew. Chem. Int. Ed.* **37**, 3036–3039.