

# 1,4-Phenylene-bis-(3,4-dichloro-5-phenyl-carbamoyl-1*H*-pyrrole-2-carboxamide) bis(tetrabutylammonium chloride) acetonitrile disolvate

Mark E. Light,\* Philip A. Gale and Korakot Navakhun

School of Chemistry, University of Southampton, Highfield, Southampton, England SO17 1BJ

Correspondence e-mail: light@soton.ac.uk

The title compound,  $C_{30}H_{20}N_6O_4Cl_4 \cdot 2C_{16}H_{36}N^+ \cdot 2Cl^- \cdot 2C_2H_3N$ , contains two hydrogen-bonded chloride anions bound to 1,4-phenylene-bis-(3,4-dichloro-5-phenylcarbamoyl-1*H*-pyrrole-2-carboxamide) as the tetrabutylammonium salt. There is also a short pyrrolic hydrogen bond ( $N \cdots Cl = 3.068(3) \text{ \AA}$ ), and two longer ones to the amino H atoms [ $N \cdots Cl = 3.269(3) \text{ \AA}$  and  $3.275(3) \text{ \AA}$ ]. The neutral molecule lies on an inversion centre situated at the centre of the central benzene ring.

Received 22 March 2005

Accepted 6 April 2005

Online 9 April 2005


## Key indicators

Single-crystal X-ray study  
 $T = 120 \text{ K}$   
Mean  $\sigma(C-C) = 0.005 \text{ \AA}$   
Disorder in solvent or counterion  
 $R$  factor = 0.060  
 $wR$  factor = 0.144  
Data-to-parameter ratio = 18.0

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

## Comment

1,4-Phenylene-bis-(3,4-dichloro-5-phenylcarbamoyl-1*H*-pyrrole-2-carboxamide) crystallizes from tetrabutylammonium chloride acetonitrile solution as a tetrabutylammonium chloride acetonitrile solvate, (I).



The receptor adopts an S-shaped conformation around a centre of inversion with one chloride bound on each side. The pyrrole and terminal benzene ring pairs are coplanar, and the angle between the central and terminal benzene rings is  $32.02(4)^\circ$ . Of the three hydrogen bonds to the chloride, the pyrrolic one is the shortest, with an  $N \cdots Cl$  distance of  $3.068(3) \text{ \AA}$ , whilst the two either side are longer, with distances of  $3.269(3) \text{ \AA}$  and  $3.275(3) \text{ \AA}$ .

## Experimental

The receptor molecule, 1,4-phenylene-bis-(3,4-dichloro-5-phenyl-carbamoyl-1*H*-pyrrole-2-carboxamide), (1), was synthesized according to literature methods (Gale *et al.*, 2002). Crystals of the acetonitrile solvate of the tetrabutylammonium chloride complex were grown by slow evaporation of an acetonitrile solution of (1) in acetonitrile in the presence of excess tetrabutylammonium chloride.

### Crystal data

$C_{30}H_{20}N_6O_4Cl_4 \cdot 2C_{16}H_{36}N^+ \cdot 2Cl^- \cdot 2C_2H_3N$   
 $M_r = 1308.24$   
 Monoclinic,  $P2_1/c$   
 $a = 8.5720$  (2) Å  
 $b = 21.1088$  (5) Å  
 $c = 19.3520$  (6) Å  
 $\beta = 93.5560$  (10)° **precision OK?**  
 $V = 3494.90$  (16) Å<sup>3</sup>  
 $Z = 2$

$D_x = 1.243$  Mg m<sup>-3</sup>  
 Mo  $K\alpha$  radiation  
 Cell parameters from 35384 reflections  
 $\theta = 2.9$ – $26.4$ °  
 $\mu = 0.30$  mm<sup>-1</sup>  
 $T = 120$  (2) K  
 Needle, colourless  
 $0.15 \times 0.07 \times 0.05$  mm

### Data collection

Bruker–Nonius KappaCCD area-detector diffractometer  
 $\varphi$  and  $\omega$  scans  
 Absorption correction: multi-scan (SORTAV; Blessing, 1997)  
 $T_{min} = 0.906$ ,  $T_{max} = 0.990$   
 13653 measured reflections

7107 independent reflections  
 4902 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.046$   
 $\theta_{max} = 26.4$ °  
 $h = -10 \rightarrow 10$   
 $k = -26 \rightarrow 25$   
 $l = -24 \rightarrow 24$

### Refinement

Refinement on  $F^2$   
 $R[F^2 > 2\sigma(F^2)] = 0.060$   
 $wR(F^2) = 0.144$   
 $S = 1.11$   
 7107 reflections  
 395 parameters  
 H-atom parameters constrained


$w = 1/[\sigma^2(F_o^2) + (0.0389P)^2 + 4.3109P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{max} = 0.001$   
 $\Delta\rho_{max} = 0.70$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -1.05$  e Å<sup>-3</sup>  
 Extinction correction: SHELXL97  
 Extinction coefficient: 0.0023 (4)

**Table 1**

Hydrogen-bonding geometry (Å, °).

| $D-H \cdots A$      | $D-H$ | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|---------------------|-------|--------------|--------------|----------------|
| N1–H91 $\cdots$ Cl3 | 0.88  | 2.40         | 3.269 (3)    | 171.6          |
| N2–H92 $\cdots$ Cl3 | 0.88  | 2.20         | 3.068 (3)    | 166.8          |
| N3–H93 $\cdots$ Cl3 | 0.88  | 2.40         | 3.275 (3)    | 171.2          |

H atoms were identified in a difference map and then placed in calculated positions (N–H 0.88, aromatic C–H 0.95, methylene C–H 0.99, methyl C–H 0.98) and refined using a riding model [ $U_{iso}(H) = 1.2$  or  $1.5$  times  $U_{eq}(\text{parent atom})$ ]. One arm of the tetrabutyl-



**Figure 1**

Structure of the title compound, with displacement ellipsoids drawn at the 35% probability level and non-acidic H atoms omitted for clarity. Both disorder components are shown.

ammonium is disordered. It has been modelled as split over two possible orientations with one third and two thirds occupancy. C–C and C–N distances were restrained to standard values and the displacement parameters of split atom pairs were constrained to be equal. The deepest hole is located 1.28 Å from C9.

Data collection: DENZO (Otwinowski & Minor, 1997) and COLLECT (Hooft, 1998); cell refinement: DENZO and COLLECT; data reduction: DENZO and COLLECT; program(s) used to solve structure: SHELXS97 (Sheldrick, 1990); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: CAMERON (Watkin *et al.*, 1993); software used to prepare material for publication: WinGX (Farrugia, 1999).

The authors thank the EPSRC for funding the crystallographic facilities. PAG thanks the Royal Society for a University Research Fellowship and Universities UK for an ORS studentship to KN.

## References

- Blessing, R. H. (1997). *J. Appl. Cryst.* **30**, 421–426.  
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
 Gale, P. A., Navakhun, K., Camiolo, S., Light, M. E. & Hursthouse, M. B. (2002). *J. Am. Chem. Soc.* **124**, 11228–11229.  
 Hooft, R. (1998). COLLECT. Nonius BV, The Netherlands.  
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr and R. M. Sweet, pp. 307–326. New York: Academic Press.  
 Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.  
 Sheldrick, G. M. (1997). SHELXL97. University of Göttingen, Germany.  
 Watkin, D. M., Pearce, L. & Prout, C. K. (1993). CAMERON. University of Oxford, England.