

2-Benzyl-1,3-diphenylpropane-1,3-dione

Nenad Judaš* and Branko Kaitner

Laboratory of General and Inorganic Chemistry,
Chemistry Department, Faculty of Science,
University of Zagreb, Horvatovac 102a,
HR-10000 Zagreb, Croatia

Correspondence e-mail: judas@chem.pmf.hr


Key indicators

Single-crystal X-ray study
 $T = 298\text{ K}$
Mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$
 R factor = 0.051
 wR factor = 0.152
Data-to-parameter ratio = 17.2For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.

The molecule of 2-benzyl-1,3-diphenylpropane-1,3-dione, $\text{C}_{22}\text{H}_{18}\text{O}_2$, being a diketo tautomer, shows different stereochemistry from that of the closed-ring *cis*-ketoenol tautomer of the parent 1,3-diphenylpropane-1,3-dione. The molecular packing involves $\text{C}-\text{H}\cdots\text{O}$ and $\pi-\pi$ stacking interactions.

Comment

Insertion of a sterically demanding substituent into the α -position of a β -diketone molecule affects its ligating properties and the keto–enol equilibrium. As part of a study of the effects of benzyl substitution on the structures of β -diketones, we have synthesized an α -substituted derivative of 1,3-diphenylpropane-1,3-dione, Hdbm. The title compound, 2-benzyl-1,3-diphenylpropane-1,3-dione, $\alpha(\text{Bzl})\text{Hdbm}$, (I), is a potential bidentate monoanionic and neutral ligand for coordination to transition metal ions.



The asymmetric unit of (I) contains a single molecule which is located in a general position (Fig. 1) and which is best described as a distorted *U*; *cis*-diketo; *Z*, *Z* conformer (Emsley, 1984). Distortion of the molecule with respect to the ideal *U*-conformation involving atoms O1, C1, C2, C3 and O2 is best described in terms of torsion angles involving sets of atoms O1/C1/C2/C3 and O2/C3/C2/C1. The angles are $-90.42(18)^\circ$ and $25.69(23)^\circ$, respectively. Such a conformation of the molecule is dictated by steric hindrance related to the presence of the large benzyl group, which prevents the molecule from adopting the closed-ring conformation characteristic for the *cis*-ketoenol tautomer.

The bond angles about atom C2, to which two benzoyl and one benzyl groups are bonded, do not show any significant deviations from those expected for an sp^3 -hybridized C atom. A larger discrepancy from the standard tetrahedral value is observed in the angle C2–C4–C5 [$114.7(2)^\circ$] and may be explained by steric effects. Notably, similar deviations were observed in the analogous compound 3,3-dibenzylpentane-2,4-dione, $\alpha(\text{Bzl}, \text{Bzl})\text{acac}$ (Judaš & Kaitner, 1995).

Received 29 September 2005

Accepted 31 October 2005

Online 5 November 2005




Figure 1
The molecular structure of (I), with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level.

The planes defined by the terminal phenyl rings (C11–C16 and C17–C22) are twisted with respect to the planes defined by neighbouring carbonyl groups and the C_α atom (O1/C1/C2 and O2/C3/C2) by 19.56 (17) and 26.97 (8)°, respectively. Such values are larger than those observed in any known polymorph of the parent compound 1,3-diphenylpropane-1,3-dione (Hdbm) (Williams, 1966; Hollander *et al.*, 1973; Kaitner & Meštrović, 1993).

The molecules of (I) are connected by C–H···O interactions (Table 1). Four such contacts, specifically C10–H10···O2ⁱ, C2–H2···O2ⁱ, C22–H22···O2ⁱ and C21–H21···O1ⁱ, cooperatively link the molecules into chains (Fig. 2; symmetry code as in Table 1). These chains of molecules are further linked into sheets by way of two distinct sets of C–H···O interactions (Fig. 3). The sheets are interlinked by additional C–H···O interactions and are presumably stabilized by π – π stacking interactions between the terminal phenyl rings at a distance of 3.55 Å.

Experimental

The title compound was synthesized from 1,3-diphenylpropane-1,3-dione and benzyl bromide *via* the method of Kalyanam *et al.* (1979). The resulting crude product was recrystallized twice from hot ethanol. Thermal analysis (thermogravimetric analysis and differential scanning calorimetry) of the pure product were carried out using Mettler thermal analysis modules. The calorimetric thermogram exhibited two endothermic signals. The first signal is the sharper and stronger of the two and corresponds to the melting. The onset temperature of the first signal is $T_f = 373$ K with enthalpy of fusion $\Delta H_{fus} = 35.6$ kJ mol⁻¹. The second signal, related to boiling, has an onset temperature of $T_b = 583$ K. No degradation of the sample during melting and boiling processes was detectable by thermal analysis methods. Single crystals of (I) suitable for the diffraction experiment were obtained by slow evaporation of a solution of the compound in acetic acid.




Figure 2
Ball-and-stick model representation showing the cooperative linking of the α (Bzl)Hdbm molecules of (I) into chains. C–H···O interactions are shown as dashed lines.




Figure 3
Ball-and-stick model representation showing the linking of the chains into sheets *via* C–H···O interactions (dashed lines).

Crystal data

$C_{22}H_{18}O_2$	$D_x = 1.221$ Mg m ⁻³
$M_r = 314.36$	Mo $K\alpha$ radiation
Monoclinic, $P2_1/n$	Cell parameters from 44 reflections
$a = 10.2935$ (6) Å	$\theta = 10.0$ – 16.6°
$b = 10.9060$ (6) Å	$\mu = 0.08$ mm ⁻¹
$c = 15.4090$ (7) Å	$T = 298$ (1) K
$\beta = 98.562$ (4)°	Prism, colourless
$V = 1710.55$ (16) Å ³	$0.91 \times 0.80 \times 0.60$ mm
$Z = 4$	

Data collection

Philips PW1100 diffractometer	$\theta_{max} = 27.0^\circ$
ω scans	$h = -13 \rightarrow 13$
Absorption correction: none	$k = 0 \rightarrow 13$
3866 measured reflections	$l = 0 \rightarrow 19$
3733 independent reflections	3 standard reflections
2731 reflections with $I > 2\sigma(I)$	frequency: 60 min
$R_{int} = 0.065$	intensity decay: 2.8%

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.079P)^2 + 0.4571P]$
$R[F^2 > 2\sigma(F^2)] = 0.051$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.152$	$(\Delta/\sigma)_{max} < 0.001$
$S = 1.03$	$\Delta\rho_{max} = 0.26$ e Å ⁻³
3733 reflections	$\Delta\rho_{min} = -0.17$ e Å ⁻³
217 parameters	
H-atom parameters constrained	

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C10—H10 \cdots O2 ⁱ	0.93	2.63	3.498 (2)	155
C2—H2 \cdots O2 ⁱ	0.98	2.60	3.576 (2)	173
C22—H22 \cdots O2 ⁱ	0.93	2.58	3.513 (2)	177
C21—H21 \cdots O1 ⁱ	0.93	2.74	3.491 (3)	139
C6—H6 \cdots O1 ⁱⁱ	0.93	2.72	3.401 (2)	131

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + 1, -y + 1, -z$.

The positions of all H atoms were calculated, with C—H distances of 0.93 for phenyl, 0.97 for methylene and 0.98 Å for methine H atoms. The H atoms were included in the refinement using the riding-model approximation, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Data collection: *STADIA* (Stoe & Cie, 1995); cell refinement: *STADIA*; data reduction: *X-RED* (Stoe & Cie, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997), *MERCURY* (Version 1.2.1; Bruno *et al.*, 2002), *RasTop* (Valadon, 2004) and *POV-*

Ray (Version 3.6; Persistence of Vision, 2004); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This work was supported by the Ministry of Science and Technology, Croatia.

References

- Bruno, I. J., Cole, J. C., Edgington, P. R., Kessler, M., Macrae, C. F., McCabe, P., Pearson, J. & Taylor, R. (2002). *Acta Cryst.* **B58**, 389–397.
- Emsley, J. (1984). *Struct. Bonding (Berlin)*, **57**, 147–192.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hollander, F. J., Templeton, D. H. & Zalkin, A. (1973). *Acta Cryst.* **B29**, 1552–1553.
- Judaš, N. & Kaitner, B. (1995). *Acta Cryst.* **C51**, 2123–2125.
- Kaitner, B. & Meštrović, E. (1993). *Acta Cryst.* **C49**, 1523–1525.
- Kalyanam, N., Karban, J. W. & McAtee, J. L. Jr (1979). *Org. Prep. Proced. Int.* **11**, 100–101.
- Persistence of Vision (2004). *POV-Ray*. Version 3.6. Persistence of Vision Pty. Ltd., Victoria, Australia. <http://www.povray.org/>
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Stoe & Cie (1995). *STADIA* (Version 1.05b) and *X-RED* (Version 1.05). Stoe & Cie, Darmstadt, Germany.
- Valadon, P. (2004). *RasTop*. Version 2.0.3. Philippe Valadon, PO Box 928064, La Jolla, CA 92192-8064, USA. <http://www.geneinfinity.org/rastop/>
- Williams, D. E. (1966). *Acta Cryst.* **21**, 340–349.