

catena-Poly[[diaquabis[(4-chlorophenylsulfinyl)acetato- κ O]zinc(II)]- μ -4,4'-bipyridine- κ^2 N:N']

Yan-Jun Hou, Bai-Yan Li, Ying-Hui Yu, Zhi-Zhong Sun and Guang-Feng Hou*

College of Chemistry and Materials Science, Heilongjiang University, Harbin

150080, People's Republic of China

Correspondence e-mail: hgf1000@163.com


Received 22 May 2007; accepted 24 May 2007

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.004$ Å; R factor = 0.024; wR factor = 0.062; data-to-parameter ratio = 16.7.

In the title coordination polymer, $[Zn(C_8H_6ClO_3S)_2(C_{10}H_8N_2)(H_2O)_2]_n$, the Zn^{II} atom exists in an octahedral coordination environment that is formed by two carboxylate O atoms from two (4-chlorophenylsulfinyl)acetate ligands, two N atoms from two bipyridine ligands and two water molecules. The Zn^{II} atom lies on a twofold axis. Bridging by the bipyridine ligand leads to a linear chain structure and intermolecular O–H···O hydrogen bonds link the chains into a three-dimensional network.

Related literature

For the isostructural cobalt compound, see Hou *et al.* (2007b). For related literature, see: Hou *et al.* (2007a); Nobles & Thompson (1965).



Experimental

Crystal data

$[Zn(C_8H_6ClO_3S)_2(C_{10}H_8N_2)(H_2O)_2]$
 $M_r = 692.86$
Orthorhombic, $Fdd2$
 $a = 20.194 (5)$ Å
 $b = 25.528 (8)$ Å
 $c = 11.458 (6)$ Å

$V = 5907 (4)$ Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 1.20$ mm⁻¹
 $T = 293 (2)$ K
 $0.27 \times 0.25 \times 0.19$ mm

Data collection

Rigaku R-AXIS RAPID diffractometer
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)
 $T_{min} = 0.738$, $T_{max} = 0.804$

14000 measured reflections
3142 independent reflections
2962 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$
 $wR(F^2) = 0.062$
 $S = 1.06$
3142 reflections
188 parameters
1 restraint

H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.28$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³
Absolute structure: Flack (1983),
from 1371 Friedel pairs
Flack parameter: 0.017 (8)

Table 1
Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
O4–H12···O1 ⁱ	0.85	1.88	2.706 (2)	164
O4–H12···S1 ⁱ	0.85	2.99	3.8217 (18)	165
O4–H11···O2 ⁱⁱ	0.85	1.85	2.657 (2)	158

Symmetry codes: (i) $x + \frac{1}{4}, -y + \frac{7}{4}, z - \frac{1}{4}$; (ii) $-x, -y + 2, z$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Molecular Structure Corporation & Rigaku, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997a); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997a); molecular graphics: *SHELXTL* (Sheldrick, 1997b); software used to prepare material for publication: *SHELXL97*.

The authors thank the Heilongjiang University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2270).

References

- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Hou, Y.-J., Yu, Y.-H., Sun, Z.-Z., Li, B.-Y. & Hou, G.-F. (2007a). *Acta Cryst. E* **63**, m1530.
- Hou, Y.-J., Yu, Y.-H., Sun, Z.-Z., Li, B.-Y. & Hou, G.-F. (2007b). *Acta Cryst. E* **63**, m1560.
- Molecular Structure Corporation & Rigaku (2002). *CrystalStructure*, Version 3.10. MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
- Nobles, W. L. & Thompson, B. B. (1965). *J. Pharm. Sci.* **54**, 709–713.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (1997a). *SHELXL97* and *SHELXS97*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997b). *SHELXTL*. Version 5.10. Bruker AXS Inc., Madison, Wisconsin, USA.

supplementary materials

Acta Cryst. (2007). E63, m1838 [doi:10.1107/S1600536807025299]

catena-Poly[[diaquabis[(4-chlorophenylsulfinyl)acetato- κO]zinc(II)]- μ -4,4'bipyridine- $\kappa^2 N:N'$]

Y.-J. Hou, B.-Y. Li, Y.-H. Yu, Z.-Z. Sun and G.-F. Hou

Comment

Recently, we reported the crystal structure of diaquabis[(4-nitrophenylsulfinylacetato)(4,4'-bipyridine)zinc (Hou *et al.* 2007a). We also reported that of diaquabis[(4-chlorophenylsulfinylacetato)(4,4'-bipyridine)cobalt (Hou *et al.* 2007b); this paper reports the isostructural zinc compound.

The title compound has the zinc bis(4-chlorophenylsulfinylacetate) bridged by 4,4'-bipyridine into a linear chain (Fig. 1). The Zn^{II} atom shows an all *trans* octahedral coordination. The chains are connected into a three dimensional network *via* intermolecular O—H···O hydrogen bonds (Table 1), (Fig. 2).

Experimental

(4-Chlorophenylsulfanyl)acetic acid was prepared by the nucleophilic reaction of chloroacetic acid and 4-chlorothiophenol under basic conditions. It was then oxidized using 30% aqueous hydrogen peroxide in acetic anhydride solution to produce 4-chlorophenylsulfinyl acetic acid (Nobles & Thompson, 1965). Zinc nitrate hexahydrate (0.592 g, 2 mmol), 4,4'-bipyridine (0.312 g, 2 mmol) and 4-chlorophenylsulfinyl acetic acid (0.437 g, 2 mmol) were dissolved in water and the pH was adjusted to 6 with 0.01 M sodium hydroxide; colorless crystals separated from the filtered solution after several days.

Refinement

H atoms bound to C atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.93 Å (aromatic C) or C—H = 0.97 Å (methylene C), and with $U_{iso}(H) = 1.2U_{eq}(C)$. Water H atoms were initially located in a difference Fourier map but they were treated as riding on their parent atoms with O—H = 0.85 Å and with $U_{iso}(H) = 1.5U_{eq}(O)$.

The Flack parameter was refined from 1371 Friedel pairs.

Figures




Fig. 1. Part of the polymeric structure of the title complex, with the atomlabelling scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are represented as spheres of arbitrary radii. [Symmetry codes: (I) $-x, -y + 2, z$; (II) $x, y, z - 1$; (III) $-x, -y + 2, z - 1$, (IV) $x, y, z + 1$].

supplementary materials




Fig. 2. A partial packing plot of (I). Dashed lines indicate the hydrogen-bonding interactions. H atoms not involved in hydrogen bonds have been omitted.

catena-Poly[[diaquabis[(4-chlorophenylsulfinyl)acetato- κO] zinc(II)]- μ -4,4'-bipyridine- $\kappa^2 N:N'$]

Crystal data

[Zn(C ₈ H ₆ ClO ₃ S) ₂ (C ₁₀ H ₈ N ₂)(H ₂ O) ₂]	$F_{000} = 2832$
$M_r = 692.86$	$D_x = 1.558 \text{ Mg m}^{-3}$
Orthorhombic, $Fdd2$	Mo $K\alpha$ radiation
Hall symbol: F 2 -2d	$\lambda = 0.71073 \text{ \AA}$
$a = 20.194 (5) \text{ \AA}$	Cell parameters from 12841 reflections
$b = 25.528 (8) \text{ \AA}$	$\theta = 6.3\text{--}54.9^\circ$
$c = 11.458 (6) \text{ \AA}$	$\mu = 1.20 \text{ mm}^{-1}$
$V = 5907 (4) \text{ \AA}^3$	$T = 293 (2) \text{ K}$
$Z = 8$	Block, colorless
	$0.27 \times 0.25 \times 0.19 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID diffractometer	3142 independent reflections
Radiation source: fine-focus sealed tube	2962 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.024$
$T = 293(2) \text{ K}$	$\theta_{\max} = 27.5^\circ$
ω scans	$\theta_{\min} = 3.2^\circ$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$h = -26 \rightarrow 26$
$T_{\min} = 0.738$, $T_{\max} = 0.804$	$k = -33 \rightarrow 31$
14000 measured reflections	$l = -14 \rightarrow 12$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.024$	$w = 1/[\sigma^2(F_o^2) + (0.0385P)^2 + 0.7396P]$
$wR(F^2) = 0.062$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\max} = 0.001$
3142 reflections	$\Delta\rho_{\max} = 0.28 \text{ e \AA}^{-3}$
	$\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$

188 parameters	Extinction correction: none
1 restraint	Absolute structure: Flack (1983), from 1371 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.017 (8)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.02754 (11)	0.72498 (7)	0.2397 (2)	0.0334 (5)
C2	0.06452 (12)	0.70605 (8)	0.1464 (2)	0.0413 (5)
H1	0.0606	0.7211	0.0729	0.050*
C3	0.10706 (13)	0.66457 (8)	0.1647 (2)	0.0487 (6)
H2	0.1336	0.6521	0.1045	0.058*
C4	0.10952 (13)	0.64206 (9)	0.2733 (3)	0.0505 (6)
C5	0.07313 (15)	0.66025 (11)	0.3662 (3)	0.0561 (7)
H3	0.0760	0.6444	0.4391	0.067*
C6	0.03208 (14)	0.70286 (10)	0.3482 (2)	0.0485 (6)
H4	0.0076	0.7165	0.4099	0.058*
C7	0.02983 (10)	0.82874 (7)	0.1969 (2)	0.0374 (4)
H5	0.0576	0.8306	0.2658	0.045*
H6	0.0579	0.8213	0.1302	0.045*
C8	-0.00572 (11)	0.88119 (7)	0.17876 (17)	0.0337 (5)
C9	0.01985 (13)	1.04164 (8)	0.4480 (2)	0.0437 (5)
H7	0.0347	1.0708	0.4071	0.052*
C10	0.01934 (16)	1.04353 (10)	0.5687 (2)	0.0489 (6)
H8	0.0319	1.0739	0.6075	0.059*
C11	0.0000	1.0000	0.6309 (3)	0.0347 (8)
C12	0.0000	1.0000	0.7607 (3)	0.0380 (8)
C13	0.03379 (13)	0.96210 (9)	0.8246 (2)	0.0416 (5)
H9	0.0567	0.9357	0.7861	0.050*
C14	0.03333 (11)	0.96369 (8)	0.9447 (2)	0.0405 (5)
H10	0.0572	0.9384	0.9852	0.049*
Cl1	0.16118 (5)	0.58797 (3)	0.29340 (12)	0.0958 (3)
N1	0.0000	1.0000	0.3880 (2)	0.0313 (6)
N2	0.0000	1.0000	1.0064 (2)	0.0342 (6)

supplementary materials

O1	-0.06247 (10)	0.78794 (6)	0.32997 (18)	0.0598 (5)
O2	-0.06317 (9)	0.88029 (6)	0.14055 (18)	0.0539 (5)
O3	0.02837 (7)	0.92075 (5)	0.20348 (13)	0.0352 (3)
O4	0.10101 (7)	1.02337 (5)	0.19763 (13)	0.0378 (3)
H12	0.1242	1.0061	0.1491	0.057*
H11	0.0997	1.0558	0.1805	0.057*
S1	-0.03021 (3)	0.777363 (17)	0.21386 (5)	0.03840 (13)
Zn1	0.0000	1.0000	0.198249 (18)	0.02787 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0359 (11)	0.0260 (8)	0.0384 (13)	-0.0028 (7)	0.0043 (8)	0.0004 (8)
C2	0.0475 (13)	0.0375 (10)	0.0390 (11)	0.0028 (9)	0.0112 (9)	0.0060 (9)
C3	0.0496 (14)	0.0409 (11)	0.0557 (14)	0.0050 (10)	0.0141 (11)	-0.0017 (11)
C4	0.0435 (13)	0.0381 (11)	0.0700 (17)	0.0048 (10)	-0.0043 (11)	0.0100 (11)
C5	0.0664 (18)	0.0572 (14)	0.0446 (14)	0.0015 (13)	0.0001 (12)	0.0200 (12)
C6	0.0589 (15)	0.0509 (13)	0.0359 (12)	0.0054 (11)	0.0088 (10)	0.0043 (11)
C7	0.0373 (10)	0.0271 (8)	0.0478 (12)	-0.0029 (7)	0.0039 (10)	-0.0011 (9)
C8	0.0440 (12)	0.0287 (8)	0.0282 (14)	0.0033 (8)	0.0010 (9)	-0.0018 (8)
C9	0.0696 (15)	0.0359 (10)	0.0255 (11)	-0.0108 (10)	-0.0049 (10)	0.0047 (9)
C10	0.0792 (18)	0.0401 (12)	0.0274 (12)	-0.0121 (12)	-0.0103 (11)	-0.0014 (10)
C11	0.046 (2)	0.0390 (19)	0.0190 (18)	0.0008 (12)	0.000	0.000
C12	0.045 (2)	0.043 (2)	0.027 (2)	-0.0058 (13)	0.000	0.000
C13	0.0551 (14)	0.0442 (12)	0.0255 (11)	0.0101 (10)	0.0047 (9)	-0.0012 (9)
C14	0.0498 (13)	0.0422 (11)	0.0295 (12)	0.0099 (9)	0.0011 (10)	0.0037 (9)
Cl1	0.0835 (6)	0.0675 (5)	0.1364 (9)	0.0378 (5)	-0.0058 (6)	0.0273 (5)
N1	0.0389 (16)	0.0346 (14)	0.0204 (15)	0.0012 (9)	0.000	0.000
N2	0.0429 (17)	0.0363 (15)	0.0233 (15)	-0.0023 (10)	0.000	0.000
O1	0.0569 (11)	0.0459 (9)	0.0766 (13)	0.0038 (8)	0.0336 (10)	-0.0001 (9)
O2	0.0546 (11)	0.0341 (7)	0.0729 (12)	0.0044 (7)	-0.0229 (9)	-0.0095 (8)
O3	0.0453 (8)	0.0272 (6)	0.0333 (8)	0.0002 (5)	-0.0010 (7)	-0.0007 (6)
O4	0.0391 (8)	0.0364 (6)	0.0380 (8)	0.0041 (6)	0.0044 (6)	0.0040 (6)
S1	0.0353 (3)	0.0278 (2)	0.0521 (3)	-0.00156 (18)	0.0019 (2)	-0.0024 (2)
Zn1	0.03723 (16)	0.02579 (13)	0.02058 (13)	0.00031 (13)	0.000	0.000

Geometric parameters (\AA , $^\circ$)

C1—C6	1.369 (3)	C10—H8	0.9300
C1—C2	1.390 (3)	C11—C10 ⁱ	1.376 (3)
C1—S1	1.799 (2)	C11—C12	1.487 (3)
C2—C3	1.380 (3)	C12—C13 ⁱ	1.392 (3)
C2—H1	0.9300	C12—C13	1.392 (3)
C3—C4	1.371 (4)	C13—C14	1.377 (3)
C3—H2	0.9300	C13—H9	0.9300
C4—C5	1.374 (4)	C14—N2	1.346 (3)
C4—Cl1	1.746 (2)	C14—H10	0.9300
C5—C6	1.383 (4)	N1—C9 ⁱ	1.328 (3)

C5—H3	0.9300	N1—Zn1	2.174 (3)
C6—H4	0.9300	N2—C14 ⁱ	1.346 (3)
C7—C8	1.533 (3)	N2—Zn1 ⁱⁱ	2.198 (3)
C7—S1	1.797 (2)	O1—S1	1.5057 (19)
C7—H5	0.9700	O3—Zn1	2.1036 (14)
C7—H6	0.9700	O4—Zn1	2.1254 (15)
C8—O2	1.240 (3)	O4—H12	0.8499
C8—O3	1.255 (2)	O4—H11	0.8500
C9—N1	1.328 (3)	Zn1—O3 ⁱ	2.1036 (14)
C9—C10	1.385 (4)	Zn1—O4 ⁱ	2.1254 (15)
C9—H7	0.9300	Zn1—N2 ⁱⁱⁱ	2.198 (3)
C10—C11	1.376 (3)		
C6—C1—C2	121.2 (2)	C13 ⁱ —C12—C11	121.74 (16)
C6—C1—S1	119.95 (18)	C13—C12—C11	121.74 (16)
C2—C1—S1	118.72 (18)	C14—C13—C12	120.1 (2)
C3—C2—C1	119.0 (2)	C14—C13—H9	119.9
C3—C2—H1	120.5	C12—C13—H9	119.9
C1—C2—H1	120.5	N2—C14—C13	123.3 (2)
C4—C3—C2	118.8 (2)	N2—C14—H10	118.4
C4—C3—H2	120.6	C13—C14—H10	118.4
C2—C3—H2	120.6	C9—N1—C9 ⁱ	117.6 (3)
C3—C4—C5	122.8 (2)	C9—N1—Zn1	121.18 (14)
C3—C4—Cl1	118.2 (2)	C9 ⁱ —N1—Zn1	121.18 (14)
C5—C4—Cl1	119.0 (2)	C14 ⁱ —N2—C14	116.6 (3)
C4—C5—C6	118.1 (2)	C14 ⁱ —N2—Zn1 ⁱⁱ	121.68 (14)
C4—C5—H3	121.0	C14—N2—Zn1 ⁱⁱ	121.68 (14)
C6—C5—H3	121.0	C8—O3—Zn1	128.19 (13)
C1—C6—C5	120.0 (2)	Zn1—O4—H12	112.7
C1—C6—H4	120.0	Zn1—O4—H11	104.1
C5—C6—H4	120.0	H12—O4—H11	111.7
C8—C7—S1	109.64 (15)	O1—S1—C7	104.88 (11)
C8—C7—H5	109.7	O1—S1—C1	105.59 (11)
S1—C7—H5	109.7	C7—S1—C1	97.06 (10)
C8—C7—H6	109.7	O3 ⁱ —Zn1—O3	176.74 (8)
S1—C7—H6	109.7	O3 ⁱ —Zn1—O4	89.52 (6)
H5—C7—H6	108.2	O3—Zn1—O4	90.50 (6)
O2—C8—O3	127.44 (18)	O3 ⁱ —Zn1—O4 ⁱ	90.50 (6)
O2—C8—C7	118.01 (18)	O3—Zn1—O4 ⁱ	89.52 (6)
O3—C8—C7	114.54 (18)	O4—Zn1—O4 ⁱ	179.62 (8)
N1—C9—C10	122.9 (2)	O3 ⁱ —Zn1—N1	88.37 (4)
N1—C9—H7	118.6	O3—Zn1—N1	88.37 (4)
C10—C9—H7	118.6	O4—Zn1—N1	90.19 (4)
C11—C10—C9	119.4 (2)	O4 ⁱ —Zn1—N1	90.19 (4)
C11—C10—H8	120.3	O3 ⁱ —Zn1—N2 ⁱⁱⁱ	91.63 (4)
C9—C10—H8	120.3	O3—Zn1—N2 ⁱⁱⁱ	91.63 (4)

supplementary materials

C10—C11—C10 ⁱ	117.7 (3)	O4—Zn1—N2 ⁱⁱⁱ	89.81 (4)
C10—C11—C12	121.16 (16)	O4 ⁱ —Zn1—N2 ⁱⁱⁱ	89.81 (4)
C10 ⁱ —C11—C12	121.16 (16)	N1—Zn1—N2 ⁱⁱⁱ	180.000 (1)
C13 ⁱ —C12—C13	116.5 (3)		


Symmetry codes: (i) $-x, -y+2, z$; (ii) $x, y, z+1$; (iii) $x, y, z-1$.

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O4—H12···O1 ^{iv}	0.85	1.88	2.706 (2)	164
O4—H12···S1 ^{iv}	0.85	2.99	3.8217 (18)	165
O4—H11···O2 ⁱ	0.85	1.85	2.657 (2)	158

Symmetry codes: (iv) $x+1/4, -y+7/4, z-1/4$; (i) $-x, -y+2, z$.

Fig. 1



supplementary materials

Fig. 2

