

Retraction of articles

This article reports the retraction of 39 articles published in *Acta Crystallographica Section E* between 2004 and 2009.

After thorough investigation (see Harrison *et al.*, 2010), 39 additional articles are retracted as a result of problems with the data sets or incorrect atom assignments. Full details of all the articles are given in Table 1.

Table 1

Details of articles to be retracted, in order of publication.

Title	Reference	Retracted by	DOI	Refcode
<i>trans</i> -Bis[1-β-(cyclohexylamino)propyliminomethyl]-2-naphtholato]copper(II) dichloride dihydrate	Zhang (2004)	Journal	10.1107/S1600536804028296	BIPDUA
Bis(4-bromo-2-formylphenolato-κ ² O,O')copper(II)	Sun & Gao (2005)	Author	10.1107/S16005368050187X	FEYSUY
Bis(salicylaldehyde)zinc(II)	Xiong & Liu (2005)	Journal	10.1107/S1600536805010913	GAMDUU
Bis(4-bromo-2-formylphenolato-κ ² O,O')zinc(II)	Chen (2006)	Journal	10.1107/S1600536805040432	SAZCUS
Bis(2-formylphenolato-κ ² O,O')nickel(II)	Li & Chen (2006)	Journal	10.1107/S1600536806012931	IDAZAP
Bis(2-formylphenolato)cobalt(II)	Qiu (2006)	Journal	10.1107/S1600536806015704	GEJDUV
Bis(2-formylphenolato-κ ² O,O')manganese(II)	Wang & Fang (2006)	Journal	10.1107/S1600536806021039	IDOVED
Tetraqua(1,10-phenanthroline-κ ² N,N')copper(II) naphthalene-1,5-disulfonate dihydrate	Liu <i>et al.</i> (2006)	Author	10.1107/S1600536806030637	GENYOO
Tetraqua(1,10-phenanthroline-κ ² N,N')nickel(II) naphthalene-1,5-disulfonate dihydrate	Liu & Fan (2006)	Author	10.1107/S1600536806035410	KERBEP
{6,6'-Diethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-trinitratolutetium(III)copper(II)	Sui <i>et al.</i> (2006)	Journal	10.1107/S160053680604565X	HESPEP
Bis(2-formylphenolato-κ ² O,O')iron(II)	Yang <i>et al.</i> (2007)	Author	10.1107/S1600536807021721	PIFCAJ
2,6-Dimethoxybenzohydrazide	Qadeer <i>et al.</i> (2007a)	Journal	10.1107/S1600536807022593	PIFHES
2-(2,4-Dichlorophenylsulfanylmethyl)acetohydrazide	Qadeer <i>et al.</i> (2007b)	Journal	10.1107/S1600536807022891	YIFSOW
{6,6'-Diethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-trinitratoeuropium(III)zinc(II)	Hu <i>et al.</i> (2007)	Author	10.1107/S1600536807031121	WIHKEE
{μ-6,6'-Diethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-trinitratocerium(III)zinc(II)	Sui, Zhang, Hu & Yin (2007)	Author	10.1107/S1600536807032564	WIHREL
{μ-6,6'-Diethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-trinitratorpaseodymium(III)zinc(II)	Chen <i>et al.</i> (2007)	Author	10.1107/S1600536807032540	WIHRIP
{μ-6,6'-Diethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-trinitratorpaseodymium(III)nickel(II)	Sui, Li <i>et al.</i> (2007)	Author	10.1107/S1600536807032618	UFACUA
{6,6'-Dimethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato-1κ ⁴ O ^I ,O ^V ,O ⁶ :2κ ⁴ O ^I ,N,N',O ^V](methanol-1κO)-μ-nitrito-1:2κ ² O:O'-dinitrato-1κ ⁴ O,O'-cerium(III)zinc(II)	Sui, Fang, Hu & Lin (2007)	Author	10.1107/S1600536807033314	UDUYIC
{6,6'-Dimethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-methanol-μ-nitrito-dinitratosamarium(III)nickel(II)	Sui, Zhang, Hu & Jiang (2007)	Author	10.1107/S1600536807037130	AFECEU
{6,6'-Dimethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-methanol-μ-nitrito-dinitratorpaseodymium(III)zinc(II)	Sui, Fang & Yuan (2007)	Author	10.1107/S1600536807037488	AFICEY
{6,6'-Dimethoxy-2,2'-ethane-1,2-diylbis(nitrilomethylidyne)}diphenolato]-methanol-μ-nitrito-dinitratolutetium(III)zinc(II)	Sui, Sui <i>et al.</i> (2007)	Author	10.1107/S1600536807037737	AFEFOH
catena-Poly[<i>μ</i> chloridonickel(II)- <i>μ</i> -chlorido- <i>μ</i> chloridonickel(II)]-μ-4,4'-methylenebis(3,5-dimethylpyrazole)- <i>n</i> ² N ² :N ²]	Huang & Chen (2007)	Author	10.1107/S1600536807039384	VIJYOD
{2,2'- <i>o</i> -Phenylenebis(nitrilomethylidyne)}diphenolatozinc(II)	Liu <i>et al.</i> (2007a)	Author	10.1107/S1600536807040640	DIKYUS
trans-Bis(ethylenediamine-2 ^{N,N'})bis(nitrato-κO)zinc(II)	Liu, Zeng & Chen (2007)	Author	10.1107/S1600536807042390	XIKYEW
[N,N'- <i>o</i> -Phenylenebis(picolinamido)-κ ² N,N',N'',N''']cobalt(II)	Liu & Zeng (2007a)	Author	10.1107/S1600536807044571	XILFII
[N,N'- <i>o</i> -Phenylenedipicolinamide-κ ⁴ N]nickel(II)	Liu & Zeng (2007b)	Author	10.1107/S1600536807048386	WINWEW
{2,2'- <i>o</i> -Phenylenebis(nitrilomethylidyne)}diphenolato)manganese(II)	Liu <i>et al.</i> (2007b)	Author	10.1107/S1600536807052993	VIQPIV
N-(2-Amino-3-pyridyl)urea monohydrate	Li <i>et al.</i> (2007)	Author	10.1107/S1600536807047526	SIMFEA
N-(2-Fluorophenyl)carbamic acid monohydrate	Yang (2007)	Author	10.1107/S1600536807052464	WINMOW
Aqua(dimethylglyoxime-κ ² N,N')(3,5-dinitro-2-oxidobenzoato-κ ² O ^I ,O ²)-copper(II)	Liu & Wen (2007)	Author	10.1107/S1600536807054244	HIQCAM
μ-Acetoato-tri-μ-ferrocenecarboxylatobis[(N,N-dimethylformamide)-copper(II)]	Liu, Lin <i>et al.</i> (2007)	Journal	10.1107/S1600536807059041	HIQQEE

addenda and errata

Table 1 (continued)

Title	Reference	Retracted by	DOI	Refcode
{ μ -6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato- μ -nitro-dinitratoeuropium(III)zinc(II)	Hu <i>et al.</i> (2008)	Author	10.1107/S160053680706151X	MIRPAF
Bis(4-chloro-2-formylphenolato)nickel(II)	Li <i>et al.</i> (2008)	Author	10.1107/S1600536807056309	RISTET
{ μ -6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato- μ -nitro-dinitratoterbium(III)zinc(II)	Chen <i>et al.</i> (2008)	Author	10.1107/S1600536808006958	QIXHIP
Bis(2-ethoxy-6-formylphenolato- κ^2O^1,O^6)nickel(II)	Han (2008)	Journal	10.1107/S160053680800809X	QIXLIT
{ μ -6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato- μ -nitro-dinitratoholmium(III)zinc(II)	Xiao, Sui <i>et al.</i> (2008)	Author	10.1107/S1600536808013743	BIZTUA
{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato-trinitratoholmium(III)nickel(II)	Xiao, Fu <i>et al.</i> (2008)	Author	10.1107/S1600536808013755	BIZVAI
Hydrogen-bonding patterns in the cocrystal terephthalic acid-4,4'-bipyridine (2/1)	Wang <i>et al.</i> (2009)	Journal	10.1107/S160053680903236X	DUCZEH
{ μ 6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato- $1k^4O^1,O^r,O^6,O^{\prime\prime}:2k^4O^1,N,N',O^{\prime\prime}\}$ (ethanol- $1kO$)- μ -nitro- $1:2k^2O:O'$ -dinitrato- $1k^2O,O'$ -samarium(III)zinc(II)	Huang <i>et al.</i> (2009)	Journal	10.1107/S1600536809033558	YUCWAV

References

- Chen, Q. (2006). *Acta Cryst.* **E62**, m56–m57.
- Chen, J.-R., Sui, Y., Luo, Q.-Y. & Jiang, R.-Q. (2007). *Acta Cryst.* **E63**, m2091–m2092.
- Chen, J.-R., Sui, Y., Wen, J.-W. & Yin, L.-Y. (2008). *Acta Cryst.* **E64**, m562–m563.
- Han, Z.-Q. (2008). *Acta Cryst.* **E64**, m592.
- Harrison, W. T. A., Simpson, J. & Weil, M. (2010). *Acta Cryst.* **E66**, e1–e2.
- Hu, R.-H., Sui, Y., Chen, L. & He, C.-M. (2008). *Acta Cryst.* **E64**, m8–m9.
- Hu, R.-H., Sui, Y., Fang, X.-N. & Chen, H.-M. (2007). *Acta Cryst.* **E63**, m2039–m2040.
- Huang, C.-F. & Chen, H.-L. (2007). *Acta Cryst.* **E63**, m2356–m2357.
- Huang, Q., Sui, Y.-H. & Zhang, G.-X. (2009). *Acta Cryst.* **E65**, m1161–m1162.
- Li, Y.-G. & Chen, H.-J. (2006). *Acta Cryst.* **E62**, m1038–m1039.
- Li, N.-G., Tao, R.-M. & Fu, B.-F. (2007). *Acta Cryst.* **E63**, o4228.
- Li, Z., Zhang, X. & Pu, X. (2008). *Acta Cryst.* **E64**, m215.
- Liu, J.-T. & Fan, S.-D. (2006). *Acta Cryst.* **E62**, m2507–m2508.
- Liu, J.-T., Fan, S.-D. & Li, D.-Q. (2006). *Acta Cryst.* **E62**, m2165–m2166.
- Liu, D., Lin, J., Xu, Y., Huang, C. & Li, X. (2007). *Acta Cryst.* **E63**, m3094.
- Liu, Y.-Q. & Wen, H.-R. (2007). *Acta Cryst.* **E63**, m2928.
- Liu, Y.-Q. & Zeng, X.-R. (2007a). *Acta Cryst.* **E63**, m2547.
- Liu, Y.-Q. & Zeng, X.-R. (2007b). *Acta Cryst.* **E63**, m2684.
- Liu, Y.-Q., Zeng, X.-R. & Chen, W.-T. (2007). *Acta Cryst.* **E63**, m2462.
- Liu, Y.-Q., Zeng, X.-R., Luo, Q.-Y. & Xu, Y.-P. (2007a). *Acta Cryst.* **E63**, m2396.
- Liu, Y.-Q., Zeng, X.-R., Luo, Q.-Y. & Xu, Y.-P. (2007b). *Acta Cryst.* **E63**, m2854.
- Qadeer, G., Rama, N. H. & Chen, W.-T. (2007a). *Acta Cryst.* **E63**, o2892.
- Qadeer, G., Rama, N. H. & Chen, W.-T. (2007b). *Acta Cryst.* **E63**, o2932.
- Qiu, X.-Y. (2006). *Acta Cryst.* **E62**, m1190–m1191.
- Sui, Y., Fang, X.-N., Hu, P. & Lin, J. (2007). *Acta Cryst.* **E63**, m2135–m2136.
- Sui, Y., Fang, X.-N. & Yuan, M.-W. (2007). *Acta Cryst.* **E63**, m2275–m2276.
- Sui, Y., Li, X.-F., Huang, G.-S. & Wang, G.-J. (2007). *Acta Cryst.* **E63**, m2093–m2094.
- Sui, Y., Sui, Y.-H., Luo, Q.-Y. & Wang, Y.-D. (2007). *Acta Cryst.* **E63**, m2277–m2278.
- Sui, Y., Xiao, Y.-A., Fang, X.-N., Zeng, X.-R. & Li, M.-H. (2006). *Acta Cryst.* **E62**, m3205–m3207.
- Sui, Y., Zhang, J.-H., Hu, R.-H. & Jiang, R.-Q. (2007). *Acta Cryst.* **E63**, m2256–m2257.
- Sui, Y., Zhang, J.-H., Hu, R.-H. & Yin, L.-Y. (2007). *Acta Cryst.* **E63**, m2089–m2090.
- Sun, Y.-X. & Gao, G.-Z. (2005). *Acta Cryst.* **E61**, m354–m355.
- Wang, Q. & Fang, Z.-N. (2006). *Acta Cryst.* **E62**, m1492–m1493.
- Wang, S., Yang, T., Li, Z. & Yu, X. (2009). *Acta Cryst.* **E65**, o2198.
- Xiao, Y.-A., Fu, X.-K., Sui, Y., Wu, Q. & Xiong, S.-H. (2008). *Acta Cryst.* **E64**, m806–m807.
- Xiao, Y.-A., Sui, Y., Yi, X.-G., Wu, J.-H. & Zhang, L.-P. (2008). *Acta Cryst.* **E64**, m804–m805.
- Xiong, Z.-Y. & Liu, L.-J. (2005). *Acta Cryst.* **E61**, m863–m864.
- Yang, X.-M. (2007). *Acta Cryst.* **E63**, o4453.
- Yang, Y.-M., Lu, P.-C., Zhu, T.-T. & Liu, C.-H. (2007). *Acta Cryst.* **E63**, m1613.
- Zhang, P. (2004). *Acta Cryst.* **E60**, m1808–m1810.

{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}trinitratorpaseodymium(III)-zinc(II)

Jing-Rong Chen,^a Yan Sui,^{b*} Qiu-Yan Luo^b and Run-Qing Jiang^b

^aCollege of Chemistry and Chemical Engineering, Southwest University, 400715 Beibei, Chongqing, People's Republic of China, and ^bJiangXi Province Key Laboratory of Coordination Chemistry, College of Chemistry and Chemical Engineering, JingGangShan University, 343009 Ji'an, JiangXi, People's Republic of China

Correspondence e-mail: ysui@163.com


Received 1 July 2007; accepted 4 July 2007

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.027; wR factor = 0.061; data-to-parameter ratio = 16.7.

In the title heteronuclear Zn^{II} – Pr^{III} complex (systematic name: {6,6'-diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato- κ^4O^1,O^1,O^6,O^6 : κ^2O^1,N,N',O^1 }trinitrato- κ^6O,O' -praseodymium(III)zinc(II)), $[PrZn(C_{20}H_{22}N_2O_4)(NO_3)_3]$, with the hexadentate Schiff base compartmental ligand N,N' -bis(3-ethoxysalicylidene)ethylenediamine (H_2L), the Zn^{II} and Pr^{III} atoms are doubly bridged by two phenolate O atoms provided by the Schiff base ligand. The coordination of the Zn^{II} atom is approximately square planar, involving the two imine N atoms and the two phenolate O atoms. The Pr^{III} centre has a decacoordination environment of O atoms, formed by the phenolate ligands, the two ethoxy O atoms and two O atoms from each of the three nitrates. Some weak C–H···O and O···Zn [3.159 (4) Å] interactions generate a two-dimensional zigzag sheet.

Related literature

For related literature, see: Baggio *et al.* (2000); Caravan *et al.* (1999); Edder *et al.* (2000); Knoer *et al.* (2005); Sui *et al.* (2006).



Experimental

Crystal data

$[PrZn(C_{20}H_{22}N_2O_4)(NO_3)_3]$

$M_r = 746.71$

Orthorhombic, $P2_12_12_1$

$a = 8.6317$ (7) Å

$b = 13.8782$ (11) Å

$c = 21.1267$ (16) Å

$V = 2530.8$ (3) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 2.93$ mm⁻¹

$T = 293$ (2) K

$0.21 \times 0.15 \times 0.13$ mm

Data collection

Bruker APEXII area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2004)

$T_{min} = 0.605$, $T_{max} = 0.683$

19198 measured reflections

6058 independent reflections

4803 reflections with $I > 2\sigma(I)$

$R_{int} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$

$wR(F^2) = 0.061$

$S = 1.01$

6058 reflections

363 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.50$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.62$ e Å⁻³

Absolute structure: Flack (1983), with 2527 Friedel pairs

Flack parameter: 0.021 (12)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C20–H20A···O9	0.96	2.44	3.148 (5)	131
C17–H17A···O8 ⁱ	0.97	2.59	3.528 (5)	163
C9–H9A···O7 ⁱⁱ	0.97	2.40	3.266 (5)	148
C7–H7···O7 ⁱⁱⁱ	0.93	2.36	3.276 (5)	168

Symmetry codes: (i) $x - \frac{1}{2}, -y + \frac{5}{2}, -z$; (ii) $x - 1, y, z$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2*; data reduction: *APEX2*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *APEX2*; software used to prepare material for publication: *APEX2* and *publCIF* (Westrip, 2007).

metal-organic compounds

The authors gratefully acknowledge financial support from the Department of Education, JiangXi Province (grant Nos. 2007317 and 05YB195).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2254).

References

- Baggio, R., Garland, M. T., Moreno, Y., Pena, O., Perec, M. & Spodine, E. (2000). *J. Chem. Soc. Dalton Trans.* pp. 2061–2066.
- Bruker (2004). *APEX2* (Version 1.22) and *SADABS* (Version 1.22). Bruker AXS Inc., Madison, Wisconsin, USA.
- Caravan, P., Ellison, J. J., McMurry, T. J. & Lauffer, R. B. (1999). *Chem. Rev.* **99**, 2293–2352.
- Edder, C., Piguet, C., Bernardinelli, G., Mareda, J., Bochet, C. G., Bunzli, J.-C. G. & Hopfgartner, G. (2000). *Inorg. Chem.* **39**, 5059–5073.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Knoer, R., Lin, H.-H., Wei, H.-H. & Mohanta, S. (2005). *Inorg. Chem.* **44**, 3524–3536.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sui, Y., Fang, X.-N., Xiao, Y.-A., Luo, Q.-Y. & Li, M.-H. (2006). *Acta Cryst. E* **62**, m2230–m2232.
- Westrip, S. P. (2007). *publCIF*. In preparation.

Article retracted

supplementary materials

Article retracted

Acta Cryst. (2007). E63, m2091-m2092 [doi:10.1107/S1600536807032540]

{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}trinitratopraseodymium(III)zinc(II)

J.-R. Chen, Y. Sui, Q.-Y. Luo and R.-Q. Jiang

Comment

The potential applications of trivalent lanthanide complexes as contrast agent for magnetic resonance imaging and stains for fluorescence imaging have prompted considerable interest in the preparation, magnetic and optical properties of 3 d–4f heterometallic dinuclear complexes (Baggio *et al.*, 2000; Caravan *et al.*, 1999; Edder *et al.*, 2000; Knoer *et al.*, 2005). As part of our investigations into the structure and applications of 3 d–4f heterometallic Schiff base complexes (Sui *et al.*, 2006), we report here the synthesis and X-ray crystal structure analysis of the title complex, (I), a new Zn^{II}–Pr^{III} complex with salen-type Schiff base *N,N'*-bis(3-ethoxysalicylidene) ethylenediamine(H₂L).

Complex (I) crystallizes in the space group *P*2₁2₁2₁, with zinc and praseodymium doubly bridged by two phenolate O atoms provided by a salen-type Schiff base ligand. The inner salen-type cavity is occupied by zinc(II), while praseodymium(III) is present in the open and larger portion of the dinucleating compartmental Schiff base ligand. The dihedral angles between the mean planes of Zn1/O1/O2 and Pr1/O1/O2 is 3.45 (15)^o suggesting that the bridging moiety is almost planar; the deviation of atoms from the least squares Zn1/O1/O2/Pr1 plane being 0.0293 (3) Å for Zn, 0.0199 (3) Å for Pr, –0.0244 (4) Å for O1 and –0.0248 (4) Å for O2.

The praseodymium(III) center in (I) has a decacoordination environment of O atoms. In addition to the phenolate ligands, two ethoxy O atoms coordinate to this metal center, two O atoms from each of the three nitrates chelate to praseodymium to complete the decacoordination. The three kinds of Pr—O bond distances are significantly different, the shortest being the Pr—O(phenolate) and longest being the Pr—O(ethoxy) separations.

The coordination of zinc(II) is approximately square planar. The donor centers are alternatively above and below the mean N₂O₂ plane with an average deviation from the plane of 0.0901 (3) Å, while Zn1 is 0.0437 (3) Å below this square plane.

Adjacent molecules are held together by weak interactions (O10···Zn1 = 3.159 (5) Å, C7—H7···O7ⁱ = 3.276 (5), C9—H9A···O7ⁱⁱ = 3.266 (5) and C17—H17A···O8ⁱⁱⁱ = 3.528 (5); symmetry codes: (i) 1 – *x*, *y* – 1/2, 1/2 – *z*; (ii) *x* – 1, *y*, *z*; (iii) *x* – 1/2, 5/2 – *y*, *z*) these link the molecules into a two-dimensional zigzag sheet (Fig. 2).

Experimental

H₂L was prepared by the 2:1 condensation of 3-ethoxysalicylaldehyde and ethylenediamine in methanol. Complex (I) was obtained by the treatment of zinc(II) acetate dihydrate (0.188 g, 1 mmol) with H₂L(0.356 g, 1 mmol) in methanol solution (100 ml) under reflux for 3 h and then for another 3 h after the addition of praseodymium(III) nitrate hexahydrate (0.435 g, 1 mmol). The reaction mixture was cooled and the resulting precipitate was filtered off, washed with diethyl ether and dried *in vacuo*. Single crystals of (I) suitable for X-ray analysis were obtained by slow evaporation at room temperature of

supplementary materials

a methanol solution. Analysis calculated for $C_{20}H_{22}N_5O_{13}PrZn$: C 32.17, H 2.97, N 9.38, Pr 18.87, Zn 8.76%; found: C 32.28, H 2.95, N 9.31, Pr 19.05, Zn 8.78%. IR (KBr, cm^{-1}): 1643 (C=N), 1385, 1491 (nitrate).

Refinement

The H atoms were positioned geometrically and treated as riding on their parent atoms, with C—H distances of 0.97 (methylene) and 0.96 Å (methyl), and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and $1.2U_{\text{eq}}(\text{C})$ for other H atoms.

Figures




Fig. 1. The molecular structure of (I), showing 30% probability displacement ellipsoids. All the H atoms on carbon have been omitted for clarity.





Fig. 2. The packing diagram of (I), viewed along the b axis; hydrogen bonds are shown as dashed lines.

Table 1. Selected geometric parameters (Å).



Crystal data



$F_{000} = 1480$

$M_r = 746.71$

$D_x = 1.960 \text{ Mg m}^{-3}$

Orthorhombic, $P2_12_12_1$

Mo $K\alpha$ radiation

Hall symbol: P 2ac 2ab

$\lambda = 0.71073 \text{ \AA}$

$a = 8.6317 (7) \text{ \AA}$

Cell parameters from 19198 reflections

$b = 13.8782 (11) \text{ \AA}$

$\theta = 1.9\text{--}28.2^\circ$

$c = 21.1267 (16) \text{ \AA}$

$\mu = 2.93 \text{ mm}^{-1}$

$V = 2530.8 (3) \text{ \AA}^3$

$T = 293 (2) \text{ K}$

$Z = 4$

Block, yellow

$0.21 \times 0.15 \times 0.13 \text{ mm}$

Data collection

Bruker APEXII area-detector

6058 independent reflections

diffractometer

Radiation source: fine-focus sealed tube

4803 reflections with $I > 2\sigma(I)$

Monochromator: graphite

 $R_{\text{int}} = 0.031$ Detector resolution: 0 pixels mm⁻¹ $\theta_{\text{max}} = 28.2^\circ$ $T = 293(2)$ K $\theta_{\text{min}} = 1.9^\circ$ φ and ω scans $h = -11 \rightarrow 11$ Absorption correction: multi-scan
(SADABS; Bruker, 2004) $k = -18 \rightarrow 17$ $T_{\text{min}} = 0.605$, $T_{\text{max}} = 0.683$ $l = -27 \rightarrow 28$

19198 measured reflections

*Refinement*Refinement on F^2

Hydrogen site location: inferred from neighbouring sites

Least-squares matrix: full

H-atom parameters constrained

 $R[F^2 > 2\sigma(F^2)] = 0.027$ $w = 1/[\sigma^2(F_o^2) + (0.0273P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $wR(F^2) = 0.061$ $(\Delta/\sigma)_{\text{max}} = 0.002$ $S = 1.01$ $\Delta\rho_{\text{max}} = 0.50 \text{ e } \text{\AA}^{-3}$

6058 reflections

 $\Delta\rho_{\text{min}} = -0.62 \text{ e } \text{\AA}^{-3}$

363 parameters

Extinction correction: none

Primary atom site location: structure-invariant direct methods

Absolute structure: Flack (1983), with how many Friedel pairs?

Secondary atom site location: difference Fourier map

Flack parameter: 0.021 (12)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O7	0.9052 (4)	1.0762 (3)	0.27853 (15)	0.0806 (12)
O10	1.2389 (4)	1.0397 (3)	0.07303 (19)	0.0945 (13)
O13	0.5733 (4)	1.0012 (3)	-0.08812 (14)	0.0756 (9)
O1	0.5859 (3)	0.87492 (16)	0.13666 (12)	0.0373 (6)
Pr1	0.758724 (19)	0.999692 (14)	0.094892 (8)	0.03144 (5)
Zn1	0.43316 (5)	0.94378 (3)	0.18296 (2)	0.03984 (11)
O2	0.5377 (3)	1.05631 (17)	0.15394 (11)	0.0344 (6)

supplementary materials

C16	0.4779 (4)	1.1447 (2)	0.15604 (17)	0.0324 (8)
N2	0.3000 (3)	1.0138 (2)	0.23919 (15)	0.0399 (8)
C2	0.7087 (4)	0.7432 (2)	0.08988 (17)	0.0328 (8)
N1	0.3204 (4)	0.8318 (2)	0.20905 (16)	0.0372 (8)
O3	0.8110 (3)	0.81322 (18)	0.06945 (13)	0.0377 (6)
C1	0.5866 (4)	0.7798 (2)	0.12626 (18)	0.0326 (8)
C15	0.5382 (4)	1.2125 (2)	0.11381 (18)	0.0333 (9)
O4	0.6539 (3)	1.17627 (17)	0.07518 (12)	0.0365 (6)
C10	0.2792 (4)	1.1050 (3)	0.23906 (18)	0.0389 (9)
H10	0.2104	1.1305	0.2686	0.047*
C11	0.3548 (5)	1.1715 (3)	0.19636 (19)	0.0381 (10)
C6	0.4752 (4)	0.7162 (3)	0.15047 (18)	0.0362 (9)
C8	0.1862 (5)	0.8551 (3)	0.2493 (2)	0.0450 (10)
H8A	0.1678	0.8036	0.2794	0.054*
H8B	0.0941	0.8630	0.2235	0.054*
C7	0.3493 (4)	0.7452 (3)	0.1915 (2)	0.0402 (10)
H7	0.2837	0.6971	0.2063	0.048*
C9	0.2226 (4)	0.9483 (3)	0.28391 (18)	0.0432 (9)
H9A	0.1278	0.9774	0.2995	0.052*
H9B	0.2897	0.9355	0.3198	0.052*
O11	0.7691 (3)	1.0125 (2)	-0.02391 (13)	0.0544 (7)
N5	0.6277 (4)	0.9941 (3)	-0.03462 (16)	0.0501 (8)
O6	0.8208 (4)	0.9677 (2)	0.21236 (15)	0.0540 (9)
O5	0.8639 (3)	1.1122 (2)	0.18006 (15)	0.0514 (7)
O8	1.0052 (4)	1.0817 (3)	0.04956 (16)	0.0674 (10)
O9	1.0450 (3)	0.9619 (2)	0.11152 (15)	0.0542 (8)
O12	0.5486 (3)	0.9684 (2)	0.01210 (14)	0.0600 (9)
N3	0.8645 (4)	1.0526 (3)	0.22519 (19)	0.0494 (9)
N4	1.1002 (4)	1.0277 (3)	0.07785 (18)	0.0536 (11)
C3	0.7170 (5)	0.6468 (3)	0.07493 (19)	0.0439 (10)
H3	0.7985	0.6234	0.0506	0.053*
C17	0.7260 (5)	1.2413 (3)	0.02942 (18)	0.0425 (9)
H17A	0.6464	1.2807	0.0099	0.051*
H17B	0.7748	1.2035	-0.0036	0.051*
C19	0.9463 (4)	0.7813 (3)	0.03462 (19)	0.0441 (10)
H19A	0.9888	0.8351	0.0110	0.053*
H19B	0.9158	0.7321	0.0045	0.053*
C4	0.6018 (5)	0.5849 (3)	0.0968 (2)	0.0499 (11)
H4	0.6048	0.5200	0.0859	0.060*
C5	0.4870 (5)	0.6179 (3)	0.1333 (2)	0.0448 (10)
H5	0.4125	0.5749	0.1478	0.054*
C20	1.0685 (5)	0.7415 (3)	0.0778 (2)	0.0585 (12)
H20A	1.0960	0.7892	0.1087	0.088*
H20B	1.1584	0.7244	0.0535	0.088*
H20C	1.0292	0.6852	0.0989	0.088*
C14	0.4813 (4)	1.3053 (3)	0.1114 (2)	0.0417 (10)
H14	0.5216	1.3497	0.0829	0.050*
C12	0.3023 (5)	1.2676 (3)	0.1951 (2)	0.0468 (11)
H12	0.2262	1.2876	0.2233	0.056*

C13	0.3623 (5)	1.3311 (3)	0.1527 (2)	0.0507 (11)
H13	0.3229	1.3935	0.1512	0.061*
C18	0.8445 (5)	1.3056 (3)	0.0590 (2)	0.0572 (13)
H18A	0.7941	1.3503	0.0870	0.086*
H18B	0.8984	1.3405	0.0265	0.086*
H18C	0.9170	1.2673	0.0826	0.086*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O7	0.086 (3)	0.112 (3)	0.0432 (18)	0.047 (2)	-0.0212 (19)	-0.025 (2)
O10	0.0346 (17)	0.142 (4)	0.107 (3)	-0.021 (2)	0.0064 (19)	0.004 (3)
O13	0.103 (2)	0.074 (2)	0.0496 (18)	-0.005 (2)	-0.0228 (18)	0.000 (2)
O1	0.0430 (15)	0.0200 (12)	0.0490 (16)	-0.0035 (11)	0.0155 (13)	0.0011 (11)
Pr1	0.03149 (9)	0.02595 (9)	0.03689 (9)	-0.00118 (11)	0.00386 (8)	0.00233 (10)
Zn1	0.0393 (2)	0.0319 (2)	0.0484 (2)	-0.00127 (19)	0.0103 (2)	0.0038 (2)
O2	0.0319 (13)	0.0252 (12)	0.0461 (14)	0.0024 (11)	0.0110 (12)	0.0032 (12)
C16	0.035 (2)	0.0256 (18)	0.037 (2)	0.0001 (14)	-0.0055 (17)	0.0018 (16)
N2	0.0380 (15)	0.046 (2)	0.0360 (17)	-0.0022 (14)	0.0057 (13)	0.0045 (16)
C2	0.0385 (19)	0.0280 (18)	0.0318 (19)	-0.0010 (13)	-0.0040 (17)	0.0032 (16)
N1	0.0367 (18)	0.0365 (19)	0.0385 (19)	-0.0051 (14)	0.0046 (15)	0.0092 (15)
O3	0.0353 (13)	0.0325 (14)	0.0455 (16)	0.0012 (11)	0.0085 (12)	-0.0013 (12)
C1	0.035 (2)	0.0272 (19)	0.035 (2)	-0.0006 (15)	-0.0058 (18)	0.0001 (16)
C15	0.032 (2)	0.0285 (19)	0.040 (2)	-0.0009 (14)	-0.0042 (17)	-0.0023 (16)
O4	0.0435 (15)	0.0250 (13)	0.0411 (16)	-0.0022 (11)	0.0063 (12)	0.0077 (11)
C10	0.037 (2)	0.040 (2)	0.040 (2)	0.0069 (17)	0.0043 (18)	0.0001 (18)
C11	0.036 (2)	0.037 (2)	0.042 (3)	0.0026 (16)	0.0013 (19)	0.0026 (18)
C6	0.044 (2)	0.0284 (19)	0.036 (2)	-0.0056 (15)	-0.0051 (18)	0.0051 (17)
C8	0.036 (2)	0.054 (3)	0.045 (2)	-0.0030 (18)	0.0102 (19)	0.012 (2)
C7	0.038 (2)	0.035 (2)	0.047 (3)	-0.0098 (16)	-0.003 (2)	0.0125 (19)
C9	0.042 (2)	0.044 (2)	0.043 (2)	0.0011 (18)	0.0110 (19)	0.0094 (19)
O11	0.0550 (17)	0.062 (2)	0.0461 (15)	-0.0113 (17)	0.0047 (13)	0.0002 (14)
N5	0.065 (2)	0.0392 (18)	0.046 (2)	0.001 (2)	-0.0037 (18)	-0.011 (2)
O6	0.059 (2)	0.056 (2)	0.0468 (18)	0.0084 (15)	0.0059 (15)	0.0152 (15)
O5	0.0562 (18)	0.0475 (17)	0.0505 (18)	0.0016 (14)	-0.0056 (16)	-0.0008 (16)
O8	0.0447 (18)	0.086 (3)	0.072 (2)	-0.0162 (16)	-0.0007 (17)	0.033 (2)
O9	0.0414 (16)	0.0446 (16)	0.077 (2)	0.0007 (13)	0.0054 (16)	0.0056 (15)
O12	0.0477 (16)	0.084 (3)	0.0482 (17)	-0.0121 (15)	0.0016 (15)	-0.0030 (16)
N3	0.0357 (19)	0.064 (3)	0.049 (2)	0.0208 (19)	-0.0028 (17)	-0.010 (2)
N4	0.038 (2)	0.069 (3)	0.054 (2)	-0.0086 (17)	0.0078 (18)	-0.0151 (19)
C3	0.058 (3)	0.031 (2)	0.044 (2)	0.0052 (18)	0.001 (2)	-0.0029 (17)
C17	0.054 (2)	0.0320 (19)	0.041 (2)	-0.0025 (17)	0.009 (2)	0.0081 (16)
C19	0.039 (2)	0.041 (2)	0.052 (3)	0.0045 (17)	0.008 (2)	-0.004 (2)
C4	0.065 (3)	0.028 (2)	0.057 (3)	-0.0013 (17)	-0.006 (2)	0.001 (2)
C5	0.057 (3)	0.024 (2)	0.053 (2)	-0.0102 (17)	-0.003 (2)	0.0066 (18)
C20	0.050 (3)	0.050 (3)	0.075 (3)	0.012 (2)	-0.004 (3)	-0.008 (2)
C14	0.043 (2)	0.035 (2)	0.048 (2)	-0.0004 (16)	-0.004 (2)	0.0052 (19)
C12	0.046 (2)	0.037 (2)	0.058 (3)	0.0104 (17)	0.007 (2)	-0.006 (2)

supplementary materials

C13	0.059 (3)	0.027 (2)	0.066 (3)	0.0111 (18)	-0.002 (2)	0.001 (2)
C18	0.063 (3)	0.040 (3)	0.069 (3)	-0.011 (2)	0.020 (2)	-0.003 (2)

Geometric parameters (\AA , $^{\circ}$)

O7—N3	1.225 (5)	C6—C5	1.417 (5)
O10—N4	1.213 (4)	C6—C7	1.447 (6)
O13—N5	1.228 (4)	C8—C9	1.519 (6)
O1—C1	1.339 (4)	C8—H8A	0.9700
O1—Zn1	1.900 (2)	C8—H8B	0.9700
O1—Pr1	2.450 (2)	C7—H7	0.9300
Pr1—O2	2.411 (2)	C9—H9A	0.9700
Pr1—O11	2.518 (3)	C9—H9B	0.9700
Pr1—O5	2.549 (3)	O11—N5	1.267 (4)
Pr1—O9	2.550 (3)	N5—O12	1.252 (4)
Pr1—O12	2.557 (3)	O6—N3	1.266 (5)
Pr1—O6	2.578 (3)	O5—N3	1.262 (5)
Pr1—O8	2.596 (3)	O8—N4	1.262 (4)
Pr1—O4	2.645 (2)	O9—N4	1.252 (4)
Pr1—O3	2.681 (2)	C3—C4	1.394 (6)
Zn1—O2	1.905 (2)	C3—H3	0.9300
Zn1—N1	1.915 (3)	C17—C18	1.494 (6)
Zn1—N2	1.917 (3)	C17—H17A	0.9700
O2—C16	1.332 (4)	C17—H17B	0.9700
C16—C15	1.397 (5)	C19—C20	1.500 (6)
C16—C11	1.411 (5)	C19—H19A	0.9700
N2—C10	1.279 (5)	C19—H19B	0.9700
N2—C9	1.471 (5)	C4—C5	1.335 (5)
C2—C3	1.376 (5)	C4—H4	0.9300
C2—O3	1.383 (4)	C5—H5	0.9300
C2—C1	1.400 (5)	C20—H20A	0.9600
N1—C7	1.282 (5)	C20—H20B	0.9600
N1—C8	1.473 (5)	C20—H20C	0.9600
O3—C19	1.449 (4)	C14—C13	1.394 (6)
C1—C6	1.401 (5)	C14—H14	0.9300
C15—C14	1.379 (5)	C12—C13	1.360 (6)
C15—O4	1.384 (4)	C12—H12	0.9300
O4—C17	1.462 (4)	C13—H13	0.9300
C10—C11	1.446 (5)	C18—H18A	0.9600
C10—H10	0.9300	C18—H18B	0.9600
C11—C12	1.409 (5)	C18—H18C	0.9600
C1—O1—Zn1	125.7 (2)	N2—C10—C11	124.7 (4)
C1—O1—Pr1	129.5 (2)	N2—C10—H10	117.6
Zn1—O1—Pr1	104.62 (9)	C11—C10—H10	117.6
O2—Pr1—O1	64.04 (8)	C12—C11—C16	118.7 (4)
O2—Pr1—O11	121.38 (9)	C12—C11—C10	118.1 (4)
O1—Pr1—O11	115.51 (9)	C16—C11—C10	123.3 (3)
O2—Pr1—O5	73.57 (9)	C1—C6—C5	117.6 (4)
O1—Pr1—O5	113.29 (9)	C1—C6—C7	124.0 (3)

O11—Pr1—O5	130.36 (10)	C5—C6—C7	118.4 (3)
O2—Pr1—O9	139.68 (9)	N1—C8—C9	107.6 (3)
O1—Pr1—O9	113.26 (9)	N1—C8—H8A	110.2
O11—Pr1—O9	96.75 (10)	C9—C8—H8A	110.2
O5—Pr1—O9	71.56 (9)	N1—C8—H8B	110.2
O2—Pr1—O12	81.25 (9)	C9—C8—H8B	110.2
O1—Pr1—O12	72.21 (10)	H8A—C8—H8B	108.5
O11—Pr1—O12	49.85 (9)	N1—C7—C6	125.5 (4)
O5—Pr1—O12	147.08 (10)	N1—C7—H7	117.3
O9—Pr1—O12	138.30 (10)	C6—C7—H7	117.3
O2—Pr1—O6	73.90 (9)	N2—C9—C8	108.1 (3)
O1—Pr1—O6	70.01 (10)	N2—C9—H9A	110.1
O11—Pr1—O6	164.72 (10)	C8—C9—H9A	110.1
O5—Pr1—O6	49.60 (10)	N2—C9—H9B	110.1
O9—Pr1—O6	68.31 (10)	C8—C9—H9B	110.1
O12—Pr1—O6	140.99 (10)	H9A—C9—H9B	108.4
O2—Pr1—O8	134.14 (11)	N5—O11—Pr1	97.5 (2)
O1—Pr1—O8	160.35 (10)	O13—N5—O12	122.7 (4)
O11—Pr1—O8	64.67 (10)	O13—N5—O11	121.1 (4)
O5—Pr1—O8	72.54 (11)	O12—N5—O11	116.2 (3)
O9—Pr1—O8	49.23 (10)	N3—O6—Pr1	96.2 (2)
O12—Pr1—O8	113.79 (10)	N3—O5—Pr1	97.7 (2)
O6—Pr1—O8	105.08 (11)	N4—O8—Pr1	95.6 (2)
O2—Pr1—O4	60.58 (8)	N4—O9—Pr1	98.1 (2)
O1—Pr1—O4	120.21 (8)	N5—O12—Pr1	96.0 (2)
O11—Pr1—O4	77.86 (9)	O7—N3—O5	121.4 (4)
O5—Pr1—O4	70.47 (9)	O7—N3—O6	122.1 (5)
O9—Pr1—O4	122.93 (8)	O5—N3—O6	116.5 (4)
O12—Pr1—O4	78.87 (9)	O10—N4—O9	121.6 (4)
O6—Pr1—O4	112.47 (9)	O10—N4—O8	121.3 (4)
O8—Pr1—O4	79.40 (10)	O9—N4—O8	117.1 (3)
O2—Pr1—O3	123.47 (8)	C2—C3—C4	119.1 (4)
O1—Pr1—O3	59.51 (7)	C2—C3—H3	120.5
O11—Pr1—O3	82.10 (9)	C4—C3—H3	120.5
O5—Pr1—O3	132.26 (9)	O4—C17—C18	112.5 (3)
O9—Pr1—O3	70.49 (8)	O4—C17—H17A	109.1
O12—Pr1—O3	79.54 (9)	C18—C17—H17A	109.1
O6—Pr1—O3	89.53 (9)	O4—C17—H17B	109.1
O8—Pr1—O3	102.19 (10)	C18—C17—H17B	109.1
O4—Pr1—O3	156.94 (9)	H17A—C17—H17B	107.8
O1—Zn1—O2	85.30 (10)	O3—C19—C20	111.7 (3)
O1—Zn1—N1	95.31 (12)	O3—C19—H19A	109.3
O2—Zn1—N1	177.18 (13)	C20—C19—H19A	109.3
O1—Zn1—N2	171.90 (13)	O3—C19—H19B	109.3
O2—Zn1—N2	93.89 (12)	C20—C19—H19B	109.3
N1—Zn1—N2	85.89 (15)	H19A—C19—H19B	107.9
C16—O2—Zn1	124.1 (2)	C5—C4—C3	120.6 (4)
C16—O2—Pr1	128.6 (2)	C5—C4—H4	119.7
Zn1—O2—Pr1	105.91 (10)	C3—C4—H4	119.7

Article retracted

supplementary materials


O2—C16—C15	117.0 (3)	C4—C5—C6	122.1 (4)
O2—C16—C11	123.6 (3)	C4—C5—H5	118.9
C15—C16—C11	119.3 (3)	C6—C5—H5	118.9
C10—N2—C9	123.3 (3)	C19—C20—H20A	109.5
C10—N2—Zn1	125.8 (3)	C19—C20—H20B	109.5
C9—N2—Zn1	110.9 (3)	H20A—C20—H20B	109.5
C3—C2—O3	125.3 (3)	C19—C20—H20C	109.5
C3—C2—C1	121.2 (3)	H20A—C20—H20C	109.5
O3—C2—C1	113.4 (3)	H20B—C20—H20C	109.5
C7—N1—C8	121.7 (3)	C15—C14—C13	118.7 (4)
C7—N1—Zn1	125.4 (3)	C15—C14—H14	120.7
C8—N1—Zn1	112.8 (3)	C13—C14—H14	120.7
C2—O3—C19	117.3 (3)	C13—C12—C11	120.3 (4)
C2—O3—Pr1	120.5 (2)	C13—C12—H12	119.9
C19—O3—Pr1	122.2 (2)	C11—C12—H12	119.9
O1—C1—C2	116.8 (3)	C12—C13—C14	121.7 (4)
O1—C1—C6	123.9 (3)	C12—C13—H13	119.2
C2—C1—C6	119.3 (3)	C14—C13—H13	119.2
C14—C15—O4	125.1 (3)	C17—C18—H18A	109.5
C14—C15—C16	121.3 (3)	C17—C18—H18B	109.5
O4—C15—C16	113.7 (3)	H18A—C18—H18B	109.5
C15—O4—C17	118.2 (3)	C17—C18—H18C	109.5
C15—O4—Pr1	119.4 (2)	H18A—C18—H18C	109.5
C17—O4—Pr1	122.0 (2)	H18B—C18—H18C	109.5

Hydrogen-bond geometry (\AA , °)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C20—H20A···O9	0.96	2.44	3.148 (5)	131
C17—H17A···O8 ⁱ	0.97	2.59	3.528 (5)	163
C9—H9A···O7 ⁱⁱ	0.97	2.40	3.266 (5)	148
C7—H7···O7 ⁱⁱⁱ	0.93	2.36	3.276 (5)	168

Symmetry codes: (i) $x-1/2, -y+5/2, -z$; (ii) $x-1, y, z$; (iii) $-x+1, y-1/2, -z+1/2$.

Fig. 1



Article

supplementary materials

Fig. 2

