

## Retraction of articles by H. Zhong *et al.*

H. Zhong,<sup>a\*</sup> S.-H. Duan,<sup>a</sup> Y.-P. Hong,<sup>a</sup> M.-L. Li,<sup>a</sup> Y.-Q. Liu,<sup>a</sup> C.-J. Luo,<sup>a</sup> Q.-Y. Luo,<sup>a</sup> S.-Z. Xiao,<sup>a</sup> H.-L. Xie,<sup>a</sup> Y.-P. Xu,<sup>a</sup> X.-M. Yang,<sup>b,a</sup> X.-R. Zeng<sup>a</sup> and Q. Y. Zhong<sup>c</sup>

<sup>a</sup>College of Chemistry and Chemical Engineering, Provincial Key Laboratory of Coordination Chemistry, Jinggangshan University, Jian 343009, People's Republic of China, <sup>b</sup>Institute of Applied Materials, Jiangxi University of Finance and Economics, Nanchang 330032, People's Republic of China, and <sup>c</sup>Jian Training School, Jian 343000, People's Republic of China  
Correspondence e-mail: huazhong06@126.com

Received 20 November 2009; accepted 15 December 2009

A series of 41 papers by H. Zhong *et al.* are retracted.

As a result of problems with the data sets and incorrect atom assignments, 41 papers by H. Zhong *et al.* are retracted. Full details of all the articles are given in Table 1.

**Table 1**

Details of articles to be retracted, in order of publication.

Title	Reference	DOI	Refcode
<i>Aquachlorobis(1,10-phenanthroline)cobalt(II) chloride thiourea solvate</i>	Zhong, Zeng, Liu & Luo (2006a)	10.1107/S1600536806041122	KERQEE
<i>cis-Dichlorobis(1,10-phenanthroline)cobalt(II)</i>	Zhong, Zeng & Luo (2006)	10.1107/S1600536806047295	MEQFOE
<i>Tris(quinolin-8-olato-κ<sup>2</sup>N,O)cobalt(III) glyoxal hemisolvate monohydrate</i>	Zhong, Zeng, Liu & Luo (2006b)	10.1107/S1600536806050240	MEQHEW
<i>(8-Quinolinol-κ<sup>2</sup>N,O)bis(8-quinolinolato-κ<sup>2</sup>N,O)nickel(II) glyoxal hemisolvate monohydrate</i>	Zhong, Zeng, Liu & Luo (2007)	10.1107/S1600536806053232	METVUD
<i>Aquachlorobis(1,10-phenanthroline)cobalt(II) chloride thioacetamide solvate</i>	Zhong, Zeng & Luo (2007)	10.1107/S1600536806053530	METOQM
<i>(8-Quinolinol-κ<sup>2</sup>N,O)-bis(8-quinolinolato-κ<sup>2</sup>N,O)zinc(II) glyoxal hemisolvate monohydrate</i>	Zhong, Zeng, Luo, Li & Xiao (2007)	10.1107/S1600536807001171	DEXTEG
<i>(Dimethylglyoxime-κ<sup>2</sup>N,N')bis(1,10-phenanthroline-κ<sup>2</sup>N,N')nickel(II) dinitrate dihydrate</i>	Zhong, Zeng, Yang, Luo & Li (2007a)	10.1107/S1600536807004102	YEYGOZ
<i>(Dimethylglyoxime-κ<sup>2</sup>N,N')bis(1,10-phenanthroline-κ<sup>2</sup>N,N')zinc(II) dinitrate dihydrate</i>	Zhong, Zeng, Yang, Luo & Li (2007b)	10.1107/S1600536807004096	YEYGUF
<i>Chlorodibis(1,10-phenanthroline-κN,N')copper(I) hexahydrate</i>	Zhong, Zeng, Yang, Luo & Xiao (2007)	10.1107/S160053680700791X	HEGKOU01
<i>Tetrakis(pyridine-κN)bis(thiocyanato-κN)cobalt(II)</i>	Zhong, Zeng, Yang & Luo (2007a)	10.1107/S1600536807017461	ITCP001
<i>Tetrakis(pyridine-κN)bis(thiocyanato-κN)copper(II)</i>	Zhong, Zeng, Yang & Luo (2007b)	10.1107/S160053680701879X	AVUJEG02
<i>Tetrakis(nitrato-κ<sup>2</sup>O,O')bis(4-phenylpyridine-κN)cerium(IV)</i>	Zhong, Zeng, Yang & Luo (2007c)	10.1107/S1600536807018831	CICDOI
<i>Bis(4,4'-bipyridine-κ<sup>2</sup>N,N')tetrakis(nitrato-κ<sup>2</sup>O,O')cerium(IV)</i>	Zhong, Zeng, Yang & Luo (2007d)	10.1107/S1600536807021502	YIDNEF
<i>(1,10-Phenanthroline)tris(phenoxyacetato)lanthanum(III)</i>	Zhong, Zeng, Yang, Luo & Xu (2007)	10.1107/S1600536807027171	EDUROL
<i>(1,10-Phenanthroline)tris(phenoxyacetato)cerium(III)</i>	Zhong, Yang, Luo & Xu (2007a)	10.1107/S1600536807028061	EDUTUT
<i>(1,10-Phenanthroline)tri(3-phenylpropanoato)lanthanum(III)</i>	Zhong, Yang, Luo & Xu (2007b)	10.1107/S1600536807028693	RIGQEE
<i>(1,10-Phenanthroline-κ<sup>2</sup>N,N')tris(phenoxyacetato)-κO;κO;κO,O'-neodymium(III)</i>	Zhong, Yang, Luo & Xu (2007c)	10.1107/S1600536807030371	UDUMEM
<i>Bis(2,2'-bipyridyl-κ<sup>2</sup>N,N')bis(thiocyanato-κN)nickel(II)</i>	Zhong, Yang, Luo & Xu (2007d)	10.1107/S1600536807031613	YEJGOJ01
<i>Bis(2,2'-bipyridyl-κ<sup>2</sup>N,N')bis(isothiocyanato-κN)copper(II)</i>	Zhong, Yang, Luo & Xu (2007e)	10.1107/S1600536807033181	UFAPOH
<i>Bis(2,2'-bipyridyl-κ<sup>2</sup>N,N')bis(thiocyanato-κN)zinc(II)</i>	Zhong, Yang, Luo & Xu (2007f)	10.1107/S1600536807035337	TIGFAR
<i>(1,10-Phenanthroline-κ<sup>2</sup>N,N')tris(3-phenylpropanoato-κO)neodymium(III)</i>	Zhong, Yang, Luo & Xu (2007g)	10.1107/S1600536807035350	TIGFEV
<i>2-Fluoro-3,5-dinitrobenzamide monohydrate</i>	Zhong, Yang, Xie & Luo (2007j)	10.1107/S1600536807038676	VIKGAY
<i>2-Fluoro-3,5-dinitrobenzoic acid-ammonia (1/1)</i>	Zhong, Yang, Xie & Luo (2007k)	10.1107/S1600536807039724	KILKIA
<i>1-Hydroxy-4,6-dinitropyridine-2-carboxamide monohydrate</i>	Zhong, Yang, Xie & Luo (2007l)	10.1107/S1600536807040779	AFETAH
<i>N-(2-Hydroxyphenyl)carbamic acid-ammonia (1/1)</i>	Zhong, Yang, Xie & Luo (2007m)	10.1107/S160053680704086X	AFINAF
<i>catena-Poly[[bis(μ-anilinoacetato-κ<sup>2</sup>O,O')bis(μ-anilinoacetato-κ<sup>2</sup>O,O')bis(1,10-phenanthroline-κ<sup>2</sup>N,N')samarium(III)]-μ-anilinoacetato-κ<sup>2</sup>O,O']</i>	Zhong, Yang, Xie & Luo (2007a)	10.1107/S1600536807043528	PILDAQ
<i>2-Hydroxy-5-nitrobenzene-1,3-dicarboxylic acid monohydrate</i>	Zhong, Yang, Xie & Luo (2007n)	10.1107/S1600536807045199	XILWIZ
<i>catena-Poly[[tetra-μ-anilinoacetato-bis(1,10-phenanthroline)-neodymium(III)]-di-μ-anilinoacetato]</i>	Zhong, Yang, Xie & Luo (2007b)	10.1107/S1600536807048489	WIMWEV
<i>Hexaaqua copper(II) bis(4-methylbenzenesulfonate)</i>	Zhong, Yang, Xie & Luo (2007c)	10.1107/S1600536807049525	TOLSCV01

# addenda and errata

**Table 1 (continued)**

Title	Reference	DOI	Refcode
<i>catena-Poly[[tetra-<math>\mu</math>-anilinoacetato-bis(1,10-phenanthroline)-dilanthanum(III)]-di-<math>\mu</math>-anilinoacetato]</i>	Zhong, Yang, Xie & Luo (2007d)	10.1107/S1600536807051240	GIMZEI
<i>Hexaaquachromium(II) bis(4-methylbenzenesulfonate)</i>	Zhong, Yang, Xie & Luo (2007e)	10.1107/S1600536807051227	GIMZIM
<i>Hexaaquamanganese(II) bis(4-methylbenzenesulfonate)</i>	Zhong, Yang, Xie & Luo (2007f)	10.1107/S1600536807052051	QUKQES01
<i>catena-Poly[[acetato-<math>\kappa</math>O](1,10-phenanthroline-<math>\kappa^2</math>N,N')cobalt(II)]-<math>\mu</math>-acetato-<math>\kappa^2</math>O:O'</i>	Zhong, Yang, Xie & Luo (2007g)	10.1107/S1600536807053494	NIQLAB
<i>Hexaaquanickel(II) bis(4-aminobenzenesulfonate)</i>	Zhong, Zhong, Xie & Luo (2007a)	10.1107/S1600536807054372	HIPZOW
<i>catena-Poly[[acetato-<math>\kappa</math>O](1,10-phenanthroline-<math>\kappa^2</math>N,N')copper(II)]-<math>\mu</math>-acetato-<math>\kappa^2</math>O:O'</i>	Zhong, Yang, Xie & Luo (2007h)	10.1107/S160053680705622X	XIRGOV
<i>Hexaaquazinc(II) bis(4-aminobenzenesulfonate)</i>	Zhong, Zhong, Xie & Luo (2007b)	10.1107/S1600536807056498	XIRJEQ
<i>catena-Poly[[acetato-<math>\kappa</math>O](1,10-phenanthroline-<math>\kappa^2</math>N,N')nickel(II)]-<math>\mu</math>-acetato-<math>\kappa^2</math>O:O'</i>	Zhong, Yang, Xie & Luo (2007i)	10.1107/S1600536807058540	HIQJOH
<i>Hexaaquacobalt(II) bis(4-aminobenzenesulfonate)</i>	Zhong, Xie & Luo (2007)	10.1107/S1600536807058527	HIQJUN
<i>catena-Poly[[tetra-<math>\mu</math>-anilinoacetato-bis(1,10-phenanthroline)-dieuropium(III)]-di-<math>\mu</math>-anilinoacetato]</i>	Zhong, Yang, Duan & Hong (2007)	10.1107/S1600536807060643	YIQMAN
<i>(Dimethylglyoxime-<math>\kappa^2</math>N,N')bis(1,10-phenanthroline-<math>\kappa^2</math>N,N')copper(II) dinitrate dihydrate</i>	Zhong, Yang, Luo & Li (2007)	10.1107/S1600536807061193	YIQNUI
<i>catena-Poly[[1,10-phenanthroline-<math>\kappa^2</math>N,N')praseodymium(III)]-di-<math>\mu</math>-phenoxyacetato-<math>\kappa^4</math>O:O'-[(1,10-phenanthroline-<math>\kappa^2</math>N,N')-praseodymium(III)]-di-<math>\mu</math>-phenoxyacetato-<math>\kappa^4</math>O:O'-di-<math>\mu</math>-phenoxyacetato-<math>\kappa^3</math>O,O':O;<math>\kappa^3</math>O:O,O'</i>	Zhong, Yang, Luo & Xu (2008)	10.1107/S1600536807068614	GISJIC

## References

- Zhong, H., Xie, H.-L. & Luo, C.-J. (2007). *Acta Cryst.* **E63**, m3054.  
Zhong, H., Yang, X.-M., Duan, S.-H. & Hong, Y.-P. (2007). *Acta Cryst.* **E63**, m3142–m3143.  
Zhong, H., Yang, X.-M., Luo, C.-J. & Li, M.-L. (2007). *Acta Cryst.* **E63**, m3160–m3161.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007a). *Acta Cryst.* **E63**, m1885–m1886.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007b). *Acta Cryst.* **E63**, m1909.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007c). *Acta Cryst.* **E63**, m2019.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007d). *Acta Cryst.* **E63**, m2062.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007e). *Acta Cryst.* **E63**, m2141.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007f). *Acta Cryst.* **E63**, m2208.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007g). *Acta Cryst.* **E63**, m2209–m2210.  
Zhong, H., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2008). *Acta Cryst.* **E64**, m317–m318.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007a). *Acta Cryst.* **E63**, m2508–m2509.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007b). *Acta Cryst.* **E63**, m2680–m2681.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007c). *Acta Cryst.* **E63**, m2724–m2725.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007d). *Acta Cryst.* **E63**, m2772–m2773.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007e). *Acta Cryst.* **E63**, m2774.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007f). *Acta Cryst.* **E63**, m2825.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007g). *Acta Cryst.* **E63**, m2895–m2896.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007h). *Acta Cryst.* **E63**, m2979.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007i). *Acta Cryst.* **E63**, m3053.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007j). *Acta Cryst.* **E63**, o3780.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007k). *Acta Cryst.* **E63**, o3831.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007l). *Acta Cryst.* **E63**, o3881.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007m). *Acta Cryst.* **E63**, o3882.  
Zhong, H., Yang, X.-M., Xie, H.-L. & Luo, C.-J. (2007n). *Acta Cryst.* **E63**, o4191.  
Zhong, H., Zeng, X.-R., Liu, Y.-Q. & Luo, Q.-Y. (2006a). *Acta Cryst.* **E62**, m2925–m2927.  
Zhong, H., Zeng, X.-R., Liu, Y.-Q. & Luo, Q.-Y. (2006b). *Acta Cryst.* **E62**, m3557–m3559.  
Zhong, H., Zeng, X.-R., Liu, Y.-Q. & Luo, Q.-Y. (2007). *Acta Cryst.* **E63**, m187–m189.  
Zhong, H., Zeng, X.-R. & Luo, Q.-Y. (2006). *Acta Cryst.* **E62**, m3330–m3332.  
Zhong, H., Zeng, X.-R. & Luo, Q.-Y. (2007). *Acta Cryst.* **E63**, m221–m223.  
Zhong, H., Zeng, X.-R., Luo, Q.-Y., Li, M.-L. & Xiao, S.-Z. (2007). *Acta Cryst.* **E63**, m492–m494.  
Zhong, H., Zeng, X.-R., Yang, X.-M. & Luo, Q.-Y. (2007a). *Acta Cryst.* **E63**, m1379.  
Zhong, H., Zeng, X.-R., Yang, X.-M. & Luo, Q.-Y. (2007b). *Acta Cryst.* **E63**, m1445.  
Zhong, H., Zeng, X.-R., Yang, X.-M. & Luo, Q.-Y. (2007c). *Acta Cryst.* **E63**, m1455.  
Zhong, H., Zeng, X.-R., Yang, X.-M. & Luo, Q.-Y. (2007d). *Acta Cryst.* **E63**, m1592–m1593.  
Zhong, H., Zeng, X.-R., Yang, X.-M., Luo, Q.-Y. & Li, M.-L. (2007a). *Acta Cryst.* **E63**, m639–m641.  
Zhong, H., Zeng, X.-R., Yang, X.-M., Luo, Q.-Y. & Li, M.-L. (2007b). *Acta Cryst.* **E63**, m642–m644.  
Zhong, H., Zeng, X.-R., Yang, X.-M., Luo, Q.-Y. & Xiao, S.-Z. (2007). *Acta Cryst.* **E63**, m826–m828.  
Zhong, H., Zeng, X.-R., Yang, X.-M., Luo, Q.-Y. & Xu, Y.-P. (2007). *Acta Cryst.* **E63**, m1868–m1869.  
Zhong, H., Zhong, Q.-Y., Xie, H.-L. & Luo, C.-J. (2007a). *Acta Cryst.* **E63**, m2913–m2914.  
Zhong, H., Zhong, Q.-Y., Xie, H.-L. & Luo, C.-J. (2007b). *Acta Cryst.* **E63**, m2990.

## Hexaaqua<sup>2+</sup>zinc(II) bis(4-aminobenzene-sulfonate)

H. Zhong,<sup>a\*</sup> Q.-Y. Zhong,<sup>b</sup> H.-L. Xie<sup>a</sup> and C.-J. Luo<sup>a</sup>

<sup>a</sup>College of Chemistry and Chemical Engineering, Provincial Key Laboratory of Coordination Chemistry, Jinggangshan University, Jian 343009, People's Republic of China, and <sup>b</sup>Jian Training School, Jian 343000, People's Republic of China  
Correspondence e-mail: huazhong06@126.com


Received 2 November 2007; accepted 7 November 2007

Key indicators: single-crystal X-ray study;  $T = 273$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.037;  $wR$  factor = 0.109; data-to-parameter ratio = 14.4.

The asymmetric unit of the title compound,  $[\text{Zn}(\text{H}_2\text{O})_6]\cdot(\text{C}_6\text{H}_5\text{NO}_3\text{S})_2$ , contains one half-cation and one anion; the Zn atom lies on an inversion centre. In the crystal structure, intermolecular O—H···O and O—H···S hydrogen bonds result in the formation of a supramolecular network; an intramolecular C—H···O hydrogen bond is also present.

### Related literature

For a related structure, see: Zhong *et al.* (2007). For bond-length data, see: Allen *et al.* (1987).



### Experimental

#### Crystal data

$[\text{Zn}(\text{H}_2\text{O})_6](\text{C}_6\text{H}_5\text{NO}_3\text{S})_2$

$M_r = 517.82$

Monoclinic,  $P2_1/n$

$a = 6.9902$  (19) Å

$b = 6.2974$  (16) Å

$c = 25.002$  (2) Å

$\beta = 91.449$  (2)°

$V = 1100.2$  (4) Å<sup>3</sup>

$Z = 2$

Mo  $K\alpha$  radiation

$\mu = 1.36$  mm<sup>-1</sup>

$T = 273$  (2) K

$0.49 \times 0.38 \times 0.19$  mm

#### Data collection

Bruker APEXII area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.552$ ,  $T_{\max} = 0.778$

6988 measured reflections  
2265 independent reflections  
1940 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.109$

$S = 1.04$

2265 reflections

157 parameters

9 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.39$  e Å<sup>-3</sup>

$\Delta\rho_{\text{min}} = -0.28$  e Å<sup>-3</sup>

**Table 1**  
Selected geometric parameters (Å, °).

Zn1—O1	2.052 (2)	Zn1—O3	2.042 (2)
Zn1—O2	2.0187 (19)		
O1 <sup>i</sup> —Zn1—O1	180	O2 <sup>i</sup> —Zn1—O2	180
O1—Zn1—O2 <sup>i</sup>	89.06 (9)	O2 <sup>i</sup> —Zn1—O3	89.02 (10)
O1—Zn1—O2	90.94 (9)	O2—Zn1—O3	90.98 (10)
O1—Zn1—O3 <sup>i</sup>	88.00 (12)	O3—Zn1—O3 <sup>i</sup>	180
O1—Zn1—O3	91.99 (12)		

Symmetry code: (i)  $-x + 1, -y, -z$ .

**Table 2**  
Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1B···S1 <sup>ii</sup>	0.837 (18)	3.12 (2)	3.885 (3)	153 (3)
O1—H1A···O4 <sup>iii</sup>	0.782 (17)	1.986 (18)	2.763 (3)	172 (3)
O1—H1B···O5 <sup>ii</sup>	0.837 (18)	2.01 (2)	2.813 (3)	161 (4)
O2—H2A···O4 <sup>ii</sup>	0.806 (17)	1.967 (18)	2.773 (3)	179 (3)
O2—H2B···O6 <sup>iv</sup>	0.795 (17)	1.951 (17)	2.746 (3)	178 (3)
O3—H3A···S1 <sup>v</sup>	0.786 (18)	3.17 (2)	3.921 (2)	162 (3)
O3—H3B···O5 <sup>iv</sup>	0.823 (18)	1.99 (2)	2.797 (3)	168 (4)
O3—H3A···O6 <sup>v</sup>	0.786 (18)	1.992 (18)	2.775 (3)	174 (4)
C3—H3···O5	0.93	2.57	2.927 (4)	103

Symmetry codes: (ii)  $-x + 1, -y + 1, -z + 1$ ; (iii)  $-x + 1, -y, -z + 1$ ; (iv)  $-x, -y + 1, -z + 1$ ; (v)  $-x, -y, -z + 1$ .

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Siemens, 1996); software used to prepare material for publication: *SHELXTL*.

We thank the Science and Technology Program of Jinggangshan University for financial support of this work (grant No. 2007).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2468).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2005). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Siemens (1996). *SAINT* and *SHELXTL*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Zhong, H., Zhong, Q. Y., Xie, H. L. & Luo, C. J. (2007). *Acta Cryst. E* **63**, m2913–m2914.

**supplementary materials**

Article retracted

Acta Cryst. (2007). E63, m2990 [doi:10.1107/S1600536807056498]

### Hexaaquazinc(II) bis(4-aminobenzenesulfonate)

H. Zhong, Q.-Y. Zhong, H.-L. Xie and C.-J. Luo

#### Comment

The crystal structure of Hexaaquanickel(II) bis(4-aminobenzenesulfonate), (II), has previously been reported (Zhong *et al.*, 2007). The crystal structure determination of the title compound, (I), has been carried out in order to elucidate the molecular conformation and to compare it with that of (II). We report herein the crystal structure of (I).

In the molecule of (I) (Fig. 1), the ligand bond lengths and angles are within normal ranges (Allen *et al.*, 1987). The title compound,  $[\text{Zn}(\text{H}_2\text{O})_6](\text{C}_6\text{H}_6\text{NO}_3\text{S})_2$ , contains one half-cation and one anion; the Zn atom lies on an inversion centre, as in (II).

In the crystal structure, intermolecular O—H···O and O—H···S hydrogen bonds (Fig. 2 and Table 2) result in the formation of a supramolecular network structure; an intramolecular C—H···O hydrogen bond is also present, as in (II).

The both compounds, (I) and (II), are isostructural.

#### Experimental

Crystals of the title compound were synthesized using hydrothermal method in a 23 ml Teflon-lined Parr bomb, which was then sealed. Europium (III) nitrate pentahydrate (213.9 mg, 0.5 mmol), zinc nitrate hexahydrate (148.7 mg, 0.5 mmol), *p*-aminobenzenesulfonic acid (346.4 mg, 2 mmol), ammonia (0.5 mol/l, 4 ml) and distilled water (8 g) were placed into the bomb and sealed. The bomb was then heated under autogenous pressure up to 453 K over the course of 7 d and allowed to cool at room temperature for 24 h. Upon opening the bomb, a clear colourless solution was decanted from small colourless crystals. These crystals were washed with distilled water followed by ethanol, and allowed to air-dry at room temperature.

#### Refinement

H atoms (for  $\text{H}_2\text{O}$ ) were located in difference syntheses and refined isotropically [ $\text{O}—\text{H} = 0.782$  (17) and 0.837 (18) Å,  $U_{\text{iso}}(\text{H}) = 0.050$  (9) and 0.075 (8) Å<sup>2</sup>]. The other H atoms were positioned geometrically, with  $\text{N}—\text{H} = 0.86$  Å (for  $\text{NH}_2$ ) and  $\text{C}—\text{H} = 0.93$  Å for aromatic H, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C},\text{N})$  for aromatic and amino H atoms, or  $1.5U_{\text{eq}}(\text{O})$  for water molecules' H atoms.

#### Figures

Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level [symmetry code (A): 3/2 -  $x$ , 1/2 ±  $y$ , 1/2 -  $z$ ].

# supplementary materials



Fig. 2. A packing diagram of (I). Hydrogen bonds are shown as dashed lines.

## Hexaaquazinc(II) bis(4-aminobenzenesulfonate)

### Crystal data

$[\text{Zn}(\text{H}_2\text{O})_6](\text{C}_6\text{H}_6\text{NO}_3\text{S})_2$

$M_r = 517.82$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 6.9902$  (19) Å

$b = 6.2974$  (16) Å

$c = 25.002$  (2) Å

$\beta = 91.449$  (2)°

$V = 1100.2$  (4) Å<sup>3</sup>

$Z = 2$

$F_{000} = 536$

$D_x = 1.563 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 5547 reflections

$\theta = 2.1\text{--}26.6^\circ$

$\mu = 1.36 \text{ mm}^{-1}$

$T = 273$  (2) K

Prism, colourless

$0.49 \times 0.38 \times 0.19$  mm

### Data collection

Bruker APEX-II area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 273$ (2) K

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.552$ ,  $T_{\max} = 0.778$

6988 measured reflections

2265 independent reflections

1940 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 26.8^\circ$

$\theta_{\min} = 3.3^\circ$

$h = -8 \rightarrow 8$

$k = -7 \rightarrow 7$

$l = -31 \rightarrow 31$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.109$

$S = 1.04$

2265 reflections

157 parameters

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0655P)^2 + 0.444P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.39 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.27 \text{ e \AA}^{-3}$

9 restraints

Extinction correction: none

Primary atom site location: structure-invariant direct  
methods*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.0000	0.0000	0.04348 (17)
S1	-0.01480 (9)	0.60833 (10)	0.90519 (3)	0.04055 (19)
O1	0.7237 (4)	-0.1030 (4)	0.04800 (11)	0.0715 (7)
O2	0.5086 (3)	0.2886 (3)	0.03511 (9)	0.0527 (5)
O3	0.3033 (4)	-0.1032 (4)	0.05310 (12)	0.0714 (7)
O4	0.1556 (3)	0.5131 (3)	0.93092 (9)	0.0525 (5)
O5	-0.0143 (3)	0.8403 (3)	0.90891 (8)	0.0518 (5)
O6	-0.1927 (3)	0.5160 (3)	0.92413 (9)	0.0533 (5)
N1	0.0313 (7)	0.3787 (11)	0.67178 (14)	0.1237 (17)
H1AA	0.0048	0.4709	0.6473	0.148*
H1BB	0.0684	0.2534	0.6631	0.148*
C1	0.0126 (5)	0.4361 (7)	0.72961 (14)	0.0676 (9)
C2	-0.0455 (5)	0.6320 (7)	0.74500 (14)	0.0738 (10)
H2	-0.0808	0.7305	0.7188	0.089*
C3	-0.0542 (5)	0.6905 (6)	0.79837 (13)	0.0633 (8)
H3	-0.0934	0.8262	0.8079	0.076*
C4	-0.0037 (4)	0.5444 (5)	0.83687 (11)	0.0447 (6)
C5	0.0529 (5)	0.3427 (5)	0.82224 (12)	0.0595 (8)
H5	0.0847	0.2427	0.8483	0.071*
C6	0.0621 (6)	0.2902 (6)	0.76880 (13)	0.0702 (9)
H6	0.1020	0.1551	0.7591	0.084*
H1A	0.756 (4)	-0.221 (3)	0.0511 (12)	0.075 (8)*
H2A	0.606 (3)	0.347 (5)	0.0453 (12)	0.053 (9)*
H3A	0.271 (5)	-0.217 (3)	0.0619 (15)	0.073 (12)*
H1B	0.819 (4)	-0.025 (4)	0.0535 (15)	0.070 (10)*
H2B	0.419 (3)	0.347 (5)	0.0473 (11)	0.050 (9)*
H3B	0.219 (4)	-0.017 (4)	0.0598 (15)	0.062 (11)*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.0416 (3)	0.0326 (3)	0.0562 (3)	0.00076 (16)	0.00184 (18)	0.00150 (17)
S1	0.0403 (3)	0.0304 (3)	0.0509 (4)	0.0005 (2)	0.0026 (2)	-0.0005 (2)
O1	0.0702 (15)	0.0340 (12)	0.1080 (19)	0.0050 (11)	-0.0417 (14)	0.0012 (12)
O2	0.0447 (11)	0.0364 (11)	0.0771 (14)	-0.0004 (9)	0.0052 (10)	-0.0139 (9)
O3	0.0753 (16)	0.0350 (12)	0.1060 (19)	-0.0001 (11)	0.0465 (14)	0.0061 (12)
O4	0.0560 (12)	0.0394 (11)	0.0613 (12)	0.0057 (8)	-0.0120 (10)	-0.0008 (8)
O5	0.0533 (11)	0.0301 (10)	0.0722 (13)	0.0007 (8)	0.0043 (9)	-0.0018 (9)
O6	0.0542 (11)	0.0388 (11)	0.0677 (13)	-0.0041 (8)	0.0178 (10)	-0.0019 (8)
N1	0.116 (3)	0.193 (5)	0.063 (2)	0.010 (3)	0.011 (2)	-0.004 (3)
C1	0.0522 (17)	0.094 (3)	0.0566 (19)	-0.0061 (18)	0.0016 (14)	-0.0020 (18)
C2	0.066 (2)	0.097 (3)	0.0590 (18)	0.007 (2)	0.0008 (16)	0.0232 (19)
C3	0.0640 (18)	0.060 (2)	0.0657 (19)	0.0094 (15)	0.0040 (15)	0.0136 (16)
C4	0.0394 (13)	0.0442 (15)	0.0504 (14)	0.0004 (11)	0.0011 (11)	0.0020 (11)
C5	0.074 (2)	0.0495 (18)	0.0551 (16)	0.0088 (15)	-0.0006 (14)	-0.0025 (14)
C6	0.081 (2)	0.072 (2)	0.0584 (18)	0.0041 (19)	0.0054 (16)	-0.0145 (17)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Zn1—O1	2.052 (2)	O3—H3B	0.823 (18)
Zn1—O2	2.0187 (19)	N1—C1	1.499 (5)
Zn1—O3	2.042 (2)	N1—H1AA	0.8600
Zn1—O2 <sup>i</sup>	2.0187 (19)	N1—H1BB	0.8600
Zn1—O3 <sup>i</sup>	2.042 (2)	C1—C2	1.358 (6)
Zn1—O1 <sup>i</sup>	2.052 (2)	C1—C6	1.381 (6)
S1—O6	1.463 (2)	C2—C3	1.387 (5)
S1—O5	1.464 (2)	C2—H2	0.9300
S1—O4	1.468 (2)	C3—C4	1.371 (4)
S1—C4	1.758 (3)	C3—H3	0.9300
O1—H1A	0.782 (17)	C4—C5	1.382 (4)
O1—H1B	0.837 (18)	C5—C6	1.379 (4)
O2—H2A	0.806 (17)	C5—H5	0.9300
O2—H2B	0.795 (17)	C6—H6	0.9300
O3—H3A	0.786 (18)		
O1 <sup>i</sup> —Zn1—O1	180.0	H2A—O2—H2B	110 (2)
O1—Zn1—O2 <sup>i</sup>	89.06 (9)	Zn1—O3—H3A	132 (3)
O1—Zn1—O2	90.94 (9)	Zn1—O3—H3B	115 (2)
O1—Zn1—O3 <sup>i</sup>	88.00 (12)	H3A—O3—H3B	110 (3)
O1—Zn1—O3	91.99 (12)	C1—N1—H1AA	120.0
O2 <sup>i</sup> —Zn1—O2	180.0	C1—N1—H1BB	120.0
O2 <sup>i</sup> —Zn1—O3	89.02 (10)	H1AA—N1—H1BB	120.0
O2—Zn1—O3	90.98 (10)	C2—C1—C6	118.4 (3)
O3—Zn1—O3 <sup>i</sup>	180.0	C2—C1—N1	121.8 (4)
O2 <sup>i</sup> —Zn1—O1 <sup>i</sup>	90.94 (9)	C6—C1—N1	119.8 (4)

O2 <sup>i</sup> —Zn1—O3 <sup>i</sup>	90.98 (10)	C1—C2—C3	122.3 (3)
O2—Zn1—O3 <sup>i</sup>	89.02 (10)	C1—C2—H2	118.8
O3 <sup>i</sup> —Zn1—O1 <sup>i</sup>	91.99 (12)	C3—C2—H2	118.8
O2 <sup>i</sup> —Zn1—O1	89.06 (9)	C4—C3—C2	118.7 (3)
O3 <sup>i</sup> —Zn1—O1	88.01 (12)	C4—C3—H3	120.6
O6—S1—O5	112.14 (12)	C2—C3—H3	120.6
O6—S1—O4	112.52 (13)	C3—C4—C5	120.1 (3)
O5—S1—O4	112.31 (12)	C3—C4—S1	120.8 (2)
O6—S1—C4	106.39 (13)	C5—C4—S1	119.2 (2)
O5—S1—C4	106.86 (13)	C6—C5—C4	119.8 (3)
O4—S1—C4	106.09 (13)	C6—C5—H5	120.1
Zn1—O1—H1A	125 (2)	C4—C5—H5	120.1
Zn1—O1—H1B	120 (2)	C5—C6—C1	120.7 (4)
H1A—O1—H1B	109 (2)	C5—C6—H6	119.6
Zn1—O2—H2A	124 (2)	C1—C6—H6	119.6
Zn1—O2—H2B	124 (2)		

Symmetry codes: (i)  $-x+1, -y, -z$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )


$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1—H1B···S1 <sup>ii</sup>	0.837 (18)	3.12 (2)	3.885 (3)	153 (3)
O1—H1A···O4 <sup>iii</sup>	0.782 (17)	1.986 (18)	2.763 (3)	172 (3)
O1—H1B···O5 <sup>ii</sup>	0.837 (18)	2.01 (2)	2.813 (3)	161 (4)
O2—H2A···O4 <sup>ii</sup>	0.806 (17)	1.967 (18)	2.773 (3)	179 (3)
O2—H2B···O6 <sup>iv</sup>	0.795 (17)	1.951 (17)	2.746 (3)	178 (3)
O3—H3A···S1 <sup>v</sup>	0.786 (18)	3.17 (2)	3.921 (2)	162 (3)
O3—H3B···O5 <sup>iv</sup>	0.823 (18)	1.99 (2)	2.797 (3)	168 (4)
O3—H3A···O6 <sup>v</sup>	0.786 (18)	1.992 (18)	2.775 (3)	174 (4)
C3—H3···O5	0.93	2.57	2.927 (4)	103

Symmetry codes: (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $-x+1, -y, -z+1$ ; (iv)  $-x, -y+1, -z+1$ ; (v)  $-x, -y, -z+1$ .

## **supplementary materials**


---

**Fig. 1**



Article retracted

Fig. 2



Article retrac