

Retraction of articles by T. Liu *et al.*

T. Liu,^{a*} Y.-X. Wang,^b Z.-W. Wang,^a Z.-P. Xie^{a,c} and J. Y. Zhu^d

^aCollege of Engineering, Jinggangshan University, Jian 343009, People's Republic of China, ^bCollege of Mathematics and Physics, Jinggangshan University, Jian

343009, People's Republic of China, ^cDepartment of Chemistry, Jiangxi University of Science and Technology, Ganzhou 341000, People's Republic of China, and

^dDepartment of Information Engineering, Jiangxi University of Science and Technology, Nanchang 330013, People's Republic of China

Correspondence e-mail: taoliu07@126.com

Received 20 November 2009; accepted 15 December 2009

A series of 29 papers by Liu *et al.* are retracted.

As a result of problems with the data sets and incorrect atom assignments, 29 papers by Liu *et al.* are retracted. Full details of all the articles are given in Table 1.

Table 1

Details of articles to be retracted, in order of publication.

Title	Reference	DOI	Refcode
<i>Tetrakis(pyrazine-κN)bis(thiocyanato-κN)manganese(II)</i> <i>(Dihydroxyglyoxime-κ²N,N')bis(I,10-phenanthroline-κ²N,N')copper(II) dinitrate dihydrate</i>	Liu & Xie (2007a) Liu, Wang, Wang & Xie (2007b)	10.1107/S1600536807026852 10.1107/S1600536807028255	EDUMAS EDUVAB
<i>Tetrakis(pyrazine-κN)bis(thiocyanato-κN)zinc(II)</i>	Liu & Xie (2007b)	10.1107/S1600536807028735	RIGQAA
<i>Tetrakis(μ-2-pyridyloxyacetato)bis[(I,10-phenanthroline)(2-pyridyloxyacetato)-lanthanum(III)]</i>	Liu, Wang, Wang & Xie (2007c)	10.1107/S1600536807030917	UDUMIQ
<i>Polymeric KNO₂</i> <i>(Dihydroxyglyoxime-κ²N,N')bis(I,10-phenanthroline-κ²N,N')cobalt(II) dinitrate dihydrate</i>	Liu Wang, Wang & Xie (2007a) Liu, Wang, Wang & Xie (2007d)	10.1107/S1600536807027195 10.1107/S1600536807031224	ICSD 240891 WIHIED
<i>Tetrakis(μ-2-pyridyloxyacetato)bis[(I,10-phenanthroline)(2-pyridyloxyacetato)-praseodymium(III)]</i>	Liu, Wang, Wang & Xie (2007e)	10.1107/S1600536807032679	WIHQEK
<i>Tetrakis[μ-(2-pyridyloxyacetato-κ²O:O')bis(I,10-phenanthroline-κ²N,N')-(2-pyridyloxyacetato-κO)neodymium(III)]</i>	Liu, Wang, Wang & Xie (2007f)	10.1107/S1600536807035349	TIGDAP
<i>(Dihydroxyglyoxime-κ²N,N')bis(I,10-phenanthroline-κ²N,N')manganese(II) dinitrate dihydrate</i>	Liu, Wang, Wang & Xie (2007g)	10.1107/S1600536807035076	TIGDET
<i>2-Amino-3,5-dinitrobenzoic acid-ammonium (I/I)</i>	Liu & Zhu (2007j)	10.1107/S1600536807040068	KIKQAX
<i>2-Hydroxy-3,5-dinitrobenzamide monohydrate</i>	Liu & Zhu (2007k)	10.1107/S1600536807039712	KIKQEB
<i>2-(1-Hydroxy-2-pyridyl)acetamide monohydrate</i>	Liu & Zhu (2007l)	10.1107/S1600536807040652	CIKQOD
<i>Bis(2,2'-bipyridine-κN,N')bis(thiocyanato-κN)iron(II)</i>	Liu & Zhu (2007a)	10.1107/S1600536807043486	XIFXOA
<i>catena-Poly[hexakis(μ₂-anilinoacetamide)bis(I,10-phenanthroline)disamarium(III)]</i>	Liu & Zhu (2007b)	10.1107/S1600536807045485	XILNAI
<i>3-Hydroxy-2,4,6-trinitropyridine monohydrate</i>	Liu & Zhu (2007m)	10.1107/S1600536807045230	PILNOO
<i>catena-Poly[hexakis(μ₂-anilinoacetamide)bis(I,10-phenanthroline)-dipraseodymium(III)]</i>	Liu & Zhu (2007c)	10.1107/S1600536807047733	SILZET
<i>catena-Poly[[tetra-μ-anilinoacetamido-bis(I,10-phenanthroline)dicerium(III)-di-μ-anilinoacetamido]</i>	Liu & Zhu (2007d)	10.1107/S1600536807050969	GIMZOS
<i>Tetrakis(pyridine-κN)bis(thiocyanato-κN)chromium(II)</i>	Liu & Zhu (2007e)	10.1107/S1600536807051756	WINFAB
<i>2-Ammonio-3-carboxy-5-nitrobenzoate monohydrate</i>	Liu & Zhu (2007n)	10.1107/S1600536807048477	GINFEP
<i>2-(Benzoylhydrazinocarbonyl)benzoic acid</i>	Liu & Zhu (2007o)	10.1107/S160053680705204X	TINZIA
<i>Tetrakis(pyridine-κN)bis(thiocyanato-κN)vanadium(II)</i>	Liu & Zhu (2007f)	10.1107/S1600536807054529	HIPZIQ
<i>catena-Poly[[nitrato-κO](I,10-phenanthroline-κ²N,N')nickel(II)-μ-acetamido-κ²O:N]</i>	Liu & Zhu (2007g)	10.1107/S1600536807056504	XIRGIP
<i>catena-Poly[[nitrato-κO](I,10-phenanthroline-κ²N,N')copper(II)-μ-acetamido-κ²O:N]</i>	Liu & Zhu (2007h)	10.1107/S1600536807059077	HIQROP
<i>catena-Poly[[nitrato-κO](I,10-phenanthroline-κ²N,N')cobalt(II)-μ-acetamido-κ²O:N]</i>	Liu & Zhu (2007i)	10.1107/S1600536807060631	YIQMER
<i>N'-Benzoyl-4-nitronicotinohydrazide</i>	Liu & Zhu (2007p)	10.1107/S1600536807053068	CIPVON
<i>N'-(3-Nitro-4-pyridylcarbonyl)pyridine-4-carbohydrazide</i>	Liu & Zhu (2007q)	10.1107/S1600536807054876	RIRWEV

addenda and errata

Table 1 (continued)

Title	Reference	DOI	Refcode
Ethylenediammonium sulfate	Liu & Zhu (2007r)	10.1107/S1600536807056280	ETDAMS03
Ethylenediammonium perchlorate	Liu & Zhu (2007s)	10.1107/S1600536807059909	HIRYEN
catena-Poly[μ (nitro- κO)(1,10-phenanthroline- $\kappa^2 N,N'$)manganese(II)]- μ -nitroato- $\kappa^2 O:O'$]	Liu & Zhu (2008)	10.1107/S160053680706254X	MIRROV

References

- Liu, T., Wang, Z.-W., Wang, Y.-X. & Xie, Z.-P. (2007a). *Acta Cryst.* E63, i170.
Liu, T., Wang, Z.-W., Wang, Y.-X. & Xie, Z.-P. (2007b). *Acta Cryst.* E63, m1887–m1888.
Liu, T., Wang, Z.-W., Wang, Y.-X. & Xie, Z.-P. (2007c). *Acta Cryst.* E63, m2020–m2021.
Liu, T., Wang, Z.-W., Wang, Y.-X. & Xie, Z.-P. (2007d). *Acta Cryst.* E63, m2027–m2028.
Liu, T., Wang, Z.-W., Wang, Y.-X. & Xie, Z.-P. (2007e). *Acta Cryst.* E63, m2080–m2081.
Liu, T., Wang, Z.-W., Wang, Y.-X. & Xie, Z.-P. (2007f). *Acta Cryst.* E63, m2196–m2197.
Liu, T., Wang, Z.-W., Wang, Y.-X. & Xie, Z.-P. (2007g). *Acta Cryst.* E63, m2198–m2199.
Liu, T. & Xie, Z.-P. (2007a). *Acta Cryst.* E63, m1820.
Liu, T. & Xie, Z.-P. (2007b). *Acta Cryst.* E63, m1908.
Liu, T. & Zhu, J. Y. (2007a). *Acta Cryst.* E63, m2506–m2507.
Liu, T. & Zhu, J.-Y. (2007b). *Acta Cryst.* E63, m2592–m2593.
Liu, T. & Zhu, J. Y. (2007c). *Acta Cryst.* E63, m2659–m2660.
Liu, T. & Zhu, J. Y. (2007d). *Acta Cryst.* E63, m2775–m2776.
Liu, T. & Zhu, J. Y. (2007e). *Acta Cryst.* E63, m2809.
Liu, T. & Zhu, J. Y. (2007f). *Acta Cryst.* E63, m2912.
Liu, T. & Zhu, J.-Y. (2007g). *Acta Cryst.* E63, m2977–m2978.
Liu, T. & Zhu, J. Y. (2007h). *Acta Cryst.* E63, m3108.
Liu, T. & Zhu, J. Y. (2007i). *Acta Cryst.* E63, m3144.
Liu, T. & Zhu, J.-Y. (2007j). *Acta Cryst.* E63, o3829.
Liu, T. & Zhu, J. Y. (2007k). *Acta Cryst.* E63, o3830.
Liu, T. & Zhu, J.-Y. (2007l). *Acta Cryst.* E63, o3860.
Liu, T. & Zhu, J. Y. (2007m). *Acta Cryst.* E63, o4112.
Liu, T. & Zhu, J. Y. (2007n). *Acta Cryst.* E63, o4267.
Liu, T. & Zhu, J. Y. (2007o). *Acta Cryst.* E63, o4441.
Liu, T. & Zhu, J. Y. (2007p). *Acta Cryst.* E63, o4527.
Liu, T. & Zhu, J. Y. (2007q). *Acta Cryst.* E63, o4574.
Liu, T. & Zhu, J.-Y. (2007r). *Acta Cryst.* E63, o4660.
Liu, T. & Zhu, J. Y. (2007s). *Acta Cryst.* E63, o4874.
Liu, T. & Zhu, J. Y. (2008). *Acta Cryst.* E64, m28.

N'-Benzoyl-4-nitronicotinohydrazide**T. Liu^{a*} and J. Y. Zhu^b**

^aCollege of Engineering, Jinggangshan University, Jian 343009, People's Republic of China, and ^bDepartment of Information Engineering, Jiangxi University of Science and Technology, Nanchang 330013, People's Republic of China

Correspondence e-mail: taoliu07@126.com


Received 20 October 2007; accepted 25 October 2007

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.049; wR factor = 0.139; data-to-parameter ratio = 13.5.

In the crystal structure of the title compound, $\text{C}_{13}\text{H}_{10}\text{N}_4\text{O}_4$, all bond lengths and angles are within normal ranges and the benzene and pyridine rings are oriented at a dihedral angle of $7.988(4)^\circ$. $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds result in the formation of a supramolecular network.

Related literature

For general background, see: Desiraju (1995, 1997); Braga *et al.* (1998); Li *et al.* (2005); Liu *et al.* (2004); Pan & Xu (2004); Wu *et al.* (2003). For bond-length data, see: Allen *et al.* (1987).

**Experimental***Crystal data* $\text{C}_{13}\text{H}_{10}\text{N}_4\text{O}_4$ $M_r = 286.25$ Monoclinic, $P2_1/n$ $a = 8.290(2)\text{ \AA}$ $b = 12.7981(13)\text{ \AA}$ $c = 13.0073(11)\text{ \AA}$ $\beta = 92.804(3)^\circ$ $V = 1378.4(4)\text{ \AA}^3$ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.11\text{ mm}^{-1}$ $T = 273(2)\text{ K}$ $0.28 \times 0.17 \times 0.15\text{ mm}$ **Data collection**

Bruker APEXII area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.971$, $T_{\max} = 0.984$

8484 measured reflections
 2586 independent reflections
 1276 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.139$
 $S = 1.01$
 2586 reflections

191 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.38\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N4—H4A \cdots O1 ⁱ	0.86	2.11	2.903 (3)	153
N3—H3A \cdots O2 ⁱⁱ	0.86	1.98	2.831 (3)	170

Symmetry codes: (i) $-x + 2, -y, -z + 1$; (ii) $-x + 1, -y, -z + 1$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Siemens, 1996); software used to prepare material for publication: *SHELXTL*.

We thank the Youth Program of Jinggangshan University for financial support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2444).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Braga, D., Grepioni, F. & Desiraju, G. R. (1998). *Chem. Rev.* **98**, 1375–1386.
- Bruker (2005). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 2311–2315.
- Desiraju, G. R. (1997). *J. Chem. Soc. Chem. Commun.*, pp. 1475–1476.
- Liu, B.-X., Su, J.-R. & Xu, D.-J. (2004). *Acta Cryst. C* **60**, m183–m185.
- Li, H., Yin, K.-L. & Xu, D.-J. (2005). *Acta Cryst. C* **61**, m19–m21.
- Pan, T.-T. & Xu, D.-J. (2004). *Acta Cryst. E* **60**, m56–m58.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Siemens (1996). *SAINT* and *SHELXTL*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Wu, Z.-Y., Xue, Y.-H. & Xu, D.-J. (2003). *Acta Cryst. E* **59**, m809–m811.

supplementary materials

Article retracted

Acta Cryst. (2007). E63, o4527 [doi:10.1107/S1600536807053068]

N'-Benzoyl-4-nitronicotinohydrazide

T. Liu and J. Y. Zhu

Comment

In the synthesis of crystal structures by design, the assembly of molecular units in predefined arrangements is a key goal (Desiraju, 1995, 1997; Braga *et al.*, 1998). Due to pyridine groups are one of the most important classes of biological ligands, the coordination of metal-pyridine groups complexes are of critical importance in biological systems, organic materials and coordination chemistry. Recently, pyridine groups with variable coordination modes have been used to construct metal-organic supramolecular structures (Wu *et al.*, 2003; Pan & Xu, 2004; Liu *et al.*, 2004; Li *et al.*, 2005). We originally attempted to synthesize complexes featuring Cu metal chains by reaction of the copper(II) ion with *N*-(4-nitropyridine-3-carboxamido)-benzamide ligand. Unfortunately, we obtained only the title compound, (I), and we report herein its crystal structure.

In the molecule of (I) (Fig. 1), the ligand bond lengths and angles are within normal ranges (Allen *et al.*, 1987). In the crystal structure, the N—H···O hydrogen bonds (Table 1, Fig. 2) result in the formation of a supramolecular network structure.

Experimental

Crystals of the title compound were synthesized using hydrothermal method in a 23 ml Teflon-lined Parr bomb, which was then sealed. Copper dinitrate hexahydrate (147.8 mg, 0.5 mmol), *N*-(4-nitropyridine-3-carboxamido)-benzamide (286.3 mg, 1 mmol) and distilled water (5 g) were placed into the bomb and sealed. The bomb was then heated under autogenous pressure up to 453 K over the course of 7 d and allowed to cool at room temperature for 24 h. Upon opening the bomb, a clear colourless solution was decanted from small colourless crystals. These crystals were washed with distilled water followed by ethanol, and allowed to air-dry at room temperature.

Refinement

The H atoms were positioned geometrically, with N—H = 0.86 Å (for NH) and C—H = 0.93 Å for aromatic H, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

supplementary materials



Fig. 2. A packing diagram of (I). Hydrogen bonds are shown as dashed lines.

N¹-Benzoyl-4-nitronicotinohydrazide

Crystal data

C₁₃H₁₀N₄O₄

M_r = 286.25

Monoclinic, P2₁/n

Hall symbol: -P 2yn

a = 8.290 (2) Å

b = 12.7981 (13) Å

c = 13.0073 (11) Å

β = 92.804 (3)°

V = 1378.4 (4) Å³

Z = 4

F₀₀₀ = 592

D_x = 1.379 Mg m⁻³

Mo Kα radiation

λ = 0.71073 Å

Cell parameters from 1769 reflections

θ = 2.3–21.2°

μ = 0.11 mm⁻¹

T = 273 (2) K

Prism, colourless

0.28 × 0.17 × 0.15 mm

Data collection

Bruker APEXII area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 273(2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

T_{min} = 0.971, T_{max} = 0.984

8484 measured reflections

2586 independent reflections

1276 reflections with I > 2σ(I)

R_{int} = 0.043

θ_{max} = 26.0°

θ_{min} = 2.2°

h = -10→10

k = -15→15

l = -16→15

Refinement

Refinement on F²

Hydrogen site location: inferred from neighbouring sites

Least-squares matrix: full

H-atom parameters constrained

R[F² > 2σ(F²)] = 0.049

w = 1/[σ²(F_o²) + (0.0593P)² + 0.15P]

where P = (F_o² + 2F_c²)/3

wR(F²) = 0.139

(Δ/σ)_{max} < 0.001

S = 1.01

Δρ_{max} = 0.38 e Å⁻³

2586 reflections

Δρ_{min} = -0.26 e Å⁻³

191 parameters

Extinction correction: SHELXL97 (Sheldrick, 1997),

F_c* = kF_c[1+0.001xF_c²λ³/sin(2θ)]^{1/4}

Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.0085 (16)

Secondary atom site location: difference Fourier map

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O3	0.7539 (4)	-0.37730 (18)	0.5475 (2)	0.1317 (10)
O4	0.8686 (3)	-0.23377 (16)	0.51348 (15)	0.0856 (7)
O1	0.9288 (2)	-0.01114 (13)	0.62696 (13)	0.0636 (5)
O2	0.5682 (2)	0.15823 (13)	0.51217 (15)	0.0682 (6)
N3	0.7074 (3)	-0.02935 (15)	0.51904 (16)	0.0626 (6)
H3A	0.6180	-0.0619	0.5061	0.075*
N4	0.7559 (3)	0.04934 (15)	0.45320 (16)	0.0632 (6)
H4A	0.8303	0.0384	0.4107	0.076*
N2	0.7830 (3)	-0.28636 (18)	0.56606 (18)	0.0714 (7)
C1	0.7145 (3)	-0.24006 (17)	0.65641 (18)	0.0529 (6)
C2	0.6407 (3)	-0.30418 (18)	0.7262 (2)	0.0609 (7)
H2	0.6335	-0.3756	0.7138	0.073*
C3	0.5793 (3)	-0.2641 (2)	0.8121 (2)	0.0631 (7)
H3	0.5313	-0.3082	0.8585	0.076*
N1	0.5872 (3)	-0.1581 (2)	0.8315 (2)	0.0929 (8)
C4	0.6614 (3)	-0.0939 (2)	0.76155 (19)	0.0611 (7)
H4	0.6682	-0.0226	0.7749	0.073*
C5	0.7258 (3)	-0.13250 (17)	0.67237 (18)	0.0508 (6)
C6	0.8002 (3)	-0.05385 (17)	0.6022 (2)	0.0512 (6)
C7	0.6825 (3)	0.14358 (19)	0.45821 (19)	0.0537 (6)
C8	0.7451 (3)	0.22485 (19)	0.38880 (19)	0.0563 (7)
C9	0.8229 (3)	0.2020 (2)	0.3001 (2)	0.0704 (8)
H9	0.8404	0.1328	0.2822	0.085*
C10	0.8756 (4)	0.2827 (3)	0.2370 (3)	0.0913 (10)
H10	0.9276	0.2677	0.1770	0.110*
C11	0.8490 (5)	0.3840 (3)	0.2656 (4)	0.1109 (14)
H11	0.8825	0.4382	0.2240	0.133*
C12	0.7744 (5)	0.4065 (2)	0.3537 (4)	0.1046 (13)
H12	0.7616	0.4758	0.3730	0.126*
C13	0.7176 (4)	0.3280 (2)	0.4150 (3)	0.0803 (9)

supplementary materials

H13 0.6615 0.3442 0.4731 0.096*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O3	0.216 (3)	0.0550 (14)	0.130 (2)	-0.0236 (15)	0.0768 (18)	-0.0343 (13)
O4	0.1122 (17)	0.0719 (13)	0.0763 (14)	0.0048 (12)	0.0411 (12)	-0.0029 (11)
O1	0.0679 (12)	0.0460 (10)	0.0778 (12)	-0.0083 (9)	0.0122 (9)	-0.0001 (8)
O2	0.0728 (13)	0.0579 (12)	0.0753 (13)	-0.0045 (9)	0.0167 (10)	0.0018 (9)
N3	0.0641 (14)	0.0513 (12)	0.0728 (15)	-0.0121 (11)	0.0072 (12)	0.0217 (11)
N4	0.0730 (15)	0.0479 (13)	0.0704 (14)	-0.0018 (11)	0.0212 (11)	0.0171 (10)
N2	0.0987 (19)	0.0471 (14)	0.0704 (16)	0.0043 (13)	0.0260 (14)	-0.0045 (12)
C1	0.0622 (16)	0.0395 (13)	0.0576 (15)	0.0054 (12)	0.0094 (12)	0.0006 (11)
C2	0.0760 (19)	0.0363 (13)	0.0711 (18)	0.0008 (12)	0.0104 (14)	0.0096 (12)
C3	0.082 (2)	0.0484 (16)	0.0599 (17)	0.0002 (14)	0.0158 (14)	0.0135 (13)
N1	0.110 (2)	0.084 (2)	0.0866 (18)	0.0032 (16)	0.0166 (15)	0.0102 (14)
C4	0.0780 (19)	0.0450 (14)	0.0621 (17)	-0.0015 (13)	0.0217 (14)	0.0005 (12)
C5	0.0562 (16)	0.0389 (13)	0.0578 (16)	0.0009 (11)	0.0089 (12)	0.0045 (11)
C6	0.0586 (16)	0.0337 (13)	0.0627 (17)	0.0002 (13)	0.0171 (13)	-0.0005 (11)
C7	0.0573 (17)	0.0486 (16)	0.0551 (16)	-0.0056 (13)	0.0019 (13)	0.0028 (12)
C8	0.0595 (17)	0.0479 (15)	0.0610 (17)	-0.0084 (12)	-0.0009 (13)	0.0109 (12)
C9	0.073 (2)	0.0670 (18)	0.071 (2)	-0.0065 (15)	0.0033 (15)	0.0219 (15)
C10	0.084 (2)	0.110 (3)	0.080 (2)	-0.015 (2)	0.0005 (17)	0.041 (2)
C11	0.123 (3)	0.086 (3)	0.119 (3)	-0.041 (2)	-0.030 (3)	0.057 (3)
C12	0.149 (4)	0.0486 (19)	0.113 (3)	-0.026 (2)	-0.025 (3)	0.023 (2)
C13	0.098 (2)	0.0501 (18)	0.091 (2)	-0.0077 (15)	-0.0154 (17)	0.0086 (15)

Geometric parameters (\AA , $^\circ$)

O3—N2	1.211 (3)	N1—C4	1.392 (3)
O4—N2	1.213 (3)	C4—C5	1.391 (3)
O1—C6	1.227 (3)	C4—H4	0.9300
O2—C7	1.220 (3)	C5—C6	1.511 (3)
N3—C6	1.333 (3)	C7—C8	1.487 (3)
N3—N4	1.394 (3)	C8—C9	1.380 (4)
N3—H3A	0.8600	C8—C13	1.385 (4)
N4—C7	1.354 (3)	C9—C10	1.402 (4)
N4—H4A	0.8600	C9—H9	0.9300
N2—C1	1.456 (3)	C10—C11	1.370 (6)
C1—C2	1.388 (3)	C10—H10	0.9300
C1—C5	1.395 (3)	C11—C12	1.360 (5)
C2—C3	1.351 (4)	C11—H11	0.9300
C2—H2	0.9300	C12—C13	1.379 (5)
C3—N1	1.381 (3)	C12—H12	0.9300
C3—H3	0.9300	C13—H13	0.9300
C6—N3—N4	119.7 (2)	O1—C6—N3	124.6 (2)
C6—N3—H3A	120.1	O1—C6—C5	120.9 (2)
N4—N3—H3A	120.1	N3—C6—C5	114.2 (2)

C7—N4—N3	118.0 (2)	O2—C7—N4	121.9 (2)
C7—N4—H4A	121.0	O2—C7—C8	123.1 (2)
N3—N4—H4A	121.0	N4—C7—C8	115.0 (2)
O4—N2—O3	122.5 (2)	C9—C8—C13	119.8 (2)
O4—N2—C1	119.4 (2)	C9—C8—C7	123.4 (2)
O3—N2—C1	118.1 (2)	C13—C8—C7	116.8 (3)
C2—C1—C5	121.0 (2)	C8—C9—C10	120.4 (3)
C2—C1—N2	119.3 (2)	C8—C9—H9	119.8
C5—C1—N2	119.7 (2)	C10—C9—H9	119.8
C3—C2—C1	120.8 (2)	C11—C10—C9	118.6 (3)
C3—C2—H2	119.6	C11—C10—H10	120.7
C1—C2—H2	119.6	C9—C10—H10	120.7
C2—C3—N1	120.5 (2)	C10—C11—C12	121.0 (3)
C2—C3—H3	119.7	C10—C11—H11	119.5
N1—C3—H3	119.7	C12—C11—H11	119.5
C3—N1—C4	118.6 (2)	C11—C12—C13	121.0 (3)
C5—C4—N1	122.4 (2)	C11—C12—H12	119.5
C5—C4—H4	118.8	C13—C12—H12	119.5
N1—C4—H4	118.8	C12—C13—C8	119.1 (4)
C4—C5—C1	116.7 (2)	C12—C13—H13	120.4
C4—C5—C6	116.9 (2)	C8—C13—H13	120.4
C1—C5—C6	126.5 (2)		


Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N4—H4A···O1 ⁱ	0.86	2.11	2.903 (3)	153
N3—H3A···O2 ⁱⁱ	0.86	1.98	2.831 (3)	170

Symmetry codes: (i) $-x+2, -y, -z+1$; (ii) $-x+1, -y, -z+1$.

supplementary materials

Fig. 1



Article retracted

Fig. 2

