

Bis[2-(benzyliminomethyl)-4-chlorophenolato- $\kappa^2 N,O$]nickel(II)

Zong-Xiao Li, Xin-Li Zhang and Xiao-Hua Pu*

Department of Chemistry, Baoji University of Arts and Science, Baoji, Shaanxi 721007, People's Republic of China
Correspondence e-mail: mingtian8001@163.com


Received 23 October 2007; accepted 31 October 2007

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.043; wR factor = 0.124; data-to-parameter ratio = 13.2.

In the mononuclear centrosymmetric title compound, $[\text{Ni}(\text{C}_{14}\text{H}_{11}\text{ClNO})_2]$, the Ni^{II} atom, lying on a center of symmetry, is four-coordinated by two O atoms and two N atoms from two Schiff base ligands, forming a slightly distorted square-planar environment. The dihedral angle between the two aromatic rings of the ligand is $72.0(2)^\circ$. No significant hydrogen bonding or $\pi-\pi$ stacking interactions are observed.

Related literature

For bond-length data, see: Allen *et al.* (1987). For related literature, see: Christensen *et al.* (1997); Costes *et al.* (2005); Hu *et al.* (2005); Liu *et al.* (2006); Wallis & Cummings (1974); Yu (2006).



Experimental

Crystal data

$[\text{Ni}(\text{C}_{14}\text{H}_{11}\text{ClNO})_2]$
 $M_r = 548.09$
Monoclinic, $P2_1/c$
 $a = 13.6785(17)\text{ \AA}$
 $b = 10.5986(14)\text{ \AA}$
 $c = 8.6560(13)\text{ \AA}$
 $\beta = 107.529(2)^\circ$
 $V = 1196.6(3)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.06\text{ mm}^{-1}$
 $T = 298(2)\text{ K}$
 $0.56 \times 0.44 \times 0.32\text{ mm}$

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.587$, $T_{\max} = 0.727$
5718 measured reflections
2110 independent reflections
1506 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.124$
 $S = 1.08$
2110 reflections
160 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.53\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Ni1—O1	1.817 (2)	Ni1—N1	1.926 (3)
O1—Ni1—O1 ⁱ	180	O1—Ni1—N1	92.61 (11)
O1—Ni1—N1 ⁱ	87.39 (11)		

Symmetry code: (i) $-x, -y + 1, -z$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Bruker, 2001); software used to prepare material for publication: *SHELXTL*.

Financial support by the Phytochemistry Key Laboratory of Shaanxi province (grant No. 02js40) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2496).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, S1–19.
Bruker (1998). *SMART*. Version 5.0. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2001). *SAINT* (Version 6.02) and *SHELXTL* (Version 6.12). Bruker AXS Inc., Madison, Wisconsin, USA.
Christensen, A., Jensen, H. S., McKee, V., McKenzie, C. J. & Munch, M. (1997). *Inorg. Chem.* **36**, 6080–6085.
Costes, J. P., Lamere, J. F., Lepetit, C., Lacroix, P. G. & Dahan, F. (2005). *Inorg. Chem.* **44**, 1973–1982.
Hu, Z.-Q., Li, W.-H., Ding, Y. & Wu, Y. (2005). *Acta Cryst. E61*, m2526–m2527.
Liu, H.-Y., Gao, F., Lu, Z.-S. & Wang, H.-Y. (2006). *Acta Cryst. E62*, m1306–m1308.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (1997). *SHELXL97* and *SHELXS97*. University of Göttingen, Germany.
Wallis, W. N. & Cummings, S. C. (1974). *Inorg. Chem.* **13**, 991–994.
Yu, Y.-Y. (2006). *Acta Cryst. E62*, m948–m949.

supplementary materials

Acta Cryst. (2008). E64, m202 [doi:10.1107/S1600536807054943]

Bis[2-(benzyliminomethyl)-4-chlorophenolato- κ^2N,O]nickel(II)

Z.-X. Li, X.-L. Zhang and X.-H. Pu

Comment

Recently, we have reported a Schiff base nickel(II) complex (Hu *et al.*, 2005). Owing to the nickel complexes derived from Schiff base ligands possess interesting structures and wide applications (Costes *et al.*, 2005; Wallis & Cummings, 1974; Christensen *et al.*, 1997; Liu *et al.*, 2006); Yu, 2006), we report here the crystal structure of a new Schiff base nickel(II) complex, title compound, (I),

Compound (I) is a mononuclear centrosymmetric Ni^{II} complex (Fig. 1) The Ni atom, lying on the center of symmetry, is four-coordinated by two O atoms and two N atoms from two Schiff base ligands, forming a slightly distorted square-planar environment (Table 1). The bond lengths and angles of the ligands show normal values (Allen *et al.*, 1987). The dihedral angle between the two aromatic rings of the ligand is 72.0 (2)^o. No significant hydrogen bonding or π - π stacking interactions are observed.

Experimental

5-Chlorosalicylaldehyde (0.1 mmol, 15.7 mg), Ni(NO₃)₂.6H₂O (0.1 mmol, 29.0 mg) and benzylamine (0.1 mmol, 10.7 mg) were dissolved in methanol (10 ml). The mixture was stirred for 30 min at room temperature to give a clear brown solution. After allowing the resulting solution to stand in air for 11 d, brown block-shaped crystals of compound (I) were formed on slow evaporation of the solvent. The crystals were collected, washed with methanol and dried in a vacuum desiccator using anhydrous CaCl₂ (yield 54%). Analysis found: C 61.30, H 4.0%; calculated for Ni(C₁₄H₁₁ClO)₂: C 61.34, H 4.01%.

Refinement

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures



Fig. 1. The molecular structure of (I), showing 30% probability displacement ellipsoids. H atoms have been omitted for clarity. Unlabelled atoms are related to other labelled atoms by the symmetry operation (-x, 1 - y, -z).




Fig. 2. The crystal packing of (I), viewed along the c axis.

supplementary materials

Bis[2-(benzyliminomethyl)-4-chlorophenolato- κ^2N,O]nickel(II)

Crystal data

[Ni(C ₁₄ H ₁₁ ClNO) ₂]	$F_{000} = 564$
$M_r = 548.09$	$D_x = 1.521 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 13.6785 (17) \text{ \AA}$	Cell parameters from 1825 reflections
$b = 10.5986 (14) \text{ \AA}$	$\theta = 2.5\text{--}25.2^\circ$
$c = 8.6560 (13) \text{ \AA}$	$\mu = 1.06 \text{ mm}^{-1}$
$\beta = 107.529 (2)^\circ$	$T = 298 (2) \text{ K}$
$V = 1196.6 (3) \text{ \AA}^3$	Rhombus, green
$Z = 2$	$0.56 \times 0.44 \times 0.32 \text{ mm}$

Data collection

Bruker SMART CCD diffractometer	2110 independent reflections
Radiation source: fine-focus sealed tube	1506 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.039$
$T = 298(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -16 \rightarrow 12$
$T_{\text{min}} = 0.587$, $T_{\text{max}} = 0.727$	$k = -12 \rightarrow 8$
5718 measured reflections	$l = -10 \rightarrow 10$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.043$	H-atom parameters constrained
$wR(F^2) = 0.124$	$w = 1/[\sigma^2(F_o^2) + (0.057P)^2 + 0.6708P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.08$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2110 reflections	$\Delta\rho_{\text{max}} = 0.53 \text{ e \AA}^{-3}$
160 parameters	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.0000	0.5000	0.0000	0.0329 (2)
Cl1	0.48272 (9)	0.77502 (16)	0.44350 (18)	0.0919 (6)
N1	0.0104 (2)	0.6504 (3)	0.1288 (3)	0.0316 (7)
O1	0.13607 (19)	0.4664 (3)	0.0829 (3)	0.0472 (7)
C1	0.0949 (3)	0.7006 (4)	0.2141 (4)	0.0354 (8)
H1	0.0899	0.7737	0.2705	0.043*
C2	0.1963 (3)	0.6549 (4)	0.2309 (4)	0.0359 (9)
C3	0.2107 (3)	0.5393 (4)	0.1629 (4)	0.0365 (9)
C4	0.3123 (3)	0.4982 (4)	0.1878 (5)	0.0467 (10)
H4	0.3240	0.4208	0.1461	0.056*
C5	0.3939 (3)	0.5699 (5)	0.2720 (5)	0.0528 (11)
H5	0.4603	0.5418	0.2855	0.063*
C6	0.3779 (3)	0.6840 (5)	0.3372 (5)	0.0520 (11)
C7	0.2809 (3)	0.7283 (4)	0.3197 (5)	0.0451 (10)
H7	0.2710	0.8048	0.3653	0.054*
C8	-0.0831 (3)	0.7179 (4)	0.1351 (4)	0.0369 (9)
H8A	-0.1254	0.7353	0.0253	0.044*
H8B	-0.0633	0.7982	0.1893	0.044*
C9	-0.1461 (2)	0.6457 (3)	0.2219 (4)	0.0316 (8)
C10	-0.1084 (3)	0.5490 (4)	0.3262 (4)	0.0393 (9)
H10	-0.0411	0.5225	0.3438	0.047*
C11	-0.1688 (3)	0.4897 (4)	0.4064 (5)	0.0469 (10)
H11	-0.1422	0.4233	0.4767	0.056*
C12	-0.2678 (3)	0.5281 (4)	0.3827 (5)	0.0533 (12)
H12	-0.3087	0.4878	0.4361	0.064*
C13	-0.3059 (3)	0.6264 (5)	0.2799 (5)	0.0553 (12)
H13	-0.3725	0.6544	0.2651	0.066*
C14	-0.2457 (3)	0.6839 (4)	0.1983 (5)	0.0458 (10)
H14	-0.2727	0.7493	0.1265	0.055*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0295 (4)	0.0348 (4)	0.0367 (4)	-0.0008 (3)	0.0133 (3)	-0.0002 (3)
Cl1	0.0415 (7)	0.1240 (14)	0.1083 (11)	-0.0314 (7)	0.0197 (7)	-0.0503 (10)
N1	0.0291 (15)	0.0328 (17)	0.0363 (16)	0.0014 (13)	0.0152 (13)	0.0075 (13)
O1	0.0307 (14)	0.0475 (19)	0.0590 (17)	0.0032 (11)	0.0070 (13)	-0.0124 (13)
C1	0.040 (2)	0.031 (2)	0.040 (2)	-0.0020 (16)	0.0191 (17)	0.0023 (16)
C2	0.0311 (19)	0.043 (2)	0.035 (2)	-0.0034 (17)	0.0123 (15)	0.0036 (18)
C3	0.033 (2)	0.045 (2)	0.035 (2)	-0.0018 (16)	0.0137 (17)	0.0024 (17)
C4	0.035 (2)	0.053 (3)	0.054 (2)	0.0008 (19)	0.0156 (18)	-0.010 (2)
C5	0.031 (2)	0.069 (3)	0.059 (3)	-0.001 (2)	0.0156 (19)	-0.005 (2)
C6	0.034 (2)	0.073 (3)	0.050 (2)	-0.014 (2)	0.0137 (18)	-0.008 (2)
C7	0.037 (2)	0.049 (3)	0.051 (2)	-0.0103 (18)	0.0162 (18)	-0.008 (2)
C8	0.038 (2)	0.032 (2)	0.043 (2)	0.0018 (16)	0.0163 (17)	0.0026 (17)
C9	0.0308 (18)	0.033 (2)	0.0309 (19)	-0.0009 (15)	0.0095 (15)	-0.0046 (16)
C10	0.039 (2)	0.037 (2)	0.043 (2)	0.0046 (17)	0.0142 (17)	0.0024 (18)
C11	0.055 (3)	0.043 (2)	0.047 (2)	0.003 (2)	0.0222 (19)	0.009 (2)
C12	0.049 (2)	0.067 (3)	0.053 (2)	-0.012 (2)	0.029 (2)	-0.001 (2)
C13	0.035 (2)	0.079 (4)	0.055 (3)	0.005 (2)	0.019 (2)	0.002 (2)
C14	0.040 (2)	0.055 (3)	0.043 (2)	0.0121 (19)	0.0139 (18)	0.010 (2)

Geometric parameters (\AA , $^\circ$)

Ni1—O1	1.817 (2)	C6—C7	1.372 (5)
Ni1—O1 ⁱ	1.817 (2)	C7—H7	0.93
Ni1—N1 ⁱ	1.926 (3)	C8—C9	1.510 (5)
Ni1—N1	1.926 (3)	C8—H8A	0.97
Cl1—C6	1.743 (4)	C8—H8B	0.97
N1—C1	1.284 (4)	C9—C10	1.361 (5)
N1—C8	1.480 (4)	C9—C14	1.377 (5)
O1—C3	1.301 (4)	C10—C11	1.380 (5)
C1—C2	1.434 (5)	C10—H10	0.93
C1—H1	0.93	C11—C12	1.369 (6)
C2—C3	1.399 (5)	C11—H11	0.93
C2—C7	1.414 (5)	C12—C13	1.367 (6)
C3—C4	1.409 (5)	C12—H12	0.93
C4—C5	1.365 (6)	C13—C14	1.378 (6)
C4—H4	0.93	C13—H13	0.93
C5—C6	1.379 (6)	C14—H14	0.93
C5—H5	0.93		
O1—Ni1—O1 ⁱ	180	C6—C7—C2	118.6 (4)
O1—Ni1—N1 ⁱ	87.39 (11)	C6—C7—H7	120.7
O1 ⁱ —Ni1—N1 ⁱ	92.61 (11)	C2—C7—H7	120.7
O1—Ni1—N1	92.61 (11)	N1—C8—C9	113.7 (3)
O1 ⁱ —Ni1—N1	87.39 (11)	N1—C8—H8A	108.8

N1 ⁱ —Ni1—N1	180.00 (14)	C9—C8—H8A	108.8
C1—N1—C8	114.6 (3)	N1—C8—H8B	108.8
C1—N1—Ni1	124.9 (2)	C9—C8—H8B	108.8
C8—N1—Ni1	120.5 (2)	H8A—C8—H8B	107.7
C3—O1—Ni1	129.8 (3)	C10—C9—C14	118.7 (3)
N1—C1—C2	126.4 (4)	C10—C9—C8	123.5 (3)
N1—C1—H1	116.8	C14—C9—C8	117.7 (3)
C2—C1—H1	116.8	C9—C10—C11	120.8 (3)
C3—C2—C7	120.9 (3)	C9—C10—H10	119.6
C3—C2—C1	120.5 (3)	C11—C10—H10	119.6
C7—C2—C1	118.5 (4)	C12—C11—C10	120.3 (4)
O1—C3—C2	123.8 (3)	C12—C11—H11	119.9
O1—C3—C4	118.5 (4)	C10—C11—H11	119.9
C2—C3—C4	117.6 (3)	C13—C12—C11	119.4 (4)
C5—C4—C3	121.3 (4)	C13—C12—H12	120.3
C5—C4—H4	119.3	C11—C12—H12	120.3
C3—C4—H4	119.3	C12—C13—C14	120.1 (4)
C4—C5—C6	120.1 (4)	C12—C13—H13	120.0
C4—C5—H5	120.0	C14—C13—H13	120.0
C6—C5—H5	120.0	C9—C14—C13	120.7 (4)
C7—C6—C5	121.4 (4)	C9—C14—H14	119.6
C7—C6—Cl1	118.9 (4)	C13—C14—H14	119.6
C5—C6—Cl1	119.6 (3)		
O1—Ni1—N1—C1	9.0 (3)	C4—C5—C6—C7	0.0 (7)
O1 ⁱ —Ni1—N1—C1	-171.0 (3)	C4—C5—C6—Cl1	-179.7 (3)
O1—Ni1—N1—C8	-171.1 (2)	C5—C6—C7—C2	-0.8 (6)
O1 ⁱ —Ni1—N1—C8	8.9 (2)	Cl1—C6—C7—C2	178.9 (3)
N1 ⁱ —Ni1—O1—C3	164.0 (3)	C3—C2—C7—C6	0.4 (5)
N1—Ni1—O1—C3	-16.0 (3)	C1—C2—C7—C6	179.3 (3)
C8—N1—C1—C2	179.6 (3)	C1—N1—C8—C9	-111.5 (3)
Ni1—N1—C1—C2	-0.4 (5)	Ni1—N1—C8—C9	68.5 (3)
N1—C1—C2—C3	-6.1 (6)	N1—C8—C9—C10	19.2 (5)
N1—C1—C2—C7	174.9 (3)	N1—C8—C9—C14	-163.5 (3)
Ni1—O1—C3—C2	14.1 (5)	C14—C9—C10—C11	0.4 (6)
Ni1—O1—C3—C4	-168.2 (3)	C8—C9—C10—C11	177.6 (4)
C7—C2—C3—O1	178.5 (3)	C9—C10—C11—C12	-0.5 (6)
C1—C2—C3—O1	-0.5 (5)	C10—C11—C12—C13	-0.5 (6)
C7—C2—C3—C4	0.7 (5)	C11—C12—C13—C14	1.5 (7)
C1—C2—C3—C4	-178.2 (3)	C10—C9—C14—C13	0.7 (6)
O1—C3—C4—C5	-179.4 (4)	C8—C9—C14—C13	-176.8 (4)
C2—C3—C4—C5	-1.5 (6)	C12—C13—C14—C9	-1.6 (7)
C3—C4—C5—C6	1.1 (6)		

Symmetry codes: (i) $-x, -y+1, -z$.

supplementary materials

Fig. 1




Fig. 2

