

## Cinchonidinium chloride monohydrate

Shi-Feng Ni,<sup>a</sup> Lin Lin Ma,<sup>b</sup> Gui Fang Zhao,<sup>a</sup> Wen Ji Sun<sup>a</sup> and Zhi Min Jin<sup>b\*</sup>

<sup>a</sup>Key Laboratory of Resource Biology and Biotechnology in Western China (Northwest University), Ministry of Education, College of Life Sciences, Northwest University, Xi'an 710069, People's Republic of China, and <sup>b</sup>College of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014, People's Republic of China  
Correspondence e-mail: apharm@sina.com


Received 23 November 2007; accepted 26 November 2007

Key indicators: single-crystal X-ray study;  $T = 273\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$ ; disorder in main residue;  $R$  factor = 0.069;  $wR$  factor = 0.179; data-to-parameter ratio = 12.8.

In the title salt,  $\text{C}_{19}\text{H}_{23}\text{N}_2\text{O}^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$ , the ions and the water molecule are held together by  $\text{O}-\text{H}\cdots\text{Cl}$ ,  $\text{N}-\text{H}\cdots\text{Cl}$ ,  $\text{O}-\text{H}\cdots\text{O}$ ,  $\text{O}-\text{H}\cdots\text{N}$  and  $\text{C}-\text{H}\cdots\text{Cl}$  hydrogen bonds, forming a three-dimensional framework. The vinyl group is disordered over two orientations with refined occupancies of 0.564 (16) and 0.436 (16). The cell parameters of the title compound have been reported previously [Griffiths (1952). *Acta Cryst.* **5**, 290–291].

### Related literature

For the Cambridge Structural Database (Version 5.26), see: Allen (2002). For related literature, see: Griffiths (1952); Zhang, Lü *et al.* (2006); Zhang, Tu *et al.* (2006).



### Experimental

#### Crystal data

$\text{C}_{19}\text{H}_{23}\text{N}_2\text{O}^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$   
 $M_r = 348.86$   
Orthorhombic,  $P2_12_12_1$

$a = 10.4924(8)\text{ \AA}$   
 $b = 12.9049(10)\text{ \AA}$   
 $c = 13.3936(10)\text{ \AA}$

$V = 1813.5(2)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.22\text{ mm}^{-1}$   
 $T = 273(2)\text{ K}$   
 $0.37 \times 0.34 \times 0.12\text{ mm}$

#### Data collection

Siemens P4 diffractometer  
Absorption correction: multi-scan (*SHELXTL*; Bruker, 1998)  
 $T_{\min} = 0.722$ ,  $T_{\max} = 0.895$   
(expected range = 0.785–0.973)  
9572 measured reflections

3214 independent reflections  
3090 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$   
3 standard reflections  
every 97 reflections  
intensity decay: 5.0%

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.069$   
 $wR(F^2) = 0.179$   
 $S = 1.18$   
3214 reflections  
252 parameters  
12 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.31\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$   
Absolute structure: Flack (1983),  
1369 Friedel pairs  
Flack parameter: 0.04 (13)

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 $\cdots$ O2 <sup>i</sup>	0.82 (2)	1.84 (2)	2.653 (5)	172 (3)
O2—H2OB $\cdots$ N1 <sup>ii</sup>	0.82 (2)	2.07 (2)	2.867 (6)	165 (3)
O2—H2OA $\cdots$ Cl1	0.82 (2)	2.31 (2)	3.135 (5)	174 (6)
N2—H2 $\cdots$ Cl1	0.96 (4)	2.08 (4)	3.034 (6)	175 (3)
C5—H5 $\cdots$ Cl1	0.93	2.77	3.692 (5)	172

Symmetry codes: (i)  $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + \frac{3}{2}, -y + 1, z + \frac{1}{2}$ .

Data collection: *XSCANS* (Siemens, 1994); cell refinement: *XSCANS*; data reduction: *SHELXTL* (Bruker, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2530).

### References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Bruker (1998). *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Griffiths, P. J. F. (1952). *Acta Cryst.* **5**, 290–291.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Siemens (1994). *XSCANS*. Version 2.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Zhang, H.-M., Lü, Y.-P., Tu, B., Jin, Z.-M. & Wu, Z.-Y. (2006). *Acta Cryst.* **E62**, o3398–o3400.
- Zhang, H.-M., Tu, B., Jin, Z.-M., Wu, Z.-Y. & Hu, M.-L. (2006). *Acta Cryst.* **E62**, o2614–o2616.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o193 [doi:10.1107/S160053680706343X]

### Cinchonidinium chloride monohydrate

**S.-F. Ni, L. L. Ma, G. F. Zhao, W. J. Sun and Z. M. Jin**

#### Comment

The cell parameters ( $a = 12.8$ ,  $b = 13.3$ ,  $c = 10.6 \text{ \AA}$ ) of the title compound have been reported previously (Griffiths, 1952; CSD refcode ZZTZW, CSD, Version 5.26; Allen, 2002), but structural details are not available. Crystal structures of some compounds containing cinchonidine have also been reported, *e.g.* cinchonidinium bis(4-methylbenzenesulfonate) monohydrate (Zhang, Lü *et al.*, 2006), cinchonidinium bis(perchlorate) (Zhang, Tu *et al.*, 2006).

The geometry of the cinchonidinium unit is consistent with that observed in other cinchonidinium compounds (Zhang, Lv *et al.*, 2006; Zhang, Tu *et al.*, 2006), except for the disorder in the vinyl group. The C—C bond lengths in quinoline ring system range from 1.358 (6) to 1.429 (6)  $\text{\AA}$ , comparable with the range of 1.344 (5)–1.422 (4)  $\text{\AA}$  observed by Zhang, Lv *et al.* (2006).

As shown in Fig. 1, the chloride anion and cinchonidinium cation are linked by N—H $\cdots$ Cl and weak C—H $\cdots$ Cl hydrogen bonds, and the chloride ion and water molecule are connected by a O—H $\cdots$ Cl hydrogen bond (Table 1). The ionic pairs and water molecule are linked by O—H $\cdots$ N and O—H $\cdots$ O hydrogen bonds to form a three-dimensional framework (Fig. 2).

#### Experimental

Cinchonidine (0.01 mol, 2.94 g) and 10% hydrochloric acid (3.65 g) were mixed together with enough water, and heated to a temperature where a clear solution was resulted. Colourless single crystals of the title compound were obtained by slow evaporation of the solution at room temperature for 7 d.

#### Refinement

The vinyl group is disordered over two orientations with refined occupancies of 0.564 (16) and 0.436 (16); the corresponding C—C distances in the major and minor conformers were restrained to be equal. Atoms C18' and C19' were restrained to have the same  $U_{ij}^{ij}$  components. N- and O-bound H atoms were located in a difference map and refined with the O—H and H $\cdots$ H (in water) distances restrained to 0.82 (2) and 1.39 (2)  $\text{\AA}$ , respectively. C-bound H atoms were placed in calculated positions (C—H = 0.93–0.98  $\text{\AA}$ ) and allowed to ride on their parent atoms, with  $U_{\text{iso}}(\text{H})$  values set at 1.2 times  $U_{\text{eq}}$  of the parent atoms.

# supplementary materials

---

## Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 40% probability level. Hydrogen bonds are shown as dashed lines.




Fig. 2. The molecular packing of the title compound, viewed along the *b* axis. Hydrogen bonds are shown as dashed lines.

## Cinchonidinium chloride monohydrate

### Crystal data

$\text{C}_{19}\text{H}_{23}\text{N}_2\text{O}^+\cdot\text{Cl}^-\cdot\text{H}_2\text{O}$	$F_{000} = 744$
$M_r = 348.86$	$D_x = 1.278 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 10.4924 (8) \text{ \AA}$	Cell parameters from 31 reflections
$b = 12.9049 (10) \text{ \AA}$	$\theta = 3.3\text{--}18.6^\circ$
$c = 13.3936 (10) \text{ \AA}$	$\mu = 0.22 \text{ mm}^{-1}$
$V = 1813.5 (2) \text{ \AA}^3$	$T = 273 (2) \text{ K}$
$Z = 4$	Plate, colourless
	$0.37 \times 0.34 \times 0.12 \text{ mm}$

### Data collection

Siemens P4 diffractometer	$R_{\text{int}} = 0.031$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.0^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.2^\circ$
$T = 273(2) \text{ K}$	$h = -12 \rightarrow 11$
$\omega$ scans	$k = -9 \rightarrow 15$
Absorption correction: multi-scan (SHELXTL; Bruker, 1998)	$l = -15 \rightarrow 15$
$T_{\text{min}} = 0.722$ , $T_{\text{max}} = 0.895$	3 standard reflections
9572 measured reflections	every 97 reflections
3214 independent reflections	intensity decay: 5.0%

3090 reflections with  $I > 2\sigma(I)$

### Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.069$	$w = 1/[\sigma^2(F_o^2) + (0.0981P)^2 + 0.5633P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.179$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.18$	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
3214 reflections	$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$
252 parameters	Extinction correction: none
12 restraints	Absolute structure: Flack (1983), with 1369 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.04 (13)
Secondary atom site location: difference Fourier map	

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cl1	0.94217 (9)	0.75097 (9)	0.35617 (7)	0.0500 (3)	
O1	0.7326 (3)	0.9410 (2)	0.1041 (2)	0.0511 (8)	
O2	1.0407 (3)	0.5216 (3)	0.3556 (4)	0.0858 (13)	
N1	0.6545 (4)	0.6294 (3)	-0.1087 (2)	0.0510 (9)	
N2	0.6840 (3)	0.8504 (3)	0.3211 (2)	0.0454 (8)	
C1	0.6176 (4)	0.7257 (4)	-0.1063 (3)	0.0533 (12)	
H1A	0.5696	0.7497	-0.1599	0.064*	
C2	0.6443 (4)	0.7957 (4)	-0.0297 (3)	0.0478 (10)	
H2A	0.6143	0.8634	-0.0333	0.057*	
C3	0.7148 (3)	0.7642 (3)	0.0508 (3)	0.0381 (9)	
C4	0.7589 (3)	0.6594 (3)	0.0517 (3)	0.0346 (8)	
C5	0.8343 (4)	0.6152 (3)	0.1287 (3)	0.0422 (9)	
H5	0.8546	0.6549	0.1844	0.051*	

## supplementary materials

---

C6	0.8775 (4)	0.5162 (3)	0.1227 (3)	0.0482 (10)	
H6	0.9271	0.4890	0.1739	0.058*	
C7	0.8475 (4)	0.4552 (4)	0.0401 (3)	0.0527 (11)	
H7	0.8781	0.3877	0.0361	0.063*	
C8	0.7738 (4)	0.4939 (4)	-0.0349 (3)	0.0489 (10)	
H8	0.7533	0.4522	-0.0892	0.059*	
C9	0.7288 (4)	0.5953 (3)	-0.0310 (3)	0.0397 (9)	
C10	0.7394 (3)	0.8374 (3)	0.1368 (3)	0.0363 (8)	
H10	0.8246	0.8241	0.1638	0.044*	
C11	0.6392 (4)	0.8168 (3)	0.2187 (3)	0.0410 (9)	
H11	0.6269	0.7416	0.2216	0.049*	
C12	0.5101 (4)	0.8640 (5)	0.2001 (4)	0.0668 (15)	
H12A	0.4511	0.8109	0.1780	0.080*	
H12B	0.5163	0.9162	0.1482	0.080*	
C13	0.4614 (5)	0.9137 (5)	0.2972 (4)	0.0701 (15)	
H13	0.3743	0.9388	0.2869	0.084*	
C14	0.4621 (5)	0.8368 (4)	0.3821 (3)	0.0615 (13)	
H14	0.4334	0.8725	0.4426	0.074*	
C15	0.5996 (5)	0.8022 (4)	0.3980 (3)	0.0620 (13)	
H15A	0.6277	0.8227	0.4642	0.074*	
H15B	0.6048	0.7273	0.3934	0.074*	
C16	0.5460 (6)	1.0037 (4)	0.3240 (5)	0.0831 (18)	
H16A	0.5175	1.0346	0.3861	0.100*	
H16B	0.5420	1.0560	0.2721	0.100*	
C17	0.6826 (5)	0.9649 (4)	0.3353 (4)	0.0613 (13)	
H17A	0.7369	0.9978	0.2860	0.074*	
H17B	0.7147	0.9822	0.4011	0.074*	
C18	0.3877 (9)	0.7331 (8)	0.3733 (7)	0.050 (3)	0.564 (16)
H18	0.4218	0.6754	0.4051	0.061*	0.564 (16)
C19	0.2832 (9)	0.7210 (9)	0.3257 (8)	0.067 (3)	0.564 (16)
H19A	0.2460	0.7768	0.2929	0.081*	0.564 (16)
H19B	0.2443	0.6562	0.3239	0.081*	0.564 (16)
C18'	0.3393 (10)	0.7759 (9)	0.3550 (12)	0.080 (6)	0.436 (16)
H18'	0.2713	0.8063	0.3215	0.096*	0.436 (16)
C19'	0.338 (2)	0.6812 (10)	0.3823 (11)	0.079 (5)	0.436 (16)
H19C	0.4083	0.6535	0.4158	0.094*	0.436 (16)
H19D	0.2680	0.6398	0.3687	0.094*	0.436 (16)
H2	0.765 (4)	0.820 (3)	0.336 (3)	0.035 (10)*	
H1	0.803 (3)	0.967 (4)	0.111 (4)	0.062 (16)*	
H2OA	1.020 (4)	0.5832 (15)	0.358 (5)	0.081 (18)*	
H2OB	0.982 (3)	0.480 (2)	0.354 (4)	0.061 (15)*	

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0518 (5)	0.0549 (6)	0.0432 (5)	0.0083 (5)	-0.0100 (4)	-0.0045 (5)
O1	0.0510 (18)	0.0404 (16)	0.0618 (18)	-0.0074 (14)	-0.0066 (15)	0.0044 (14)
O2	0.0468 (19)	0.051 (2)	0.160 (4)	0.0026 (17)	0.002 (3)	0.012 (3)

N1	0.053 (2)	0.059 (2)	0.0405 (18)	-0.0011 (18)	-0.0096 (16)	-0.0068 (17)
N2	0.0440 (19)	0.053 (2)	0.0387 (17)	0.0083 (17)	-0.0003 (14)	-0.0012 (16)
C1	0.056 (2)	0.066 (3)	0.038 (2)	0.000 (2)	-0.0127 (19)	0.006 (2)
C2	0.051 (2)	0.044 (2)	0.048 (2)	0.002 (2)	-0.005 (2)	0.0043 (18)
C3	0.0346 (18)	0.047 (2)	0.0329 (18)	-0.0061 (18)	0.0032 (14)	0.0078 (17)
C4	0.0298 (17)	0.045 (2)	0.0294 (18)	-0.0013 (16)	0.0031 (14)	0.0000 (16)
C5	0.047 (2)	0.045 (2)	0.0344 (19)	-0.0088 (18)	-0.0041 (17)	-0.0029 (17)
C6	0.051 (2)	0.045 (2)	0.048 (2)	0.0081 (19)	-0.008 (2)	0.0003 (19)
C7	0.057 (3)	0.042 (2)	0.059 (3)	0.011 (2)	0.004 (2)	-0.001 (2)
C8	0.056 (2)	0.050 (2)	0.041 (2)	-0.005 (2)	-0.0007 (19)	-0.0162 (19)
C9	0.0357 (19)	0.049 (2)	0.0347 (18)	-0.0030 (18)	0.0058 (16)	-0.0030 (17)
C10	0.0369 (18)	0.0347 (18)	0.0374 (18)	-0.0003 (16)	-0.0076 (16)	0.0005 (16)
C11	0.042 (2)	0.043 (2)	0.038 (2)	-0.0058 (18)	-0.0023 (16)	0.0026 (17)
C12	0.040 (2)	0.113 (4)	0.047 (3)	0.007 (3)	0.0029 (19)	0.009 (3)
C13	0.045 (3)	0.092 (4)	0.073 (3)	0.019 (3)	0.008 (2)	0.005 (3)
C14	0.061 (3)	0.076 (3)	0.048 (3)	-0.011 (3)	0.021 (2)	-0.015 (2)
C15	0.090 (4)	0.060 (3)	0.036 (2)	0.002 (3)	0.011 (2)	0.003 (2)
C16	0.086 (4)	0.053 (3)	0.110 (5)	0.019 (3)	0.031 (4)	0.006 (3)
C17	0.073 (3)	0.056 (3)	0.055 (3)	-0.011 (2)	0.011 (2)	-0.013 (2)
C18	0.068 (6)	0.026 (6)	0.058 (5)	-0.010 (5)	0.005 (4)	0.008 (4)
C19	0.078 (7)	0.051 (7)	0.072 (7)	-0.015 (5)	0.003 (5)	-0.019 (5)
C18'	0.108 (15)	0.040 (8)	0.092 (11)	0.033 (9)	0.034 (11)	-0.006 (9)
C19'	0.115 (13)	0.046 (8)	0.075 (10)	-0.012 (9)	-0.007 (9)	-0.010 (7)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C10	1.409 (5)	C11—C12	1.505 (6)
O1—H1	0.82 (2)	C11—H11	0.98
O2—H2OA	0.823 (19)	C12—C13	1.538 (7)
O2—H2OB	0.816 (18)	C12—H12A	0.97
N1—C1	1.302 (6)	C12—H12B	0.97
N1—C9	1.372 (5)	C13—C16	1.506 (9)
N2—C17	1.490 (6)	C13—C14	1.508 (7)
N2—C15	1.495 (6)	C13—H13	0.98
N2—C11	1.512 (5)	C14—C15	1.524 (7)
N2—H2	0.96 (4)	C14—C18'	1.552 (12)
C1—C2	1.395 (6)	C14—C18	1.554 (9)
C1—H1A	0.93	C14—H14	0.98
C2—C3	1.370 (6)	C15—H15A	0.97
C2—H2A	0.93	C15—H15B	0.97
C3—C4	1.429 (6)	C16—C17	1.525 (8)
C3—C10	1.512 (5)	C16—H16A	0.97
C4—C9	1.419 (5)	C16—H16B	0.97
C4—C5	1.419 (5)	C17—H17A	0.97
C5—C6	1.358 (6)	C17—H17B	0.97
C5—H5	0.93	C18—C19	1.278 (11)
C6—C7	1.394 (6)	C18—H18	0.93
C6—H6	0.93	C19—H19A	0.93
C7—C8	1.363 (6)	C19—H19B	0.93

## supplementary materials

---

C7—H7	0.93	C18'—C19'	1.277 (12)
C8—C9	1.391 (6)	C18'—H18'	0.93
C8—H8	0.93	C19'—H19C	0.93
C10—C11	1.543 (5)	C19'—H19D	0.93
C10—H10	0.98		
C10—O1—H1	108 (4)	C13—C12—H12A	109.9
H2OA—O2—H2OB	116 (3)	C11—C12—H12B	109.9
C1—N1—C9	117.1 (3)	C13—C12—H12B	109.9
C17—N2—C15	108.6 (4)	H12A—C12—H12B	108.3
C17—N2—C11	113.4 (3)	C16—C13—C14	109.0 (5)
C15—N2—C11	108.8 (3)	C16—C13—C12	109.1 (5)
C17—N2—H2	113 (2)	C14—C13—C12	111.2 (4)
C15—N2—H2	102 (2)	C16—C13—H13	109.2
C11—N2—H2	110 (2)	C14—C13—H13	109.2
N1—C1—C2	125.2 (4)	C12—C13—H13	109.2
N1—C1—H1A	117.4	C13—C14—C15	107.6 (4)
C2—C1—H1A	117.4	C13—C14—C18'	98.8 (7)
C3—C2—C1	119.6 (4)	C15—C14—C18'	132.1 (6)
C3—C2—H2A	120.2	C13—C14—C18	120.4 (5)
C1—C2—H2A	120.2	C15—C14—C18	103.5 (5)
C2—C3—C4	117.6 (4)	C13—C14—H14	108.2
C2—C3—C10	120.4 (4)	C15—C14—H14	108.2
C4—C3—C10	122.0 (3)	C18'—C14—H14	100.1
C9—C4—C5	117.2 (4)	C18—C14—H14	108.2
C9—C4—C3	118.2 (3)	N2—C15—C14	110.0 (4)
C5—C4—C3	124.5 (3)	N2—C15—H15A	109.7
C6—C5—C4	121.4 (4)	C14—C15—H15A	109.7
C6—C5—H5	119.3	N2—C15—H15B	109.7
C4—C5—H5	119.3	C14—C15—H15B	109.7
C5—C6—C7	120.2 (4)	H15A—C15—H15B	108.2
C5—C6—H6	119.9	C13—C16—C17	108.9 (4)
C7—C6—H6	119.9	C13—C16—H16A	109.9
C8—C7—C6	120.4 (4)	C17—C16—H16A	109.9
C8—C7—H7	119.8	C13—C16—H16B	109.9
C6—C7—H7	119.8	C17—C16—H16B	109.9
C7—C8—C9	120.6 (4)	H16A—C16—H16B	108.3
C7—C8—H8	119.7	N2—C17—C16	108.8 (4)
C9—C8—H8	119.7	N2—C17—H17A	109.9
N1—C9—C8	117.8 (4)	C16—C17—H17A	109.9
N1—C9—C4	122.1 (4)	N2—C17—H17B	109.9
C8—C9—C4	120.1 (4)	C16—C17—H17B	109.9
O1—C10—C3	110.4 (3)	H17A—C17—H17B	108.3
O1—C10—C11	110.5 (3)	C19—C18—C14	125.1 (11)
C3—C10—C11	108.5 (3)	C19—C18—H18	117.4
O1—C10—H10	109.1	C14—C18—H18	117.4
C3—C10—H10	109.1	C18—C19—H19A	120.0
C11—C10—H10	109.1	C18—C19—H19B	120.0
C12—C11—N2	108.3 (4)	H19A—C19—H19B	120.0
C12—C11—C10	115.2 (3)	C19'—C18'—C14	115.1 (13)

N2—C11—C10	112.6 (3)	C19'—C18'—H18'	122.4
C12—C11—H11	106.8	C14—C18'—H18'	122.4
N2—C11—H11	106.8	C18'—C19'—H19C	120.0
C10—C11—H11	106.8	C18'—C19'—H19D	120.0
C11—C12—C13	109.1 (4)	H19C—C19'—H19D	120.0
C11—C12—H12A	109.9		
C9—N1—C1—C2	−2.0 (7)	C3—C10—C11—C12	−79.1 (5)
N1—C1—C2—C3	0.1 (7)	O1—C10—C11—N2	−82.8 (4)
C1—C2—C3—C4	0.4 (6)	C3—C10—C11—N2	156.1 (3)
C1—C2—C3—C10	−177.1 (4)	N2—C11—C12—C13	−9.2 (5)
C2—C3—C4—C9	0.9 (5)	C10—C11—C12—C13	−136.2 (4)
C10—C3—C4—C9	178.5 (3)	C11—C12—C13—C16	65.9 (6)
C2—C3—C4—C5	179.0 (4)	C11—C12—C13—C14	−54.3 (6)
C10—C3—C4—C5	−3.4 (5)	C16—C13—C14—C15	−59.0 (5)
C9—C4—C5—C6	0.9 (6)	C12—C13—C14—C15	61.3 (5)
C3—C4—C5—C6	−177.2 (4)	C16—C13—C14—C18'	161.5 (6)
C4—C5—C6—C7	−0.3 (6)	C12—C13—C14—C18'	−78.3 (6)
C5—C6—C7—C8	−0.6 (7)	C16—C13—C14—C18	−177.2 (6)
C6—C7—C8—C9	1.0 (7)	C12—C13—C14—C18	−56.9 (7)
C1—N1—C9—C8	−177.1 (4)	C17—N2—C15—C14	64.0 (5)
C1—N1—C9—C4	3.5 (6)	C11—N2—C15—C14	−59.8 (5)
C7—C8—C9—N1	−179.7 (4)	C13—C14—C15—N2	−3.6 (5)
C7—C8—C9—C4	−0.3 (6)	C18'—C14—C15—N2	116.6 (9)
C5—C4—C9—N1	178.8 (4)	C18—C14—C15—N2	125.0 (5)
C3—C4—C9—N1	−3.0 (5)	C14—C13—C16—C17	63.4 (6)
C5—C4—C9—C8	−0.6 (5)	C12—C13—C16—C17	−58.2 (6)
C3—C4—C9—C8	177.6 (4)	C15—N2—C17—C16	−59.6 (5)
C2—C3—C10—O1	−25.4 (5)	C11—N2—C17—C16	61.4 (5)
C4—C3—C10—O1	157.1 (3)	C13—C16—C17—N2	−2.9 (6)
C2—C3—C10—C11	95.8 (4)	C13—C14—C18—C19	−33.2 (12)
C4—C3—C10—C11	−81.6 (4)	C15—C14—C18—C19	−153.4 (10)
C17—N2—C11—C12	−53.9 (5)	C18'—C14—C18—C19	13.9 (13)
C15—N2—C11—C12	67.0 (5)	C13—C14—C18'—C19'	150.7 (12)
C17—N2—C11—C10	74.6 (4)	C15—C14—C18'—C19'	27.2 (17)
C15—N2—C11—C10	−164.5 (3)	C18—C14—C18'—C19'	10.5 (12)
O1—C10—C11—C12	42.0 (5)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1···O2 <sup>i</sup>	0.82 (2)	1.84 (2)	2.653 (5)	172 (3)
O2—H2OB···N1 <sup>ii</sup>	0.82 (2)	2.07 (2)	2.867 (6)	165 (3)
O2—H2OA···Cl1	0.82 (2)	2.31 (2)	3.135 (5)	174 (6)
N2—H2···Cl1	0.96 (4)	2.08 (4)	3.034 (6)	175 (3)
C5—H5···Cl1	0.93	2.77	3.692 (5)	172

Symmetry codes: (i)  $-x+2, y+1/2, -z+1/2$ ; (ii)  $-x+3/2, -y+1, z+1/2$ .

## supplementary materials

---

Fig. 1




Fig. 2

