

## Redetermination of *trans*-cyclohexane-1,4-diammonium dichloride

Guido J. Reiss\* and Sara Bajorat

Institut für Anorganische Chemie und Strukturchemie, Lehrstuhl für Material- und Strukturforschung, Heinrich-Heine-Universität Düsseldorf, Universitätsstrasse 1, 40225 Düsseldorf, Germany

Correspondence e-mail: reissg@uni-duesseldorf.de


Received 22 November 2007; accepted 30 November 2007

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.038;  $wR$  factor = 0.072; data-to-parameter ratio = 22.4.

A redetermination of the crystal structure of the title compound,  $\text{C}_6\text{H}_{16}\text{N}_2^{2+}\cdot 2\text{Cl}^-$ , was undertaken. All atomic coordinates including those of the H atoms were refined freely. The cation is located on a centre of symmetry. Important for the crystal structure are wavy hydrogen-bonded layers that are formed by ammonium groups and chloride anions, giving hydrogen-bonded  $R_6^3(12)$  rings.

### Related literature

For previous structure determinations, see: Dunitz & Strickler (1965, 1966). For the isostructural cyclohexane-1,4-diammonium dibromide, see: Rademeyer (2006). For hydrogen-bond motifs, see: Etter *et al.* (1990); Rademeyer (2006).



### Experimental

#### Crystal data

$\text{C}_6\text{H}_{16}\text{N}_2^{2+}\cdot 2\text{Cl}^-$	$V = 490.39$ (16) Å <sup>3</sup>
$M_r = 187.11$	$Z = 2$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 5.2550$ (11) Å	$\mu = 0.60$ mm <sup>-1</sup>
$b = 14.890$ (3) Å	$T = 293$ (2) K
$c = 6.3604$ (12) Å	$0.30 \times 0.24 \times 0.20$ mm
$\beta = 99.824$ (18)°	

#### Data collection

Stoe STADI CCD diffractometer	1766 independent reflections
Absorption correction: none	1562 reflections with $I > 2\sigma(I)$
13502 measured reflections	$R_{\text{int}} = 0.043$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	79 parameters
$wR(F^2) = 0.072$	All H-atom parameters refined
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.41$ e Å <sup>-3</sup>
1766 reflections	$\Delta\rho_{\text{min}} = -0.28$ e Å <sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H3}\cdots\text{Cl1}^{\text{ii}}$	0.88 (2)	2.30 (2)	3.1734 (15)	170.3 (16)
$\text{N1}-\text{H2}\cdots\text{Cl1}^{\text{iii}}$	0.86 (2)	2.33 (2)	3.1833 (13)	171.9 (17)
$\text{N1}-\text{H1}\cdots\text{Cl1}$	0.93 (2)	2.23 (2)	3.1584 (13)	173.9 (16)

Symmetry codes: (ii)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (iii)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *CrysAlis CCD* (Kuma Diffraction, 2000); cell refinement: *CrysAlis RED* (Kuma Diffraction, 2000); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *DIAMOND* (Brandenburg, 2001); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GD2030).

### References

- Brandenburg, K. (2001). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Dunitz, J. D. & Strickler, P. (1965). *Helv. Chim. Acta*, **48**, 1450–1456.
- Dunitz, J. D. & Strickler, P. (1966). *Helv. Chim. Acta*, **49**, 2502–2505.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst. B* **46**, 256–262.
- Kuma Diffraction (2000). *CrysAlis CCD* and *CrysAlis RED*. Versions 1.166. Kuma Diffraction Instruments, Wrocław, Poland.
- Rademeyer, M. (2006). *Acta Cryst. E* **62**, o5767–o5769.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.

**supplementary materials**

*Acta Cryst.* (2008). E64, o223 [ doi:10.1107/S1600536807064793 ]

## Redetermination of *trans*-cyclohexane-1,4-diammonium dichloride

G. J. Reiss and S. Bajorat

### Comment

The title compound was first crystallographically characterized by Dunitz and coworkers in 1966 (Dunitz & Strickler, 1965, 1966). This quality structure determination only lacks the fact that all hydrogen atom positions, especially those of the ammonium group, were introduced into the structure model on the basis of geometrically calculated positions, with the N—H and the C—H distances set to 1.1 Å. For X-ray data refinement of hydrogen atom positions significantly shorter values are commonly found. We now describe an improved structure model - the hydrogen atoms were reliably found and refined from quality X-ray data.

A standard refinement using reflections up to  $50^\circ / 2\Theta$  gave the following values:  $R^2 = 6.01$ ,  $R_1 = 2.97$ , GooF = 1.197. Using data with reflections up to  $65^\circ / 2\Theta$  a more stable refinement is possible and the standard uncertainties of the N—H-distances are smaller.

The title structure consists of hydrogen bonded hydrophilic layers in the *ac*-plane. These wavy layers are built by an annulated ring-motif ( $R^3_6(12)$ ; Etter, 1990) constructed by three chloride anions and three ammonium groups (Fig. 3). Each ammonium group donates three hydrogen bonds of only slightly different strength to neighbouring chloride anions (Tab. 2, Fig. 1 + 2). The title compound is therefore isostructural but not isotypic to the cyclohexane-1,4-diammonium dibromide (Rademeyer, 2006).

In terms of crystal engineering the structure of the title compound is dominated by the hydrogen bonded layers. The aliphatic cyclohexane-1,4-diyl fragments connect these layers. According to the positions of the ammonium groups in the hydrogen bonded network the cyclohexyl-fragments do not appear closely packed (Fig. 3).

### Experimental

*trans*-Cyclohexane-1,4-diammonium dichloride was prepared by the reaction of 1,4-diaminocyclohexane (+99%, Aldrich, 0.11 g) and hydrochloric acid (37%) at room temperature. From this colourless solution small block shaped crystals were obtained.

### Refinement

All hydrogen atom positions were obtained from difference fourier maps, all hydrogen atoms were refined freely and with an individual isotropic displacement parameter for each (H—*X* distance range: 0.88–1.01 Å).

## Figures




Fig. 1. The structure of the title compound (displacement ellipsoids at the 40% probability level, H-atoms drawn with arbitrary radius). Thin dashed lines show hydrogen bonds to neighbouring chloride anions. The atoms of the asymmetric unit are labeled.




Fig. 2. Hydrogen bonded layer in the *ac*-plane of the title structure, constructed from annulated  $R^3_6(12)$  motifs. (Symmetry code ' =  $-1/2 + x, 0.5 - y, -1/2 + z$ )





Fig. 3. Crystal packing seen along the *c* direction. Hydrogen bonding interactions are shown as dotted lines.

## *trans*-cyclohexane-1,4-diammonium dichloride

### Crystal data



$M_r = 187.11$

Monoclinic,  $P2_1/n$

$a = 5.2550(11) \text{ \AA}$

$b = 14.890(3) \text{ \AA}$

$c = 6.3604(12) \text{ \AA}$

$\beta = 99.824(18)^\circ$

$V = 490.39(16) \text{ \AA}^3$

$Z = 2$

$F_{000} = 200$

$D_x = 1.267 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1610 reflections

$\theta = 4.8\text{--}17.4^\circ$

$\mu = 0.60 \text{ mm}^{-1}$

$T = 293(2) \text{ K}$

Block, colourless

$0.30 \times 0.24 \times 0.20 \text{ mm}$

### Data collection

Stoe STADI CCD  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293(2) \text{ K}$

1562 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.043$

$\theta_{\text{max}} = 32.5^\circ$

$\theta_{\text{min}} = 4.3^\circ$

$\omega$  scans  $h = -7 \rightarrow 7$   
 Absorption correction: none  $k = -22 \rightarrow 22$   
 13502 measured reflections  $l = -8 \rightarrow 9$   
 1766 independent reflections

### Refinement

Refinement on  $F^2$  Hydrogen site location: difference Fourier map  
 Least-squares matrix: full All H-atom parameters refined  
 $R[F^2 > 2\sigma(F^2)] = 0.038$   $w = 1/[\sigma^2(F_o^2) + 0.3P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $wR(F^2) = 0.072$   $(\Delta/\sigma)_{\max} < 0.001$   
 $S = 1.04$   $\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$   
 1766 reflections  $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$   
 79 parameters Extinction correction: SHELXL,  
 $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$   
 Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.174 (6)  
 Secondary atom site location: difference Fourier map

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.46825 (6)	0.15413 (2)	0.46296 (5)	0.03915 (12)
N1	0.5260 (2)	0.34260 (8)	0.24820 (19)	0.0338 (2)
H1	0.522 (3)	0.2870 (13)	0.316 (3)	0.051 (5)*
H2	0.642 (4)	0.3377 (12)	0.168 (3)	0.057 (5)*
H3	0.377 (4)	0.3498 (12)	0.163 (3)	0.050 (5)*
C1	0.6023 (3)	0.50463 (8)	0.3001 (2)	0.0335 (3)
H11	0.448 (3)	0.5150 (11)	0.198 (3)	0.041 (4)*
H12	0.738 (4)	0.5022 (11)	0.221 (3)	0.043 (4)*
C2	0.5751 (2)	0.41590 (8)	0.41008 (18)	0.0281 (2)
H21	0.726 (3)	0.4018 (10)	0.503 (2)	0.029 (3)*
C3	0.3565 (3)	0.41864 (9)	0.5386 (2)	0.0358 (3)
H31	0.203 (3)	0.4276 (11)	0.445 (3)	0.042 (4)*
H32	0.355 (3)	0.3590 (11)	0.613 (3)	0.044 (4)*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.03850 (18)	0.03987 (18)	0.04168 (19)	-0.00250 (12)	0.01427 (13)	0.00190 (13)
N1	0.0361 (5)	0.0330 (5)	0.0350 (5)	-0.0006 (4)	0.0137 (4)	-0.0018 (4)
C1	0.0416 (6)	0.0331 (6)	0.0301 (6)	-0.0022 (5)	0.0179 (5)	0.0026 (4)
C2	0.0275 (5)	0.0315 (5)	0.0267 (5)	0.0006 (4)	0.0082 (4)	0.0021 (4)
C3	0.0394 (6)	0.0341 (6)	0.0390 (6)	-0.0078 (5)	0.0211 (5)	-0.0008 (5)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

N1—C2	1.4924 (16)	C2—C3	1.5201 (16)
C1—C2	1.5130 (17)	C2—H21	0.928 (15)
C1—C3 <sup>i</sup>	1.5259 (18)	C3—C1 <sup>i</sup>	1.5259 (18)
C1—H11	0.959 (17)	C3—H31	0.927 (17)
C1—H12	0.941 (19)	C3—H32	1.008 (17)
C2—N1—H1	110.2 (11)	N1—C2—C3	109.47 (10)
C2—N1—H2	114.4 (13)	N1—C2—C1	109.79 (10)
H1—N1—H2	105.8 (16)	C3—C2—C1	111.39 (10)
C2—N1—H3	111.6 (12)	N1—C2—H21	107.7 (9)
H1—N1—H3	107.8 (16)	C3—C2—H21	107.6 (9)
H2—N1—H3	106.8 (18)	C1—C2—H21	110.8 (9)
C2—C1—C3 <sup>i</sup>	110.91 (10)	C2—C3—C1 <sup>i</sup>	110.36 (10)
C2—C1—H11	108.6 (10)	C2—C3—H31	108.1 (11)
C3 <sup>i</sup> —C1—H11	110.0 (10)	C1 <sup>i</sup> —C3—H31	109.5 (10)
C2—C1—H12	110.8 (10)	C2—C3—H32	107.4 (10)
C3 <sup>i</sup> —C1—H12	110.9 (10)	C1 <sup>i</sup> —C3—H32	110.8 (10)
H11—C1—H12	105.6 (15)	H31—C3—H32	110.6 (15)
C3 <sup>i</sup> —C1—C2—N1	178.07 (10)	N1—C2—C3—C1 <sup>i</sup>	-177.94 (11)
C3 <sup>i</sup> —C1—C2—C3	56.65 (16)	C1—C2—C3—C1 <sup>i</sup>	-56.34 (16)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H3 $\cdots$ C11 <sup>ii</sup>	0.88 (2)	2.30 (2)	3.1734 (15)	170.3 (16)
N1—H2 $\cdots$ C11 <sup>iii</sup>	0.86 (2)	2.33 (2)	3.1833 (13)	171.9 (17)
N1—H1 $\cdots$ C11	0.93 (2)	2.23 (2)	3.1584 (13)	173.9 (16)

Symmetry codes: (ii)  $x-1/2, -y+1/2, z-1/2$ ; (iii)  $x+1/2, -y+1/2, z-1/2$ .

Fig. 1




Fig. 2






Fig. 3

