

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N,N'*-Bis(pyrimidin-2-yl)terephthalamide dihydrate¹**Suchada Chantrapromma,^a§ Hoong-Kun Fun,^{b*} Subrata Jana,^c Anita Hazra^c and Shyamprosod Goswami^c

^aDepartment of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^cDepartment of Chemistry, Bengal Engineering and Science University, Shibpur, Howrah 711 103, India
Correspondence e-mail: hkfun@usm.my


Received 27 November 2007; accepted 5 December 2007

Key indicators: single-crystal X-ray study; *T* = 100 K; mean $\sigma(\text{C}-\text{C})$ = 0.002 Å; *R* factor = 0.043; *wR* factor = 0.106; data-to-parameter ratio = 14.3.

The organic molecule of the title compound, C₁₆H₁₂N₆O₂·2H₂O, lies across a crystallographic inversion centre. The dihedral angle between the pyrimidine and benzene rings is 80.78 (6)°. The two pyrimidine rings are parallel by virtue of the centre of symmetry. The pyrimidine and benzene rings form dihedral angles of 41.41 (7) and 40.26 (7)°, respectively, with the amide plane. The molecules are linked by N—H···N and C—H···N hydrogen bonds into a two-dimensional network parallel to the (111) plane. O—H···O and C—H···O hydrogen bonds involving the water molecules link the adjacent layers into a three-dimensional network. In addition, a C—H··· π interaction involving the benzene ring is observed.

Related literature

For bond-length data, see: Allen *et al.* (1987). For related literature on supramolecular chemistry, see: Desiraju (1989); Lehn (1995). For related structures, see: Goswami, Jana, Das *et al.* (2007); Goswami, Jana, Hazra *et al.* (2007). For related literature on the coordination chemistry and applications of aminopyrimidine derivatives, see: Aakeroy *et al.* (2006); Etter (1990); Fun *et al.* (2006); Gallagher *et al.* (2004); Goswami & Mahapatra (1999); Smith *et al.* (1998); Wang *et al.* (2006).



¹ This paper is dedicated to His Majesty, Thai King Bhumibol Adulyadej on the occasion of his 80th Birthday Anniversary which fell on December 5th, 2007.

§ Additional correspondence author, email: suchada.c@psu.ac.th.

Experimental

Crystal data

C₁₆H₁₂N₆O₂·2H₂O
M_r = 356.35
Triclinic, *P* $\bar{1}$
a = 5.0733 (1) Å
b = 8.3233 (1) Å
c = 9.9622 (2) Å
 α = 68.220 (1)°
 β = 75.441 (1)°

γ = 82.333 (1)°
V = 377.72 (1) Å³
Z = 1
Mo *K* α radiation
 μ = 0.12 mm⁻¹
T = 100.0 (1) K
0.33 × 0.22 × 0.08 mm

Data collection

Bruker SMART APEXII CCD
area-detector diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)
T_{min} = 0.963, *T_{max}* = 0.991

12751 measured reflections
2202 independent reflections
1982 reflections with *I* > 2 σ (*I*)
R_{int} = 0.035

Refinement

$R[F^2 > 2\sigma(F^2)]$ = 0.044
 $wR(F^2)$ = 0.106
S = 1.12
2202 reflections
154 parameters

3 restraints
All H-atom parameters refined
 $\Delta\rho_{\text{max}}$ = 0.39 e Å⁻³
 $\Delta\rho_{\text{min}}$ = -0.26 e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the benzene ring.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1W—H2WA···O1W ⁱ	0.80 (4)	1.99 (4)	2.791 (2)	171 (6)
O1W—H1W···O1	0.81 (2)	1.98 (2)	2.785 (2)	177 (3)
O1W—H2WB···O1W ⁱⁱ	0.80 (6)	2.04 (5)	2.795 (2)	156 (6)
N3—H1N3···N1 ⁱⁱⁱ	0.89 (2)	2.14 (2)	3.017 (2)	169 (2)
C1—H1···N2 ^{iv}	0.98 (2)	2.61 (2)	3.206 (2)	119 (1)
C2—H2···O1W ^v	0.95 (2)	2.58 (2)	3.508 (2)	164 (1)
C3—H3···Cg1 ^v	0.97 (2)	2.98 (2)	3.914 (2)	164 (2)
C3—H3···Cg1 ⁱⁱⁱ	0.97 (2)	2.98 (2)	3.914 (2)	164 (2)

Symmetry codes: (i) $-x, -y + 2, -z + 1$; (ii) $-x + 1, -y + 2, -z + 1$; (iii) $-x + 1, -y + 1, -z$; (iv) $-x + 2, -y + 2, -z$; (v) $x + 1, y, z - 1$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 1998); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2003).

SJ, AH and SG acknowledge the DST (grant No. SR/S1/OC-13/2005) and CSIR [grant No. 01(1913)/04/EMR-II], Government of India, for financial support. SJ and AH thank the CSIR, Government of India, for research fellowships. SC thanks the Prince of Songkla University, Thailand, for support. The authors also thank the Malaysian Government and Universiti Sains Malaysia for a Scientific Advancement Grant Allocation (SAGA, grant No. 304/PFIZIK/653003/A118).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2534).

References

- Aakeroy, C. B., Schultheiss, N., Desper, J. & Moore, C. (2006). *New J. Chem.* **30**, 1452–1460.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2005). *APEX2* (Version 1.27), *SAINT* (Version 7.12a) and *SADABS* (Version 2004/1). Bruker AXS Inc., Madison, Wisconsin, USA.
- Desiraju, G. R. (1989). *Crystal Engineering. The Design of Organic Solids*. Amsterdam: Elsevier.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Fun, H.-K., Goswami, S., Jana, S. & Chantrapromma, S. (2006). *Acta Cryst. E* **62**, o5332–o5334.
- Gallagher, J. F., Goswami, S., Chatterjee, B., Jana, S. & Dutta, K. (2004). *Acta Cryst. C* **60**, o229–o231.
- Goswami, S., Jana, S., Das, N. K., Fun, H.-K. & Chantrapromma, S. (2007). *J. Mol. Struct.* doi:10.1016/j.molstruc.2007.07.005. In the press.
- Goswami, S., Jana, S., Hazra, A., Fun, H.-K. & Chantrapromma, S. (2007). *Supramol. Chem.* **19**. In the press.
- Goswami, S. P. & Mahapatra, A. K. (1999). *Supramol. Chem.* **11**, 25–33.
- Lehn, J. M. (1995). *Supramolecular Chemistry: Concepts and Perspectives*. Weinheim: VCH.
- Sheldrick, G. M. (1998). *SHELXTL*. Version 5.1. Bruker AXS Inc., Madison, Wisconsin, USA.
- Smith, G., Cloutt, B. A., Lynch, D. E., Byriel, K. A. & Kennard, C. H. L. (1998). *Inorg. Chem.* **37**, 3236–3242.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Wang, Y.-H., Chu, K.-L., Chen, H.-C., Yeh, C.-W., Chan, Z.-K., Suen, M.-C., Chen, J.-D. & Wang, J.-C. (2006). *CrystEngComm*, **8**, 84–93.

supplementary materials

Acta Cryst. (2008). E64, o267-o268 [doi:10.1107/S1600536807065683]

N,N'-Bis(pyrimidin-2-yl)terephthalamide dihydrate

S. Chantrapromma, H.-K. Fun, S. Jana, A. Hazra and S. Goswami

Comment

Substituted 2-aminopyrimidines are very important compounds in molecular recognition and supramolecular chemistry (Desiraju, 1989; Lehn, 1995) for the presence of nice donor-acceptor arrays (Aakeroy *et al.*, 2006; Etter, 1990; Fun *et al.*, 2006; Gallagher, *et al.*, 2004; Goswami & Mahapatra, 1999). The donor-acceptor arrangement of the title molecule differs from the 2-aminopyrimidine based compounds (Goswami, Jana, Das *et al.*, 2007; Goswami, Jana, Hazra *et al.*, 2007). These types of compounds are also important for the metal coordination and related studies (Smith *et al.*, 1998; Wang *et al.*, 2006). The coordination chemistry of the title compound is under investigations.

Molecules of the title compound lie across a crystallographic inversion centre (Fig. 1). All bond lengths and angles have normal values (Allen *et al.*, 1987). The *N,N*-di-pyrimidin-2-yl-terephthalamide molecule has a staggered conformation. The orientation of the pyrimidine ring (C1–C4/N1/N2) with respect to the benzene ring (C6–C8/C6A–C8A) can be described by the dihedral angle formed by these planes of 80.78 (6)° and the torsion angle C4–N3–C5–C6 of –169.50 (11)°. The two pyrimidine rings are parallel by virtue of the centre of symmetry. The pyrimidine and benzene rings form dihedral angles of 41.41 (7)° and 40.26 (7)°, respectively, with the amide plane.

The structure shows O—H···O and C—H···O hydrogen bond between water and *N,N*-di-pyrimidin-2-yl-terephthalamide molecule. In the crystal packing (Fig. 2), the molecules are linked by N—H···N and C—H···N hydrogen bonds into a two-dimensional network parallel to the (1 $\bar{1}$ 1) plane. The O—H···O and C—H···O hydrogen bonds (Table 1) involving the water molecules link the adjacent layers into a three-dimensional network. The crystal structure is further stabilized by C—H··· π interactions involving the benzene ring (centroid Cg1).

Experimental

A solution of terephthaloyl chloride (203 mg, 1 mmol), in dry CH₂Cl₂ (20 ml) was put in a round bottle flask under nitrogen atmosphere. 2-Aminopyrimidine (190 mg, 2 mmol) containing triethylamine (0.55 ml) in dry CH₂Cl₂ (15 ml) was added dropwise. The reaction mixture was stirred at room temperature for 10 h. The reaction mixture was dried, washed with sodium bicarbonate solution, and then extracted with CH₂Cl₂ (4 × 20 ml). The crude mixture was purified by column chromatography (silica gel, 100–200 mesh) using 20% ethyl acetate-petroleum ether solution as eluent to afford a white solid compound (188 mg, 65%). Single crystals of the title compound were grown by slow evaporation of a CHCl₃—CH₃OH (3:1 v/v) solution (m.p. 457–459 K).

Refinement

All H atoms were located from the difference map and isotropically refined. One of the water hydrogen atoms is disordered over two positions with occupancies each 0.50. The O—H distances were restrained to be equal within 0.03 Å.

Figures




Fig. 1. The molecular structure of the title compound, showing 60% probability displacement ellipsoids and the atomic numbering. One of the H atoms of the water molecule is disordered over two positions. The dashed line indicates a hydrogen bond. Atoms labelled with the suffix A are generated by the symmetry operation $(-x, 1 - y, 1 - z)$.




Fig. 2. The crystal packing of the title compound, viewed along the a axis. Hydrogen bonds are shown as dashed lines.

N,N'-Bis(pyrimidin-2-yl)terephthalamide dihydrate

Crystal data

$C_{16}H_{12}N_6O_2 \cdot 2H_2O$

$M_r = 356.35$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 5.0733$ (1) Å

$b = 8.3233$ (1) Å

$c = 9.9622$ (2) Å

$\alpha = 68.220$ (1)°

$\beta = 75.441$ (1)°

$\gamma = 82.333$ (1)°

$V = 377.720$ (12) Å³

$Z = 1$

$F_{000} = 186$

$D_x = 1.567$ Mg m⁻³

Melting point: 457-459 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2202 reflections

$\theta = 2.3$ – 30.0°

$\mu = 0.12$ mm⁻¹

$T = 100.0$ (1) K

Block, colourless

$0.33 \times 0.22 \times 0.08$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 8.33 pixels mm⁻¹

$T = 100.0$ (1) K

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.963$, $T_{\max} = 0.991$

12751 measured reflections

2202 independent reflections

1982 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\text{max}} = 30.0^\circ$

$\theta_{\text{min}} = 2.3^\circ$

$h = -7 \rightarrow 7$

$k = -11 \rightarrow 11$

$l = -13 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.044$$

$$wR(F^2) = 0.106$$

$$S = 1.12$$

2202 reflections

154 parameters

3 restraints

Primary atom site location: structure-invariant direct methods

All H-atom parameters refined

$$w = 1/[\sigma^2(F_o^2) + (0.0387P)^2 + 0.2638P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.39 \text{ e } \text{Å}^{-3}$$

$$\Delta\rho_{\min} = -0.26 \text{ e } \text{Å}^{-3}$$

Extinction correction: none

Special details

Experimental. The data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.40513 (19)	0.85138 (12)	0.21674 (10)	0.0152 (2)	
N1	0.7704 (2)	0.64103 (14)	-0.09155 (12)	0.0131 (2)	
N2	0.9113 (2)	0.77862 (14)	0.05021 (12)	0.0134 (2)	
N3	0.5215 (2)	0.61544 (14)	0.14450 (12)	0.0124 (2)	
H1N3	0.441 (4)	0.530 (2)	0.1395 (19)	0.013 (4)*	
C1	1.1248 (3)	0.83918 (16)	-0.06205 (14)	0.0135 (2)	
H1	1.248 (4)	0.908 (2)	-0.0464 (19)	0.016 (4)*	
C2	1.1741 (3)	0.80284 (17)	-0.19128 (14)	0.0144 (2)	
H2	1.328 (4)	0.846 (2)	-0.270 (2)	0.018 (4)*	
C3	0.9898 (3)	0.70138 (17)	-0.20052 (14)	0.0144 (2)	
H3	1.021 (4)	0.669 (2)	-0.287 (2)	0.015 (4)*	
C4	0.7440 (2)	0.68481 (15)	0.02820 (13)	0.0113 (2)	
C5	0.3775 (2)	0.69938 (16)	0.23761 (13)	0.0121 (2)	
C6	0.1803 (2)	0.59197 (16)	0.37078 (13)	0.0114 (2)	
C7	-0.0689 (3)	0.67033 (16)	0.41794 (14)	0.0131 (2)	
H7	-0.118 (4)	0.789 (2)	0.3599 (19)	0.015 (4)*	
C8	0.2492 (2)	0.42173 (16)	0.45293 (14)	0.0126 (2)	
H8	0.419 (4)	0.368 (2)	0.4211 (19)	0.014 (4)*	
O1W	0.2790 (3)	0.96683 (17)	0.45548 (13)	0.0296 (3)	
H1W	0.309 (5)	0.934 (3)	0.386 (2)	0.034 (6)*	
H2WA	0.123 (6)	0.994 (7)	0.484 (6)	0.050 (15)*	0.50

supplementary materials

H2WB 0.380 (10) 1.013 (7) 0.479 (7) 0.064 (18)* 0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0159 (4)	0.0143 (4)	0.0149 (4)	-0.0037 (3)	0.0011 (3)	-0.0063 (3)
N1	0.0123 (5)	0.0147 (5)	0.0122 (5)	-0.0017 (4)	-0.0013 (4)	-0.0054 (4)
N2	0.0125 (5)	0.0142 (5)	0.0132 (5)	-0.0022 (4)	-0.0009 (4)	-0.0053 (4)
N3	0.0117 (5)	0.0136 (5)	0.0116 (5)	-0.0040 (4)	0.0014 (4)	-0.0057 (4)
C1	0.0118 (5)	0.0128 (5)	0.0151 (6)	-0.0018 (4)	-0.0026 (4)	-0.0040 (4)
C2	0.0110 (5)	0.0168 (6)	0.0128 (5)	-0.0028 (4)	0.0002 (4)	-0.0035 (4)
C3	0.0134 (5)	0.0179 (6)	0.0116 (5)	-0.0016 (4)	-0.0009 (4)	-0.0057 (5)
C4	0.0104 (5)	0.0112 (5)	0.0111 (5)	-0.0006 (4)	-0.0008 (4)	-0.0033 (4)
C5	0.0108 (5)	0.0148 (5)	0.0105 (5)	-0.0018 (4)	-0.0013 (4)	-0.0047 (4)
C6	0.0106 (5)	0.0145 (5)	0.0098 (5)	-0.0035 (4)	-0.0006 (4)	-0.0052 (4)
C7	0.0127 (5)	0.0140 (5)	0.0123 (5)	-0.0012 (4)	-0.0014 (4)	-0.0049 (4)
C8	0.0104 (5)	0.0154 (5)	0.0124 (5)	-0.0014 (4)	-0.0006 (4)	-0.0064 (4)
O1W	0.0388 (7)	0.0353 (7)	0.0186 (5)	-0.0039 (5)	-0.0004 (5)	-0.0170 (5)

Geometric parameters (\AA , $^\circ$)

O1—C5	1.2252 (15)	C3—H3	0.970 (18)
N1—C4	1.3429 (15)	C5—C6	1.5006 (16)
N1—C3	1.3441 (16)	C6—C8	1.3967 (17)
N2—C4	1.3325 (16)	C6—C7	1.3976 (17)
N2—C1	1.3403 (16)	C7—C8 ⁱ	1.3937 (17)
N3—C5	1.3734 (15)	C7—H7	0.980 (18)
N3—C4	1.4037 (15)	C8—C7 ⁱ	1.3937 (17)
N3—H1N3	0.886 (18)	C8—H8	0.951 (18)
C1—C2	1.3850 (17)	O1W—H1W	0.807 (19)
C1—H1	0.976 (18)	O1W—H2WA	0.80 (2)
C2—C3	1.3810 (17)	O1W—H2WB	0.80 (2)
C2—H2	0.956 (18)		
C4—N1—C3	115.24 (10)	N1—C4—N3	115.16 (10)
C4—N2—C1	115.74 (11)	O1—C5—N3	123.59 (11)
C5—N3—C4	123.80 (10)	O1—C5—C6	120.82 (11)
C5—N3—H1N3	117.2 (11)	N3—C5—C6	115.59 (10)
C4—N3—H1N3	116.3 (11)	C8—C6—C7	120.14 (11)
N2—C1—C2	122.42 (11)	C8—C6—C5	121.39 (11)
N2—C1—H1	116.0 (10)	C7—C6—C5	118.30 (11)
C2—C1—H1	121.5 (10)	C8 ⁱ —C7—C6	119.91 (11)
C3—C2—C1	116.73 (11)	C8 ⁱ —C7—H7	119.9 (10)
C3—C2—H2	121.6 (11)	C6—C7—H7	120.2 (10)
C1—C2—H2	121.7 (11)	C7 ⁱ —C8—C6	119.95 (11)
N1—C3—C2	122.62 (11)	C7 ⁱ —C8—H8	119.9 (11)
N1—C3—H3	118.1 (11)	C6—C8—H8	120.2 (11)
C2—C3—H3	119.3 (11)	H1W—O1W—H2WA	116 (4)

N2—C4—N1	127.22 (11)	H1W—O1W—H2WB	128 (5)
N2—C4—N3	117.53 (11)	H2WA—O1W—H2WB	111 (6)
C4—N2—C1—C2	-1.83 (18)	C4—N3—C5—O1	10.1 (2)
N2—C1—C2—C3	0.61 (19)	C4—N3—C5—C6	-169.50 (11)
C4—N1—C3—C2	-0.68 (19)	O1—C5—C6—C8	-137.21 (13)
C1—C2—C3—N1	0.72 (19)	N3—C5—C6—C8	42.36 (16)
C1—N2—C4—N1	1.96 (19)	O1—C5—C6—C7	38.15 (17)
C1—N2—C4—N3	178.50 (11)	N3—C5—C6—C7	-142.27 (12)
C3—N1—C4—N2	-0.73 (19)	C8—C6—C7—C8 ⁱ	-0.1 (2)
C3—N1—C4—N3	-177.34 (11)	C5—C6—C7—C8 ⁱ	-175.54 (11)
C5—N3—C4—N2	36.36 (18)	C7—C6—C8—C7 ⁱ	0.1 (2)
C5—N3—C4—N1	-146.68 (12)	C5—C6—C8—C7 ⁱ	175.39 (11)

Symmetry codes: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H2WA \cdots O1W ⁱⁱ	0.80 (4)	1.99 (4)	2.791 (2)	171 (6)
O1W—H1W \cdots O1	0.81 (2)	1.98 (2)	2.785 (2)	177 (3)
O1W—H2WB \cdots O1W ⁱⁱⁱ	0.80 (6)	2.04 (5)	2.795 (2)	156 (6)
N3—H1N3 \cdots N1 ^{iv}	0.89 (2)	2.14 (2)	3.017 (2)	169 (2)
C1—H1 \cdots N2 ^v	0.98 (2)	2.61 (2)	3.206 (2)	119 (1)
C2—H2 \cdots O1W ^v	0.95 (2)	2.58 (2)	3.508 (2)	164 (1)
C3—H3 \cdots Cg1 ^{vi}	0.97 (2)	2.98 (2)	3.914 (2)	164 (2)
C3—H3 \cdots Cg1 ^{iv}	0.97 (2)	2.98 (2)	3.914 (2)	164 (2)

Symmetry codes: (ii) $-x, -y+2, -z+1$; (iii) $-x+1, -y+2, -z+1$; (iv) $-x+1, -y+1, -z$; (v) $-x+2, -y+2, -z$; (vi) $x+1, y, z-1$.

Fig. 1




Fig. 2

