

5,6-Dioxo-1,10-phenanthrolin-1-ium nitrate

Alireza Abbasi,* Alireza Badiei and Sara Tarighi

School of Chemistry, University College of Science, University of Tehran, Tehran, Iran

Correspondence e-mail: aabbasi@khayam.ut.ac.ir


Received 29 November 2007; accepted 8 December 2007

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.038; wR factor = 0.060; data-to-parameter ratio = 11.5.

In the title salt, $\text{C}_{12}\text{H}_7\text{N}_2\text{O}_2^+\cdot\text{NO}_3^-$, the monoprotonated cation is connected to the nitrate anion by a hydrogen bond. Weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds hold the planar cations together in a layer structure.

Related literature

For related literature, see Fujihara *et al.* (2004); Larsson & Ohström (2004). For the bromide salt, see: Bomfim *et al.* (2003).



Experimental

Crystal data

$\text{C}_{12}\text{H}_7\text{N}_2\text{O}_2^+\cdot\text{NO}_3^-$
 $M_r = 273.21$
 Monoclinic, $C2/c$
 $a = 14.4860$ (19) Å
 $b = 12.5177$ (13) Å
 $c = 13.4535$ (16) Å
 $\beta = 101.720$ (12)°

$V = 2388.7$ (5) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.12$ mm⁻¹
 $T = 290$ (2) K
 $0.30 \times 0.10 \times 0.08$ mm

Data collection

Oxford Diffraction Xcalibur 3 CCD diffractometer
 Absorption correction: numerical (*X-RED*; Stoe & Cie, 1997)
 $T_{\min} = 0.960$, $T_{\max} = 0.989$

7311 measured reflections
 2095 independent reflections
 635 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.068$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.060$
 $S = 0.83$
 2095 reflections

182 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.31$ e Å⁻³
 $\Delta\rho_{\min} = -0.14$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O5}$	0.86	1.95	2.703 (4)	145
$\text{C8}-\text{H8}\cdots\text{O5}^{\text{i}}$	0.93	2.48	3.396 (4)	167
$\text{C9}-\text{H9}\cdots\text{O4}^{\text{i}}$	0.93	2.71	3.359 (5)	128
$\text{C4}-\text{H4}\cdots\text{O4}^{\text{ii}}$	0.93	2.61	3.323 (4)	134
$\text{C3}-\text{H3}\cdots\text{O3}^{\text{iii}}$	0.93	2.40	3.209 (4)	146
$\text{C10}-\text{H10}\cdots\text{O1}^{\text{iii}}$	0.93	2.47	3.318 (5)	151
$\text{C5}-\text{H5}\cdots\text{O2}^{\text{iv}}$	0.93	2.37	3.266 (5)	162

Symmetry codes: (i) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (ii) $x + \frac{1}{2}, y - \frac{1}{2}, z$; (iii) $x - \frac{1}{2}, y + \frac{1}{2}, z$; (iv) $x - \frac{1}{2}, y - \frac{1}{2}, z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2003); cell refinement: *CrysAlis CCD*; data reduction: *CrysAlis RED* (Oxford Diffraction, 2003); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *DIAMOND* (Brandenburg, 2001); software used to prepare material for publication: *PLATON* (Spek, 2003).

This work was supported by grants from the University of Tehran and the Swedish Research Council. The authors are grateful to Professor Magnus Sandström for his support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2405).

References

- Bomfim, J. A. S., Filgueiras, C. A. L., Howie, R. A. & Wardell, J. L. (2003). *Acta Cryst.* **E59**, o244–o246.
 Brandenburg, K. (2001). *DIAMOND*. Version 2.1e. Crystal Impact GbR, Bonn, Germany.
 Fujihara, T., Wada, T. & Tanaka, K. (2004). *Dalton Trans.*, pp. 645–652.
 Larsson, K. & Ohström, L. (2004). *Inorg. Chim. Acta*, **357**, 657–664.
 Oxford Diffraction (2003). *CrysAlis CCD* and *CrysAlis RED*. Versions 1.171.29.2. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
 Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Stoe & Cie (1997). *X-RED*. Version 1.09. Stoe & Cie GmbH, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2008). E64, o290 [doi:10.1107/S1600536807066172]

5,6-Dioxo-1,10-phenanthroline-1-ium nitrate

A. Abbasi, A. Badiei and S. Tarighi

Comment

From a survey of the Cambridge Structural Database [version 5.28, updated Jan. 2007], we found two complexes containing O-protonated pdon (Fujihara, *et al.*, 2004; Larsson, & Ohström, 2004). However, up to now, only one molecule is reported for the N-protonated pdon ($C_{12}H_7N_2O_2^+$, Hpdon) with bromide ions as counter ion (Bomfim *et al.*, 2003).

The molecular structure and atom-labeling scheme for (I) are shown in Fig. 1. There is a relatively strong hydrogen bond between the nitrate ion and to the protonated nitrogen atom in Hpdon (N1—H1 \cdots O5, 2.703 (4) Å). Weak C—H \cdots O hydrogen bonds hold the approximately planar cations together in a layer structure (Fig. 2). The packing of the layers is similar to that in the bromide salt of Hpdon (Bomfim *et al.*, 2003) without specific directional interactions between the layers. Ring A (C1—C5/N1), B (C6—C10/N2) and C (C11—C12/C11/C7/C6) in the cation ($C_{12}H_7N_2O_2^+$, Hpdon) is approximately planar. The interplanar angle between the least-square planes defined by ring A and two other rings, B and C, are 1.69 (5)° and 1.17 (5)°, respectively. While, rings B and C are almost planar, 0.52 (5)°. However, the C—N—C angle is affected by protonation and causes to increase from 116.6 (3)° to 122.7 (3)° in the rings B and A, respectively. Similar effect has also been shown in the previously reported Hpdon bromide by increasing from 116.8 (3)° to 123.2 (3)° for the corresponding values. The comparison of the torsion angle, O1—C12—C11—O2, 2.5 (6)°, in (I) with the molecular pdon indicates that the N-protonation in (I) doesn't significantly affect on dione groups.

Experimental

The title compound was obtained during the reaction of thorium nitrate, $Th(NO_3)_4 \cdot 5H_2O$ (Merck, 99%), and 1,10-phenanthroline-5,6-dione (Aldrich, 97%) in ethanolic solution. Pale-yellow crystals of Hpdon were taken from the obtained greenish precipitates. The presence of water in the ethanolic solution and the high basicity of dione ligand can be the reason for the protonation, providing the $C_{12}H_7N_2O_2^+$, Hpdon cation. Thin fragile layer crystals were obtained by recrystallization from acetonitrile.

Refinement

All H atoms were geometrically positioned and constrained to ride on their parent atoms, with $U_{iso}(H) = 1.2$ times $U_{eq}(C,N)$.

Figures




Fig. 1. Molecular structure of (I), with 50% probability displacement ellipsoids. H atoms are shown as circles of arbitrary radii.




Fig. 2. Packing view for (I), showing the hydrogen bonds as dashed lines.

5,6-Dioxo-1,10-phenanthroline-1-ium nitrate

Crystal data

$C_{12}H_7N_2O_2^+ \cdot NO_3^-$

$M_r = 273.21$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 14.4860$ (19) Å

$b = 12.5177$ (13) Å

$c = 13.4535$ (16) Å

$\beta = 101.720$ (12)°

$V = 2388.7$ (5) Å³

$Z = 8$

$F_{000} = 1120$

$D_x = 1.519$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 7311 reflections

$\theta = 3.5\text{--}25.0^\circ$

$\mu = 0.12$ mm⁻¹

$T = 290$ (2) K

Needle, pale-yellow

$0.30 \times 0.10 \times 0.08$ mm

Data collection

Oxford Diffraction Xcalibur 3 CCD diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 12 pixels mm⁻¹

$T = 290$ (2) K

ω -scans at different ϕ

Absorption correction: numerical (X-RED; Stoe & Cie, 1997)

2095 independent reflections

635 reflections with $I > 2\sigma(I)$

$R_{int} = 0.068$

$\theta_{max} = 25.0^\circ$

$\theta_{min} = 3.8^\circ$

$h = -17 \rightarrow 17$

$k = -14 \rightarrow 14$

$T_{\min} = 0.960$, $T_{\max} = 0.989$
7311 measured reflections

$l = -10 \rightarrow 15$

Refinement

Refinement on F^2

Hydrogen site location: inferred from neighbouring sites

Least-squares matrix: full

H-atom parameters constrained

$R[F^2 > 2\sigma(F^2)] = 0.038$

$$w = 1/[\sigma^2(F_o^2) + (0.01P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$

$wR(F^2) = 0.060$

$(\Delta/\sigma)_{\max} < 0.001$

$S = 0.83$

$\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$

2095 reflections

$\Delta\rho_{\min} = -0.14 \text{ e } \text{\AA}^{-3}$

182 parameters

Extinction correction: SHELXL97 (Sheldrick, 1997),

$$F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$$

Primary atom site location: structure-invariant direct methods

Extinction coefficient: 0.00072 (4)

Secondary atom site location: difference Fourier map

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.85417 (17)	-0.13522 (19)	0.12439 (18)	0.0673 (8)
O2	0.85003 (19)	0.08376 (19)	0.1238 (2)	0.0834 (10)
O3	0.30299 (14)	0.0377 (2)	0.07097 (16)	0.0649 (7)
O4	0.22236 (15)	-0.03456 (19)	0.17219 (15)	0.0613 (6)
O5	0.34903 (16)	-0.11186 (18)	0.14632 (16)	0.0622 (7)
N1	0.52797 (19)	-0.1426 (2)	0.12265 (18)	0.0439 (9)
H1	0.4776	-0.1063	0.1230	0.053*
N2	0.5205 (2)	0.0692 (2)	0.1274 (2)	0.0488 (9)
N3	0.29121 (19)	-0.0361 (3)	0.1292 (2)	0.0476 (8)
C1	0.6089 (3)	-0.0896 (3)	0.1224 (2)	0.0370 (9)
C2	0.6903 (3)	-0.1473 (3)	0.1211 (2)	0.0374 (10)
C3	0.6870 (3)	-0.2573 (3)	0.1201 (2)	0.0528 (11)
H3	0.7411	-0.2969	0.1191	0.063*
C4	0.6018 (3)	-0.3086 (3)	0.1207 (2)	0.0550 (11)

supplementary materials

H4	0.5986	-0.3828	0.1198	0.066*
C5	0.5228 (3)	-0.2490 (3)	0.1224 (2)	0.0502 (11)
H5	0.4657	-0.2826	0.1235	0.060*
C6	0.6062 (3)	0.0275 (3)	0.1258 (2)	0.0380 (9)
C7	0.6867 (3)	0.0879 (3)	0.1278 (2)	0.0421 (10)
C8	0.6798 (3)	0.1978 (3)	0.1312 (2)	0.0534 (13)
H8	0.7320	0.2411	0.1319	0.064*
C9	0.5922 (3)	0.2415 (3)	0.1336 (2)	0.0589 (12)
H9	0.5852	0.3151	0.1369	0.071*
C10	0.5151 (3)	0.1749 (3)	0.1308 (3)	0.0589 (13)
H10	0.4568	0.2058	0.1314	0.071*
C11	0.7780 (3)	0.0351 (4)	0.1244 (2)	0.0526 (10)
C12	0.7807 (3)	-0.0894 (3)	0.1227 (2)	0.0469 (11)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0405 (19)	0.064 (2)	0.102 (2)	0.0165 (15)	0.0270 (16)	0.0046 (15)
O2	0.051 (2)	0.062 (2)	0.146 (2)	-0.0150 (16)	0.0412 (18)	-0.0002 (17)
O3	0.0736 (19)	0.0573 (18)	0.0696 (16)	0.0079 (15)	0.0282 (13)	0.0205 (15)
O4	0.0508 (18)	0.0664 (15)	0.0730 (16)	0.0081 (16)	0.0276 (13)	0.0048 (14)
O5	0.0453 (17)	0.0477 (18)	0.0977 (18)	0.0163 (14)	0.0242 (14)	0.0116 (15)
N1	0.036 (2)	0.038 (2)	0.0605 (19)	0.0038 (18)	0.0143 (17)	0.0013 (19)
N2	0.054 (2)	0.035 (2)	0.0620 (19)	0.0086 (17)	0.0214 (18)	-0.0027 (17)
N3	0.041 (2)	0.051 (2)	0.053 (2)	-0.005 (2)	0.0144 (17)	-0.001 (2)
C1	0.034 (3)	0.038 (3)	0.041 (2)	-0.007 (2)	0.012 (2)	-0.003 (2)
C2	0.037 (3)	0.033 (3)	0.044 (2)	0.008 (2)	0.011 (2)	0.006 (2)
C3	0.056 (3)	0.043 (3)	0.065 (3)	0.009 (3)	0.023 (2)	0.002 (2)
C4	0.067 (4)	0.034 (3)	0.066 (3)	0.001 (3)	0.020 (2)	0.004 (2)
C5	0.050 (3)	0.035 (3)	0.068 (3)	-0.007 (2)	0.017 (2)	0.000 (2)
C6	0.037 (3)	0.035 (3)	0.042 (2)	-0.001 (3)	0.0093 (19)	-0.010 (3)
C7	0.038 (3)	0.036 (3)	0.053 (2)	0.002 (2)	0.011 (2)	0.002 (2)
C8	0.060 (4)	0.030 (3)	0.070 (3)	-0.005 (2)	0.012 (2)	0.003 (2)
C9	0.073 (4)	0.037 (3)	0.065 (3)	0.015 (3)	0.010 (3)	-0.001 (2)
C10	0.058 (4)	0.045 (3)	0.075 (3)	0.005 (2)	0.016 (3)	-0.003 (2)
C11	0.052 (3)	0.051 (3)	0.059 (2)	-0.002 (3)	0.021 (2)	0.001 (3)
C12	0.050 (3)	0.048 (3)	0.046 (2)	0.000 (3)	0.018 (2)	-0.002 (2)

Geometric parameters (\AA , $^\circ$)

O1—C12	1.206 (4)	C3—C4	1.392 (4)
O2—C11	1.210 (3)	C3—H3	0.9300
O3—N3	1.244 (3)	C4—C5	1.371 (4)
O4—N3	1.251 (3)	C4—H4	0.9300
O5—N3	1.256 (3)	C5—H5	0.9300
N1—C5	1.335 (4)	C6—C7	1.385 (4)
N1—C1	1.347 (4)	C7—C8	1.381 (4)
N1—H1	0.8600	C7—C11	1.488 (4)
N2—C10	1.326 (4)	C8—C9	1.387 (4)

N2—C6	1.350 (4)	C8—H8	0.9300
C1—C2	1.387 (4)	C9—C10	1.389 (4)
C1—C6	1.467 (4)	C9—H9	0.9300
C2—C3	1.377 (4)	C10—H10	0.9300
C2—C12	1.492 (4)	C11—C12	1.559 (4)
C5—N1—C1	122.7 (3)	N2—C6—C7	124.1 (4)
C5—N1—H1	118.7	N2—C6—C1	114.7 (4)
C1—N1—H1	118.7	C7—C6—C1	121.2 (4)
C10—N2—C6	116.6 (3)	C8—C7—C6	118.5 (4)
O3—N3—O4	120.3 (3)	C8—C7—C11	120.9 (4)
O3—N3—O5	120.3 (3)	C6—C7—C11	120.5 (4)
O4—N3—O5	119.4 (3)	C7—C8—C9	117.8 (4)
N1—C1—C2	119.1 (4)	C7—C8—H8	121.1
N1—C1—C6	117.6 (4)	C9—C8—H8	121.1
C2—C1—C6	123.3 (4)	C8—C9—C10	119.8 (4)
C3—C2—C1	119.5 (4)	C8—C9—H9	120.1
C3—C2—C12	121.0 (4)	C10—C9—H9	120.1
C1—C2—C12	119.5 (4)	N2—C10—C9	123.1 (4)
C2—C3—C4	119.4 (4)	N2—C10—H10	118.4
C2—C3—H3	120.3	C9—C10—H10	118.4
C4—C3—H3	120.3	O2—C11—C7	123.4 (5)
C5—C4—C3	119.6 (4)	O2—C11—C12	118.6 (4)
C5—C4—H4	120.2	C7—C11—C12	118.0 (4)
C3—C4—H4	120.2	O1—C12—C2	122.5 (4)
N1—C5—C4	119.8 (4)	O1—C12—C11	120.0 (4)
N1—C5—H5	120.1	C2—C12—C11	117.5 (4)
C4—C5—H5	120.1		
C5—N1—C1—C2	0.5 (5)	N2—C6—C7—C11	-179.1 (3)
C5—N1—C1—C6	-178.2 (3)	C1—C6—C7—C11	1.2 (5)
N1—C1—C2—C3	0.0 (5)	C6—C7—C8—C9	0.7 (5)
C6—C1—C2—C3	178.6 (3)	C11—C7—C8—C9	179.5 (3)
N1—C1—C2—C12	-178.9 (3)	C7—C8—C9—C10	-1.0 (5)
C6—C1—C2—C12	-0.3 (5)	C6—N2—C10—C9	-0.5 (5)
C1—C2—C3—C4	-0.1 (5)	C8—C9—C10—N2	1.0 (6)
C12—C2—C3—C4	178.7 (3)	C8—C7—C11—O2	-0.4 (5)
C2—C3—C4—C5	-0.2 (5)	C6—C7—C11—O2	178.4 (3)
C1—N1—C5—C4	-0.8 (5)	C8—C7—C11—C12	178.9 (3)
C3—C4—C5—N1	0.6 (6)	C6—C7—C11—C12	-2.3 (5)
C10—N2—C6—C7	0.2 (5)	C3—C2—C12—O1	-0.9 (5)
C10—N2—C6—C1	179.8 (3)	C1—C2—C12—O1	178.0 (4)
N1—C1—C6—N2	-0.9 (5)	C3—C2—C12—C11	-179.6 (3)
C2—C1—C6—N2	-179.5 (3)	C1—C2—C12—C11	-0.8 (4)
N1—C1—C6—C7	178.7 (3)	O2—C11—C12—O1	2.5 (6)
C2—C1—C6—C7	0.1 (5)	C7—C11—C12—O1	-176.8 (3)
N2—C6—C7—C8	-0.3 (5)	O2—C11—C12—C2	-178.6 (3)
C1—C6—C7—C8	-179.9 (3)	C7—C11—C12—C2	2.1 (5)

supplementary materials

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O5	0.86	1.95	2.703 (4)	145
C8—H8 \cdots O5 ⁱ	0.93	2.48	3.396 (4)	167
C9—H9 \cdots O4 ⁱ	0.93	2.71	3.359 (5)	128
C4—H4 \cdots O4 ⁱⁱ	0.93	2.61	3.323 (4)	134
C3—H3 \cdots O3 ⁱⁱ	0.93	2.40	3.209 (4)	146
C10—H10 \cdots O1 ⁱⁱⁱ	0.93	2.47	3.318 (5)	151
C5—H5 \cdots O2 ^{iv}	0.93	2.37	3.266 (5)	162

Symmetry codes: (i) $x+1/2, y+1/2, z$; (ii) $x+1/2, y-1/2, z$; (iii) $x-1/2, y+1/2, z$; (iv) $x-1/2, y-1/2, z$.

Fig. 1




Fig. 2

