

## Tris(ethane-1,2-diamine)copper(II) bis(trifluoroacetate)

Elena V. Karpova,<sup>a</sup> Maxim A. Zakharov,<sup>b</sup> Alexandr I. Boltalin<sup>a</sup> and Victor B. Rybakov<sup>b\*</sup>

<sup>a</sup>Inorganic Chemistry Division, Chemistry Department, Moscow State University, 119991 Leninskie Gory 1-3, Moscow, Russian Federation, and <sup>b</sup>General Chemistry Division, Chemistry Department, Moscow State University, 119991 Leninskie Gory 1-3, Moscow, Russian Federation

Correspondence e-mail: rybakov20021@yandex.ru


Received 18 December 2007; accepted 14 January 2008

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; disorder in main residue;  $R$  factor = 0.039;  $wR$  factor = 0.092; data-to-parameter ratio = 18.0.

In the title complex,  $[\text{Cu}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_3](\text{CF}_3\text{COO})_2$ , the environment of the Cu atom is distorted octahedral, formed by six N atoms from three chelating ethane-1,2-diamine ligands. The Cu–N distances range from 2.050 (2) to 2.300 (2) Å. This complex cation and the two trifluoroacetate anions are connected by weak N–H $\cdots$ O and N–H $\cdots$ F hydrogen bonds, forming a three-dimensional framework. In both anions, the F atoms are disordered over two positions; in one the site-occupancy factors are 0.55 and 0.45, in the other the values are 0.69 and 0.31.

### Related literature

For other carboxylate complexes, see: Karpova *et al.* (1998); Karpova *et al.* (2001); Gutnikov *et al.* (2006); Karpova *et al.* (2007).



### Experimental

#### Crystal data

$[\text{Cu}(\text{C}_2\text{H}_8\text{N}_2)_3](\text{C}_2\text{F}_3\text{O}_2)_2$

$M_r = 469.90$

Triclinic,  $P\bar{1}$

$a = 8.582$  (6) Å

$b = 9.316$  (6) Å

$c = 12.859$  (7) Å

$\alpha = 74.73$  (3)°

$\beta = 84.69$  (4)°

$\gamma = 69.56$  (3)°

$V = 929.3$  (10) Å<sup>3</sup>

$Z = 2$

Mo  $K\alpha$  radiation

$\mu = 1.26$  mm<sup>-1</sup>

$T = 293$  (2) K

$0.15 \times 0.1 \times 0.08$  mm

#### Data collection

Bruker SMART CCD area-detector  
diffractometer  
Absorption correction: none  
5406 independent reflections  
4277 reflections with  $I > 2\sigma(I)$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.092$   
 $S = 0.94$   
5406 reflections  
300 parameters  
45 restraints  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.24$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2–H2B $\cdots$ F1A	0.90	2.54	3.248 (17)	136
N2–H2B $\cdots$ F3B	0.90	2.55	3.41 (2)	161
N6–H6B $\cdots$ O1	0.90	2.15	3.048 (3)	179
N1–H1A $\cdots$ O2 <sup>i</sup>	0.90	2.35	3.141 (3)	147
N1–H1A $\cdots$ O1 <sup>i</sup>	0.90	2.53	3.377 (4)	156
N4–H4A $\cdots$ O2 <sup>ii</sup>	0.90	2.34	3.175 (3)	154
N6–H6A $\cdots$ O1 <sup>i</sup>	0.90	2.56	3.326 (3)	143
N1–H1B $\cdots$ O2 <sup>iii</sup>	0.90	2.11	3.002 (3)	173
N2–H2A $\cdots$ F4A <sup>ii</sup>	0.90	2.47	3.267 (15)	148
N3–H3A $\cdots$ O3 <sup>iii</sup>	0.90	2.09	2.958 (3)	162
N5–H5A $\cdots$ O3 <sup>iii</sup>	0.90	2.36	3.126 (3)	143
N2–H2A $\cdots$ O4 <sup>iii</sup>	0.90	2.41	3.037 (3)	127
N4–H4B $\cdots$ O4 <sup>iv</sup>	0.90	2.11	2.997 (3)	168
N5–H5B $\cdots$ O3 <sup>iv</sup>	0.90	2.13	3.013 (3)	167

Symmetry codes: (i)  $-x + 2, -y + 1, -z + 2$ ; (ii)  $x, y + 1, z$ ; (iii)  $-x + 1, -y + 1, -z + 1$ ; (iv)  $-x + 2, -y + 1, -z + 1$ .

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINTE* (Bruker, 2000); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97*; molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors gratefully acknowledge Zoja A. Starikova for the X-ray data collection. This investigation was supported by the Russian Fund of Basic Research (project No. 05–03–33038-a).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2235).

### References

- Bruker (2000). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gutnikov, S. I., Karpova, E. V., Zakharov, M. A. & Boltalin, A. I. (2006). *Russ. J. Inorg. Chem.* **51**, 541–548.
- Karpova, E. V., Boltalin, A. I., Korenev, Yu. M., Zakharov, M. A. & Troyanov, S. I. (2001). *Russ. J. Coord. Chem.* **27**, 286–291.
- Karpova, E. V., Boltalin, A. I., Zakharov, M. A., Sorokina, N. I., Korenev, Yu. M. & Troyanov, S. I. (1998). *Z. Anorg. Allg. Chem.* **B624**, 741–744.
- Karpova, E. V., Zakharov, M. A., Gutnikov, S. I. & Boltalin, A. I. (2007). *Acta Cryst.* **E63**, m658–m659.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2008). E64, m373 [ doi:10.1107/S1600536808001438 ]

## Tris(ethane-1,2-diamine)copper(II) bis(trifluoroacetate)

E. V. Karpova, M. A. Zakharov, A. I. Boltalin and V. B. Rybakov

### Comment

The present investigation is a continuation of experimental work to study the structure and properties of different carboxylate complexes (Karpova *et al.*, 1998; Karpova *et al.*, 2001; Gutnikov *et al.*, 2006; Karpova *et al.*, 2007). In the title compound,  $\text{Cu}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_3 \cdot (\text{CF}_3\text{COO})_2$ , the asymmetric unit consists of a complex cation and two crystallographically independent anions. The environment of the Cu atom is distorted octahedral, formed by six N atoms from three chelate ethylenediamine groups. The two trifluoroacetate anions form N—H $\cdots$ O and N—H $\cdots$ F hydrogen bonds with  $\text{NH}_2$  groups of the cation, forming a three-dimensional framework.

### Experimental

An ethanol solution of  $[\text{Cu}(\text{CF}_3\text{COO})_2(\text{CH}_3\text{CN})]_2(\text{CH}_3\text{CN})_2$  was mixed with ethylenediamine in a 1:4 molar ratio. After several days blue prism-shaped crystals were formed in a desiccator over  $\text{P}_4\text{O}_{10}$ .

### Refinement

All H atoms were positioned geometrically and refined using a riding model (including free rotation about C—C or C—N bonds) with C—H = 0.97 Å, N—H = 0.90 Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ . The DELU option in *SHELXL97* was used with parameters 0.0001, 0.0001 for all bonds Cu1—N; for all C—O and C—F bonds. The MERG option with parameter 2 was used before refinement. The F atoms are disordered over two positions, with site occupancy ratios 0.55 (3)/0.45 (3) and 0.69 (2)/0.31 (2).

### Figures




Fig. 1. A view of the title compound, with the atom numbering scheme. Displacement ellipsoids are shown at the 50% probability level. H atoms are drawn as small spheres of arbitrary radius. Only the major disorder components are shown.

## Tris(ethane-1,2-diamine)copper(II) bis(trifluoroacetate)

### Crystal data

$[\text{Cu}(\text{C}_2\text{H}_8\text{N}_2)_3](\text{C}_2\text{F}_3\text{O}_2)_2$

$M_r = 469.90$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$Z = 2$

$F_{000} = 482$

$D_x = 1.679 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

# supplementary materials

---

$a = 8.582 (6) \text{ \AA}$	Cell parameters from 250 reflections
$b = 9.316 (6) \text{ \AA}$	$\theta = 18\text{--}24^\circ$
$c = 12.859 (7) \text{ \AA}$	$\mu = 1.26 \text{ mm}^{-1}$
$\alpha = 74.73 (3)^\circ$	$T = 293 (2) \text{ K}$
$\beta = 84.69 (4)^\circ$	Prism, blue
$\gamma = 69.56 (3)^\circ$	$0.15 \times 0.1 \times 0.08 \text{ mm}$
$V = 929.3 (10) \text{ \AA}^3$	

## Data collection

Bruker SMART CCD area-detector diffractometer	5406 independent reflections
Radiation source: fine-focus sealed tube	4277 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$\theta_{\max} = 30.0^\circ$
Detector resolution: $0.1 \text{ pixels mm}^{-1}$	$\theta_{\min} = 1.6^\circ$
$T = 293(2) \text{ K}$	$h = -11 \rightarrow 12$
$\varphi$ and $\omega$ scans	$k = 0 \rightarrow 13$
Absorption correction: none	$l = -17 \rightarrow 18$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H-atom parameters constrained
$wR(F^2) = 0.092$	$w = 1/[\sigma^2(F_o^2) + (0.0567P)^2]$
$S = 0.94$	where $P = (F_o^2 + 2F_c^2)/3$
5406 reflections	$(\Delta/\sigma)_{\max} = 0.001$
300 parameters	$\Delta\rho_{\max} = 0.24 \text{ e \AA}^{-3}$
45 restraints	$\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

## Special details

**Experimental.** The original *HKL* file was deleted and the present study was conducted using the merged data set.

**Geometry.** All s.u.'s (except the s.u.'s in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.94303 (3)	0.85236 (3)	0.778785 (18)	0.03254 (8)	
N1	0.8821 (2)	0.8734 (2)	0.93404 (13)	0.0444 (3)	
H1A	0.9694	0.8131	0.9777	0.053*	
H1B	0.8587	0.9745	0.9362	0.053*	
C1	0.7376 (3)	0.8237 (3)	0.97256 (17)	0.0486 (5)	
H1C	0.6921	0.8610	1.0363	0.058*	
H1D	0.7726	0.7092	0.9923	0.058*	
C2	0.6060 (3)	0.8885 (3)	0.88718 (19)	0.0503 (5)	
H2C	0.5139	0.8505	0.9121	0.060*	
H2D	0.5641	1.0031	0.8708	0.060*	
N2	0.6812 (2)	0.8352 (2)	0.79038 (16)	0.0495 (4)	
H2A	0.6212	0.8977	0.7314	0.059*	
H2B	0.6866	0.7350	0.7968	0.059*	
N3	0.8604 (2)	1.1007 (2)	0.71134 (15)	0.0471 (4)	
H3A	0.8004	1.1244	0.6515	0.057*	
H3B	0.7948	1.1505	0.7589	0.057*	
C3	1.0036 (3)	1.1543 (3)	0.68450 (19)	0.0539 (5)	
H3C	0.9672	1.2681	0.6739	0.065*	
H3D	1.0531	1.1292	0.6177	0.065*	
C4	1.1308 (3)	1.0764 (3)	0.77268 (19)	0.0510 (5)	
H4C	1.2266	1.1102	0.7523	0.061*	
H4D	1.0843	1.1079	0.8382	0.061*	
N4	1.1826 (2)	0.9040 (2)	0.79228 (15)	0.0457 (4)	
H4A	1.2277	0.8564	0.8583	0.055*	
H4B	1.2577	0.8695	0.7428	0.055*	
N5	0.9835 (2)	0.8096 (2)	0.62842 (13)	0.0453 (4)	
H5A	0.8923	0.8658	0.5873	0.054*	
H5B	1.0688	0.8396	0.5965	0.054*	
C5	1.0217 (3)	0.6405 (3)	0.6384 (2)	0.0540 (5)	
H5C	1.0712	0.6134	0.5721	0.065*	
H5D	0.9208	0.6139	0.6522	0.065*	
C6	1.1414 (3)	0.5510 (3)	0.7304 (2)	0.0541 (5)	
H6C	1.1649	0.4382	0.7420	0.065*	
H6D	1.2451	0.5721	0.7144	0.065*	
N6	1.0652 (2)	0.6024 (2)	0.82682 (15)	0.0482 (4)	
H6A	1.1435	0.5783	0.8766	0.058*	
H6B	0.9914	0.5539	0.8555	0.058*	
C7	0.7420 (3)	0.3504 (3)	0.91576 (18)	0.0494 (4)	
C8	0.5826 (4)	0.4304 (4)	0.8472 (3)	0.0722 (6)	
C9	0.5876 (3)	0.1740 (3)	0.4611 (2)	0.0526 (5)	
C10	0.4415 (4)	0.2533 (4)	0.5269 (3)	0.0749 (6)	
O1	0.8137 (3)	0.4407 (2)	0.92515 (17)	0.0722 (5)	
O2	0.7776 (3)	0.2083 (2)	0.95315 (17)	0.0721 (5)	
O3	0.7270 (2)	0.1248 (2)	0.50104 (16)	0.0693 (5)	
O4	0.5502 (3)	0.1706 (3)	0.37227 (17)	0.0780 (6)	

## supplementary materials

---

F1A	0.495 (2)	0.5750 (14)	0.8639 (12)	0.087 (2)	0.55 (3)
F2A	0.4769 (16)	0.3531 (13)	0.8668 (13)	0.089 (3)	0.55 (3)
F3A	0.6064 (18)	0.4591 (14)	0.7404 (6)	0.093 (3)	0.55 (3)
F1B	0.471 (2)	0.537 (2)	0.8865 (14)	0.083 (2)	0.45 (3)
F2B	0.5182 (17)	0.3267 (11)	0.8290 (14)	0.088 (2)	0.45 (3)
F3B	0.6454 (13)	0.4960 (16)	0.7584 (10)	0.084 (3)	0.45 (3)
F4A	0.4601 (19)	0.166 (3)	0.6294 (12)	0.095 (4)	0.31 (2)
F5A	0.2886 (15)	0.281 (3)	0.4998 (12)	0.090 (4)	0.31 (2)
F6A	0.460 (3)	0.3837 (19)	0.529 (2)	0.110 (5)	0.31 (2)
F4B	0.4729 (10)	0.2254 (9)	0.6304 (5)	0.0850 (12)	0.69 (2)
F5B	0.3116 (8)	0.2068 (10)	0.5228 (6)	0.0872 (13)	0.69 (2)
F6B	0.3834 (11)	0.4109 (4)	0.4916 (4)	0.0858 (16)	0.69 (2)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.03390 (11)	0.03383 (12)	0.03075 (11)	-0.01198 (8)	-0.00040 (7)	-0.00858 (8)
N1	0.0453 (9)	0.0514 (10)	0.0380 (6)	-0.0172 (8)	0.0030 (6)	-0.0134 (6)
C1	0.0495 (12)	0.0522 (12)	0.0418 (11)	-0.0159 (10)	0.0017 (9)	-0.0102 (9)
C2	0.0459 (11)	0.0535 (12)	0.0534 (12)	-0.0178 (10)	-0.0004 (9)	-0.0151 (10)
N2	0.0431 (7)	0.0582 (10)	0.0527 (10)	-0.0214 (8)	0.0005 (7)	-0.0173 (9)
N3	0.0516 (9)	0.0419 (7)	0.0458 (9)	-0.0141 (6)	-0.0022 (7)	-0.0090 (6)
C3	0.0558 (13)	0.0551 (13)	0.0501 (12)	-0.0207 (11)	-0.0030 (10)	-0.0082 (10)
C4	0.0567 (13)	0.0513 (12)	0.0484 (12)	-0.0217 (10)	-0.0007 (10)	-0.0126 (10)
N4	0.0435 (7)	0.0539 (10)	0.0436 (9)	-0.0214 (7)	0.0008 (6)	-0.0122 (8)
N5	0.0497 (9)	0.0504 (9)	0.0371 (7)	-0.0169 (8)	-0.0005 (6)	-0.0130 (7)
C5	0.0562 (13)	0.0536 (13)	0.0544 (13)	-0.0187 (10)	-0.0003 (10)	-0.0168 (11)
C6	0.0556 (13)	0.0511 (12)	0.0564 (13)	-0.0160 (10)	0.0011 (10)	-0.0178 (11)
N6	0.0492 (10)	0.0468 (10)	0.0468 (10)	-0.0151 (8)	-0.0046 (8)	-0.0085 (8)
C7	0.0437 (11)	0.0599 (10)	0.0441 (11)	-0.0142 (9)	0.0010 (8)	-0.0166 (9)
C8	0.0540 (14)	0.0761 (15)	0.0748 (13)	-0.0134 (9)	-0.0125 (11)	-0.0069 (13)
C9	0.0515 (10)	0.0506 (12)	0.0544 (11)	-0.0160 (10)	0.0046 (9)	-0.0141 (10)
C10	0.0646 (14)	0.0808 (12)	0.0766 (12)	-0.0185 (13)	0.0130 (13)	-0.0279 (14)
O1	0.0695 (12)	0.0765 (11)	0.0789 (13)	-0.0323 (10)	-0.0057 (10)	-0.0208 (10)
O2	0.0746 (12)	0.0622 (9)	0.0781 (13)	-0.0197 (9)	-0.0112 (10)	-0.0157 (9)
O3	0.0593 (10)	0.0788 (13)	0.0688 (11)	-0.0210 (9)	-0.0023 (8)	-0.0191 (10)
O4	0.0763 (13)	0.0897 (15)	0.0685 (11)	-0.0219 (11)	-0.0079 (9)	-0.0262 (11)
F1A	0.094 (5)	0.078 (3)	0.086 (5)	-0.026 (2)	-0.003 (3)	-0.018 (4)
F2A	0.085 (4)	0.089 (3)	0.097 (5)	-0.036 (3)	-0.012 (3)	-0.018 (3)
F3A	0.099 (5)	0.099 (4)	0.0789 (16)	-0.032 (3)	-0.003 (2)	-0.0216 (19)
F1B	0.077 (4)	0.089 (6)	0.085 (6)	-0.026 (4)	0.004 (3)	-0.025 (5)
F2B	0.086 (5)	0.093 (3)	0.089 (6)	-0.033 (3)	0.006 (4)	-0.029 (3)
F3B	0.078 (4)	0.102 (5)	0.075 (3)	-0.033 (3)	0.001 (2)	-0.022 (3)
F4A	0.070 (4)	0.113 (10)	0.093 (4)	-0.038 (6)	0.005 (3)	0.000 (6)
F5A	0.078 (3)	0.116 (11)	0.083 (5)	-0.038 (6)	0.005 (3)	-0.029 (6)
F6A	0.097 (9)	0.098 (5)	0.144 (12)	-0.056 (6)	0.016 (8)	-0.020 (7)
F4B	0.085 (3)	0.091 (3)	0.0800 (15)	-0.033 (2)	0.0010 (16)	-0.0179 (17)
F5B	0.080 (2)	0.091 (3)	0.090 (3)	-0.030 (3)	-0.0023 (18)	-0.020 (3)

F6B            0.082 (4)            0.0829 (12)            0.090 (2)            -0.0239 (16)            -0.0062 (19)            -0.0192 (13)

*Geometric parameters (Å, °)*

Cu1—N5	2.050 (2)	N5—H5A	0.9000
Cu1—N1	2.059 (2)	N5—H5B	0.9000
Cu1—N3	2.126 (2)	C5—C6	1.503 (3)
Cu1—N6	2.136 (2)	C5—H5C	0.9700
Cu1—N2	2.297 (2)	C5—H5D	0.9700
Cu1—N4	2.300 (2)	C6—N6	1.462 (3)
N1—C1	1.471 (3)	C6—H6C	0.9700
N1—H1A	0.9000	C6—H6D	0.9700
N1—H1B	0.9000	N6—H6A	0.9000
C1—C2	1.500 (3)	N6—H6B	0.9000
C1—H1C	0.9700	C7—O2	1.219 (3)
C1—H1D	0.9700	C7—O1	1.237 (3)
C2—N2	1.471 (3)	C7—C8	1.539 (4)
C2—H2C	0.9700	C8—F1B	1.291 (11)
C2—H2D	0.9700	C8—F3B	1.311 (7)
N2—H2A	0.9000	C8—F2A	1.312 (7)
N2—H2B	0.9000	C8—F3A	1.338 (7)
N3—C3	1.462 (3)	C8—F2B	1.345 (9)
N3—H3A	0.9000	C8—F1A	1.357 (9)
N3—H3B	0.9000	C9—O4	1.225 (3)
C3—C4	1.499 (3)	C9—O3	1.230 (3)
C3—H3C	0.9700	C9—C10	1.524 (4)
C3—H3D	0.9700	C10—F6A	1.286 (10)
C4—N4	1.466 (3)	C10—F5A	1.308 (11)
C4—H4C	0.9700	C10—F4B	1.322 (6)
C4—H4D	0.9700	C10—F5B	1.338 (5)
N4—H4A	0.9000	C10—F6B	1.338 (5)
N4—H4B	0.9000	C10—F4A	1.343 (11)
N5—C5	1.464 (3)		
N5—Cu1—N1	171.52 (7)	H5A—N5—H5B	108.3
N5—Cu1—N3	91.32 (8)	N5—C5—C6	107.7 (2)
N1—Cu1—N3	93.88 (8)	N5—C5—H5C	110.2
N5—Cu1—N6	81.82 (8)	C6—C5—H5C	110.2
N1—Cu1—N6	93.95 (8)	N5—C5—H5D	110.2
N3—Cu1—N6	169.22 (7)	C6—C5—H5D	110.2
N5—Cu1—N2	92.91 (8)	H5C—C5—H5D	108.5
N1—Cu1—N2	80.03 (8)	N6—C6—C5	108.1 (2)
N3—Cu1—N2	94.49 (8)	N6—C6—H6C	110.1
N6—Cu1—N2	94.15 (9)	C5—C6—H6C	110.1
N5—Cu1—N4	98.13 (8)	N6—C6—H6D	110.1
N1—Cu1—N4	89.38 (8)	C5—C6—H6D	110.1
N3—Cu1—N4	79.81 (8)	H6C—C6—H6D	108.4
N6—Cu1—N4	92.88 (8)	C6—N6—Cu1	107.29 (14)
N2—Cu1—N4	167.65 (7)	C6—N6—H6A	110.3
C1—N1—Cu1	110.81 (14)	Cu1—N6—H6A	110.3

## supplementary materials

---

C1—N1—H1A	109.5	C6—N6—H6B	110.3
Cu1—N1—H1A	109.5	Cu1—N6—H6B	110.3
C1—N1—H1B	109.5	H6A—N6—H6B	108.5
Cu1—N1—H1B	109.5	O2—C7—O1	129.9 (2)
H1A—N1—H1B	108.1	O2—C7—C8	115.2 (2)
N1—C1—C2	110.86 (18)	O1—C7—C8	114.9 (2)
N1—C1—H1C	109.5	F1B—C8—F3B	110.6 (7)
C2—C1—H1C	109.5	F1B—C8—F2A	86.2 (6)
N1—C1—H1D	109.5	F3B—C8—F2A	133.3 (5)
C2—C1—H1D	109.5	F1B—C8—F3A	118.9 (10)
H1C—C1—H1D	108.1	F3B—C8—F3A	28.3 (3)
N2—C2—C1	107.94 (19)	F2A—C8—F3A	105.2 (5)
N2—C2—H2C	110.1	F1B—C8—F2B	110.6 (7)
C1—C2—H2C	110.1	F3B—C8—F2B	110.6 (5)
N2—C2—H2D	110.1	F2A—C8—F2B	27.2 (3)
C1—C2—H2D	110.1	F3A—C8—F2B	82.7 (5)
H2C—C2—H2D	108.4	F1B—C8—F1A	21.0 (6)
C2—N2—Cu1	105.49 (14)	F3B—C8—F1A	90.5 (6)
C2—N2—H2A	110.6	F2A—C8—F1A	104.9 (5)
Cu1—N2—H2A	110.6	F3A—C8—F1A	103.9 (6)
C2—N2—H2B	110.6	F2B—C8—F1A	126.1 (7)
Cu1—N2—H2B	110.6	F1B—C8—C7	112.6 (9)
H2A—N2—H2B	108.8	F3B—C8—C7	98.8 (7)
C3—N3—Cu1	109.77 (15)	F2A—C8—C7	114.9 (4)
C3—N3—H3A	109.7	F3A—C8—C7	115.3 (7)
Cu1—N3—H3A	109.7	F2B—C8—C7	113.1 (5)
C3—N3—H3B	109.7	F1A—C8—C7	111.6 (7)
Cu1—N3—H3B	109.7	O4—C9—O3	128.0 (3)
H3A—N3—H3B	108.2	O4—C9—C10	114.6 (3)
N3—C3—C4	110.8 (2)	O3—C9—C10	117.4 (3)
N3—C3—H3C	109.5	F6A—C10—F5A	109.5 (6)
C4—C3—H3C	109.5	F6A—C10—F4B	79.6 (10)
N3—C3—H3D	109.5	F5A—C10—F4B	117.5 (7)
C4—C3—H3D	109.5	F6A—C10—F5B	135.3 (9)
H3C—C3—H3D	108.1	F5A—C10—F5B	28.1 (8)
N4—C4—C3	110.08 (19)	F4B—C10—F5B	105.9 (4)
N4—C4—H4C	109.6	F6A—C10—F6B	34.5 (11)
C3—C4—H4C	109.6	F5A—C10—F6B	77.3 (8)
N4—C4—H4D	109.6	F4B—C10—F6B	105.2 (4)
C3—C4—H4D	109.6	F5B—C10—F6B	105.2 (3)
H4C—C4—H4D	108.2	F6A—C10—F4A	105.5 (7)
C4—N4—Cu1	105.13 (14)	F5A—C10—F4A	106.4 (8)
C4—N4—H4A	110.7	F4B—C10—F4A	26.0 (9)
Cu1—N4—H4A	110.7	F5B—C10—F4A	86.2 (7)
C4—N4—H4B	110.7	F6B—C10—F4A	128.0 (12)
Cu1—N4—H4B	110.7	F6A—C10—C9	105.3 (6)
H4A—N4—H4B	108.8	F5A—C10—C9	120.7 (7)
C5—N5—Cu1	109.29 (14)	F4B—C10—C9	114.9 (4)
C5—N5—H5A	109.8	F5B—C10—C9	111.5 (4)



Cu1—N5—H5A	109.8	F6B—C10—C9	113.3 (3)
C5—N5—H5B	109.8	F4A—C10—C9	108.5 (9)
Cu1—N5—H5B	109.8		

*Hydrogen-bond geometry (Å, °)*

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N2—H2B···F1A	0.90	2.54	3.248 (17)	136
N2—H2B···F3B	0.90	2.55	3.41 (2)	161
N6—H6B···O1	0.90	2.15	3.048 (3)	179
N1—H1A···O2 <sup>i</sup>	0.90	2.35	3.141 (3)	147
N1—H1A···O1 <sup>i</sup>	0.90	2.53	3.377 (4)	156
N4—H4A···O2 <sup>i</sup>	0.90	2.34	3.175 (3)	154
N6—H6A···O1 <sup>i</sup>	0.90	2.56	3.326 (3)	143
N1—H1B···O2 <sup>ii</sup>	0.90	2.11	3.002 (3)	173
N2—H2A···F4A <sup>ii</sup>	0.90	2.47	3.267 (15)	148
N3—H3A···O3 <sup>ii</sup>	0.90	2.09	2.958 (3)	162
N5—H5A···O3 <sup>ii</sup>	0.90	2.36	3.126 (3)	143
N2—H2A···O4 <sup>iii</sup>	0.90	2.41	3.037 (3)	127
N4—H4B···O4 <sup>iv</sup>	0.90	2.11	2.997 (3)	168
N5—H5B···O3 <sup>iv</sup>	0.90	2.13	3.013 (3)	167

Symmetry codes: (i)  $-x+2, -y+1, -z+2$ ; (ii)  $x, y+1, z$ ; (iii)  $-x+1, -y+1, -z+1$ ; (iv)  $-x+2, -y+1, -z+1$ .

Fig. 1

