

cis-Bis(2-sulfidopyridine N-oxide)-platinum(II)

B. Ravindran Durai Nayagam,^{a*} Samuel Robinson Jebas^b and Dieter Schollmeyer^c

^aDepartment of Chemistry, Popes College, Sawyerpuram, Tamilnadu, India,

^bDepartment of Physics, Karunya University, Coimbatore 641 114, India, and

^cInstitut für Organische Chemie, Universität Mainz, Duesbergweg 10-14, 55099

Mainz, Germany

Correspondence e-mail: b_ravidurai@yahoo.com


Received 13 December 2007; accepted 18 January 2008

Key indicators: single-crystal X-ray study; $T = 193$ K; mean $\sigma(\text{C}-\text{C}) = 0.017$ Å; R factor = 0.055; wR factor = 0.114; data-to-parameter ratio = 18.0.

In the crystal structure of the title complex, $[\text{Pt}(\text{C}_5\text{H}_4\text{NOS})_2]$, the Pt atom is coordinated by two O atoms and two S atoms in a *cis* configuration, forming a distorted square-planar coordination geometry. The molecule exhibits pseudo- C_{2v} symmetry and is essentially planar, with a maximum deviation from planarity of 0.0124 (2) Å. The dihedral angle between the two pyridine rings is 5.85 (2)°.

Related literature

For related literature, see: Bovin *et al.* (1992); Chen *et al.* (1991); Dyksterhouse *et al.* (2000); Katsuyuki *et al.* (1991); Leonard *et al.* (1955); Lobana & Bhatia (1989); Lydon *et al.* (1982); Ohms *et al.* (1982); Symons & West (1985); Zhou *et al.* (2005); Shi *et al.* (1997).



Experimental

Crystal data

$[\text{Pt}(\text{C}_5\text{H}_4\text{NOS})_2]$

$M_r = 447.39$

Monoclinic, $P2_1/n$

$a = 6.9832$ (3) Å

$b = 22.3897$ (11) Å

$c = 8.1495$ (4) Å

$\beta = 113.572$ (3)°

$V = 1167.87$ (10) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 12.36$ mm⁻¹

$T = 193$ (2) K

0.27 × 0.22 × 0.08 mm

Data collection

Bruker Kappa APEXII

diffractometer

Absorption correction: multi-scan

(*SADABS*; Blessing, 1995, 1997)

$T_{\min} = 0.051$, $T_{\max} = 0.372$

25638 measured reflections

2778 independent reflections

2610 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.068$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$

$wR(F^2) = 0.114$

$S = 1.32$

2778 reflections

154 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 1.73$ e Å⁻³

$\Delta\rho_{\text{min}} = -3.87$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Pt1—O15	2.045 (7)	Pt1—S16	2.234 (3)
Pt1—O7	2.052 (7)	Pt1—S8	2.239 (3)
O15—Pt1—O7	90.0 (3)	O15—Pt1—S8	176.4 (2)
O15—Pt1—S16	86.4 (2)	O7—Pt1—S8	86.4 (2)
O7—Pt1—S16	176.3 (2)	S16—Pt1—S8	97.13 (10)

Data collection: *APEX2* (Bruker, 2006); cell refinement: *APEX2*; data reduction: *APEX2*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP2* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2232).

References

- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
 Blessing, R. H. (1997). *J. Appl. Cryst.* **30**, 421–426.
 Bovin, D. H. R., Crepon, E. & Zard, S. Z. (1992). *Bull. Soc. Chim. Fr.* **129**, 145–150.
 Bruker (2006). *APEX2*. Version 2.0. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chen, X., Hu, Y., Wu, D., Weng, L. & Kang, B. (1991). *Polyhedron*, **10**, 2651–2657.
 Dyksterhouse, R. M., Howell, B. A. & Squattrito, P. J. (2000). *Acta Cryst.* **C56**, 64–66.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Katsuyuki, N., Carter, B. J., Xu, J. & Hetch, S. M. (1991). *J. Am. Chem. Soc.* **113**, 5099–5100.
 Leonard, F., Barklay, F. A., Brown, E. V., Anderson, F. E. & Green, D. M. (1955). *Antibiot. Chemother.* pp. 261–264.
 Lobana, T. S. & Bhatia, P. K. (1989). *J. Sci. Ind. Res.* **48**, 394–401.
 Lydon, J. D., Elder, R. C. & Deutch, E. (1982). *Inorg. Chem.* **21**, 3168–3197.
 Ohms, U., Guth, H., Kutoglu, A. & Scheringer, C. (1982). *Acta Cryst.* **B38**, 831–834.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Shi, J. C., Wen, T. B., Zheng, Y., Zhong, S. J., Wu, D. X., Liu, Q. T., Kang, B. S., Wu, B. M. & Mak, T. C. W. (1997). *Polyhedron*, **16**, 369–375.
 Symons, M. C. R. & West, D.-X. (1985). *J. Chem. Soc. Dalton Trans.* pp. 379–381.
 Zhou, J.-H., Li, Y.-Z., Lui, Z. & Chen, X.-T. (2005). *Acta Cryst.* **E61**, m195–m197.

supplementary materials

Acta Cryst. (2008). E64, m425 [doi:10.1107/S1600536808002006]

***cis*-Bis(2-sulfidopyridine *N*-oxide)platinum(II)**

B. Ravindran Durai Nayagam, S. R. Jebas and D. Schollmeyer

Comment

N-oxides and their derivatives show a broad spectrum of biological activity, such as antifungal, antibacterial, antimicrobial and antibiotic activities (Lobana & Bhatia, 1989; Symons *et al.*, 1985). These compounds are also found to be involved in DNA strand scission under physiological conditions (Katsuyuki *et al.*, 1991; Bovin *et al.*, 1992). Pyridine N-oxides bearing a sulfur group at the 2-position display significant antimicrobial activity (Leonard *et al.*, 1955). The crystal structure of the title compound in the *trans* configuration has already been reported (Zhou *et al.*, 2005). The crystal structures of *cis*-bis(2-mercaptopyridine N-oxide)palladium and *cis*-bis(2-mercaptopyridine N-oxide)nickel have already been reported (Shi *et al.*, 1997; Chen *et al.*, 1991).

In the title compound, the Pt is coordinated by two O atoms and two S atoms from two mercaptopyridine N-oxide ligands in a *cis* configuration. The coordination geometry is distorted square-planar. The whole molecule exhibits pseudo-C_{2v} symmetry, with a pseudo-C₂ axis passing through the platinum atom. The average Pt—O and Pt—S distances of 2.048 (7) and 2.236 (3) Å of the title compound are comparable with the values reported in the literature (Dyksterhouse *et al.*, 2000). The mean Pt—S bond length (2.236 (3) Å,) is shorter while the mean Pt—O bond length (2.048 (7) Å) is longer than those reported for the *trans* isomer (Pt—S 2.270 Å; Pt—O 2.007 Å) (Zhou *et al.*, 2005). The C—S bond distances [1.725 (11) and 1.730 (11) Å] are slightly longer than those reported for the uncoordinated thione molecule [1.692 (2)–1.698 (2) Å; Ohms *et al.*, 1982]. The mean C—S bond length (1.727 Å) shows partial double bond character; it is shorter than the normal covalent bond distance of 1.81 (2) Å, for a C—S single bond (Lydon *et al.*, 1982). This further results in the fact that complexes with mercaptopyridine oxide (mpo) ligands have little ability to bridge another metal ion through a sulfur atom to form polynuclear complexes (Shi *et al.*, 1997). The entire molecule is essentially planar, with a maximum deviation from planarity of 0.0124 (2) Å for atom C3. The dihedral angle between the two pyridine rings is 5.85 (2)°.

Experimental

By heating a mixture of dichlorido(ethane1,2-diamine)platinum(II), Pt(en)Cl₂ (0.326 g, 1 mmol) and 1-hydroxypyridine-2-thione sodium salt (0.298 g, 2 mmol) in 20 ml of water at 338 K with magnetic stirring for 1 h, a red-orange compound was obtained. This was filtered and dried (0.32 g, 80% yield). The compound was dissolved in methanol and allowed to undergo slow evaporation. Fine red crystals were obtained after a week.

Refinement

H atoms were placed in calculated positions [C—H = 0.95 Å] and refined in the riding model approximation; $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The highest residual electron density peak is 1.724 Å from C13 and the deepest hole lies 1.688 Å from N1.

Figures



Fig. 1. View of the title compound, with displacement ellipsoids drawn at the 50% probability level.

cis-Bis(2-sulfidopyridine N-oxide)platinum(II)

Crystal data

[Pt(C ₅ H ₄ NOS) ₂]	$F_{000} = 832$
$M_r = 447.39$	$D_x = 2.545 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: $-P\ 2_1n$	$\lambda = 0.71069 \text{ \AA}$
$a = 6.9832 (3) \text{ \AA}$	Cell parameters from 9973 reflections
$b = 22.3897 (11) \text{ \AA}$	$\theta = 2.8\text{--}28^\circ$
$c = 8.1495 (4) \text{ \AA}$	$\mu = 12.36 \text{ mm}^{-1}$
$\beta = 113.572 (3)^\circ$	$T = 193 (2) \text{ K}$
$V = 1167.87 (10) \text{ \AA}^3$	Block, orange
$Z = 4$	$0.27 \times 0.22 \times 0.08 \text{ mm}$

Data collection

Bruker APEXII Kappa-CCD diffractometer	$R_{\text{int}} = 0.068$
CCD scan	$\theta_{\text{max}} = 27.9^\circ$
Absorption correction: multi-scan (SADABS; Blessing, 1995, 1997)	$\theta_{\text{min}} = 1.8^\circ$
$T_{\text{min}} = 0.051$, $T_{\text{max}} = 0.372$	$h = -9 \rightarrow 9$
25638 measured reflections	$k = -29 \rightarrow 29$
2778 independent reflections	$l = -10 \rightarrow 10$
2610 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.0153P)^2 + 17.8973P]$
$R[F^2 > 2\sigma(F^2)] = 0.054$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.114$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.32$	$\Delta\rho_{\text{max}} = 1.73 \text{ e \AA}^{-3}$
2778 reflections	$\Delta\rho_{\text{min}} = -3.86 \text{ e \AA}^{-3}$
154 parameters	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Pt1	0.75198 (6)	0.117494 (18)	0.60862 (6)	0.03763 (14)
N1	0.5870 (13)	0.1570 (4)	0.2386 (12)	0.0373 (18)
C2	0.5399 (16)	0.1976 (5)	0.1045 (15)	0.042 (2)
H2	0.5774	0.2384	0.1308	0.05*
C3	0.4376 (18)	0.1791 (6)	-0.0694 (17)	0.050 (3)
H3	0.4097	0.2066	-0.165	0.059*
C4	0.3746 (18)	0.1195 (6)	-0.1053 (16)	0.048 (3)
H4	0.3007	0.1064	-0.2251	0.058*
C5	0.4207 (16)	0.0803 (5)	0.0340 (15)	0.042 (2)
H5	0.3772	0.0399	0.0102	0.05*
C6	0.5318 (17)	0.0990 (5)	0.2124 (15)	0.039 (2)
O7	0.6952 (11)	0.1788 (3)	0.4073 (10)	0.0380 (15)
S8	0.5946 (4)	0.05080 (12)	0.3916 (4)	0.0413 (6)
N9	0.9639 (13)	0.1639 (4)	0.9685 (12)	0.0378 (19)
C10	1.0646 (17)	0.2052 (5)	1.0962 (16)	0.042 (2)
H10	1.0885	0.2441	1.0617	0.051*
C11	1.1311 (17)	0.1915 (5)	1.2724 (16)	0.044 (2)
H11	1.1986	0.2208	1.3612	0.053*
C12	1.0994 (17)	0.1342 (5)	1.3215 (15)	0.043 (2)
H12	1.147	0.1239	1.4445	0.051*
C13	0.9994 (17)	0.0921 (5)	1.1923 (15)	0.042 (2)
H13	0.9763	0.053	1.2263	0.051*
C14	0.9315 (16)	0.1069 (5)	1.0105 (14)	0.038 (2)
O15	0.8948 (11)	0.1822 (3)	0.7947 (9)	0.0381 (16)
S16	0.8174 (4)	0.05572 (12)	0.8401 (4)	0.0394 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pt1	0.0364 (2)	0.0382 (2)	0.0391 (2)	0.00055 (17)	0.01591 (17)	0.00058 (17)
N1	0.032 (4)	0.039 (5)	0.042 (5)	-0.002 (3)	0.017 (4)	-0.001 (4)
C2	0.033 (5)	0.049 (6)	0.039 (6)	0.006 (4)	0.009 (4)	0.011 (5)
C3	0.043 (6)	0.063 (7)	0.046 (7)	0.003 (5)	0.021 (5)	0.013 (6)
C4	0.040 (6)	0.062 (7)	0.041 (6)	0.001 (5)	0.014 (5)	0.000 (5)
C5	0.031 (5)	0.056 (7)	0.041 (6)	-0.003 (5)	0.016 (4)	-0.003 (5)
C6	0.037 (5)	0.041 (5)	0.045 (6)	0.001 (4)	0.023 (5)	0.000 (4)
O7	0.035 (4)	0.043 (4)	0.036 (4)	-0.003 (3)	0.016 (3)	-0.002 (3)

supplementary materials

S8	0.0440 (14)	0.0404 (13)	0.0391 (14)	-0.0015 (11)	0.0164 (12)	0.0009 (11)
N9	0.032 (4)	0.041 (5)	0.044 (5)	0.002 (3)	0.020 (4)	0.000 (4)
C10	0.039 (6)	0.039 (5)	0.050 (7)	-0.001 (4)	0.019 (5)	-0.003 (5)
C11	0.037 (6)	0.050 (6)	0.044 (6)	0.000 (5)	0.016 (5)	-0.009 (5)
C12	0.039 (6)	0.052 (6)	0.038 (6)	0.003 (5)	0.016 (5)	-0.001 (5)
C13	0.042 (6)	0.047 (6)	0.043 (6)	0.004 (5)	0.024 (5)	0.007 (5)
C14	0.032 (5)	0.044 (6)	0.040 (6)	0.001 (4)	0.017 (4)	-0.001 (4)
O15	0.038 (4)	0.042 (4)	0.034 (4)	0.000 (3)	0.014 (3)	0.005 (3)
S16	0.0399 (14)	0.0399 (13)	0.0381 (14)	-0.0016 (10)	0.0154 (11)	-0.0001 (10)

Geometric parameters (Å, °)

Pt1—O15	2.045 (7)	N9—C14	1.364 (13)
Pt1—O7	2.052 (7)	C10—C11	1.356 (16)
Pt1—S16	2.234 (3)	C11—C12	1.389 (16)
Pt1—S8	2.239 (3)	C12—C13	1.376 (16)
N1—C6	1.348 (13)	C13—C14	1.403 (15)
N1—C2	1.357 (13)	C14—S16	1.730 (11)
N1—O7	1.367 (11)	C2—H2	0.9500
C2—C3	1.372 (16)	C3—H3	0.9500
C3—C4	1.399 (17)	C4—H4	0.9500
C4—C5	1.367 (16)	C5—H5	0.9500
C5—C6	1.410 (15)	C10—H10	0.9500
C6—S8	1.725 (11)	C11—H11	0.9500
N9—C10	1.359 (14)	C12—H12	0.9500
N9—O15	1.363 (11)	C13—H13	0.9500
O15—Pt1—O7	90.0 (3)	C10—N9—C14	122.1 (10)
O15—Pt1—S16	86.4 (2)	O15—N9—C14	121.0 (9)
O7—Pt1—S16	176.3 (2)	C11—C10—N9	120.6 (10)
O15—Pt1—S8	176.4 (2)	C10—C11—C12	119.2 (11)
O7—Pt1—S8	86.4 (2)	C13—C12—C11	120.2 (11)
S16—Pt1—S8	97.13 (10)	C12—C13—C14	120.0 (10)
C6—N1—C2	123.8 (10)	N9—C14—C13	117.8 (10)
C6—N1—O7	120.7 (9)	N9—C14—S16	119.2 (8)
C2—N1—O7	115.4 (9)	C13—C14—S16	122.9 (9)
N1—C2—C3	119.2 (11)	N9—O15—Pt1	115.1 (6)
C2—C3—C4	119.6 (11)	C14—S16—Pt1	98.1 (4)
C5—C4—C3	119.3 (11)	N1—C2—H2	120.0
C4—C5—C6	121.0 (11)	C3—C2—H2	120.0
N1—C6—C5	117.0 (10)	C2—C3—H3	120.0
N1—C6—S8	120.6 (8)	C4—C3—H3	120.0
C5—C6—S8	122.4 (9)	C3—C4—H4	120.0
N1—O7—Pt1	114.7 (6)	C5—C4—H4	120.0
C6—S8—Pt1	97.5 (4)	C4—C5—H5	119.0
C10—N9—O15	116.8 (9)	C6—C5—H5	120.0

Fig. 1

