

## N-(2-Chloro-2-nitro-1-phenylpropyl)-4-methylbenzenesulfonamide

Sanjun Zhi, Tengfei Li, Guanghui An and Yi Pan\*

School of Chemistry and Chemical Engineering, State Key Laboratory of Coordination Chemistry, Nanjing University, Nanjing 210093, People's Republic of China  
Correspondence e-mail: jchyzhi2003@126.com


Received 29 November 2007; accepted 19 December 2007

Key indicators: single-crystal X-ray study;  $T = 291$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.056;  $wR$  factor = 0.125; data-to-parameter ratio = 15.3.

In the title compound,  $\text{C}_{16}\text{H}_{17}\text{ClN}_2\text{O}_4\text{S}$ , the dihedral angle between the phenyl and benzene rings is  $19.4(2)^\circ$ . The crystal packing is stabilized by intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds, as well as by intra- and intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For general background, see Kemp (1991); Qui & Silverman (2000); Orlek & Stemp (1991), Han *et al.* (2007); Li *et al.* (2007). For bond-length data, see: Allen *et al.* (1987).



### Experimental

#### Crystal data

$\text{C}_{16}\text{H}_{17}\text{ClN}_2\text{O}_4\text{S}$   
 $M_r = 368.83$   
Orthorhombic,  $Pbca$   
 $a = 7.8254(8)$  Å  
 $b = 19.610(2)$  Å  
 $c = 22.533(3)$  Å

$V = 3457.8(7)$  Å<sup>3</sup>  
 $Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 0.36$  mm<sup>-1</sup>  
 $T = 291(2)$  K  
 $0.30 \times 0.26 \times 0.24$  mm

#### Data collection

Bruker SMART APEX CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2000)  
 $T_{\min} = 0.901$ ,  $T_{\max} = 0.921$   
17527 measured reflections  
3396 independent reflections  
2467 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.070$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.125$   
 $S = 1.01$   
3396 reflections  
222 parameters  
H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.28$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.26$  e Å<sup>-3</sup>

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H3}\cdots\text{O1}$	0.98	2.46	2.933 (4)	109
$\text{C3}-\text{H3}\cdots\text{O3}$	0.98	2.42	2.779 (4)	101
$\text{C1}-\text{H1A}\cdots\text{O2}^i$	0.96	2.52	3.369 (4)	147
$\text{C1}-\text{H1B}\cdots\text{O1}^{ii}$	0.96	2.58	3.310 (4)	133
$\text{N2}-\text{H2A}\cdots\text{O2}^i$	0.89 (2)	2.32 (3)	3.141 (3)	153

Symmetry codes: (i)  $x + \frac{1}{2}, y, -z + \frac{3}{2}$ ; (ii)  $x + 1, y, z$ .

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Bruker, 2000); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2003).

This work was supported by the 863 High Technology Program (to YP). The research funds for YP from the Qing-Lan Program of Jiangsu Province and the Kua-Shi-Ji Program of the Education Ministry of China are also acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2518).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.  
Bruker (2000). SMART (Version 5.625), SAINT (Version 6.01), SHELXTL (Version 6.10) and SADABS (Version 2.03). Bruker AXS Inc., Madison, Wisconsin, USA.  
Han, J., Zhi, S., Wang, L., Pan, Y. & Li, G. (2007). *Eur. J. Org. Chem.* pp. 1332–1337.  
Kemp, J. E. G. (1991). *Comprehensive Organic Synthesis*, Vol. 3, edited by B. M. Trost & I. Fleming, pp. 469–513. Oxford: Pergamon Press.  
Li, G., Saibabu, K. S. R. S. & Timmons, C. (2007). *Eur. J. Org. Chem.* pp. 2745–2758.  
Orlek, B. S. & Stemp, G. (1991). *Tetrahedron Lett.* **32**, 4045–4048.  
Qui, J. & Silverman, R. B. (2000). *J. Med. Chem.* **43**, 706–720.  
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

**supplementary materials**

*Acta Cryst.* (2008). E64, o357 [ doi:10.1107/S1600536807067712 ]

## ***N*-(2-Chloro-2-nitro-1-phenylpropyl)-4-methylbenzenesulfonamide**

**S. Zhi, T. Li, G. An and Y. Pan**

### **Comment**

Since the vicinal haloamines are important building blocks in organic and medicinal chemistry, many attentions are attracted to the aminohalogenation reactions of functionalized alkenes (Kemp, 1991; Qui & Silverman, 2000). In recent years, many new aminohalogenation processes of several kinds of functionalized alkenes have been developed (Han *et al.*, 2007), with the different nitrogen/halogen sources in the presence of metallic catalysts (Li *et al.*, 2007). However, the aminohalogenation reactions of 2-nitro-propenyl benzene was not been well documented. Recently, we synthesized the title compound (I) from 2-nitro-propenyl benzene. As part of this study, we have undertaken the X-ray crystallographic analysis of (I) in order to elucidate the conformation and configuration of this product.

The bond lengths and angles in (I) are in good agree with expected values (Allen *et al.*, 1987). The dihedral angle between the phenyl and benzene rings is 19.4 (2)°. The packing is stabilized by intermolecular N—H···O as well as intra and intermolecular C—H···O interactions in the crystal structure (Table 1).

### **Experimental**

*N*-cChlorosuccinimide (400 mg, 3.0 mmol) was added into a solution of MnSO<sub>4</sub> (30.2 mg, 0.20 mmol), 1-benzyl-2-nitro-propene (163 mg, 1 mmol), toluenesulfonamide (513 mg, 3 mmol) and 4 Å molecular sieves (500 mg) in CH<sub>2</sub>Cl<sub>2</sub> (5.0 ml) with nitrogen atmosphere. The resulting mixture was stirred at room temperature for 48 h. Reaction was quenched with saturated aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution. The solid precipitates were filtered off and washed with ethyl acetate (3 × 10 ml). The organic solution was concentrated and then purified *via* flash chromatography (ethyl acetate/ hexane, 1:4, *v/v*) provide the title compound (I) as white solid (276 mg) in yield of 75%. A colourless crystal of (I) for X-ray analysis was obtained by slow evaporation from ethyl acetate solution system.

### **Refinement**

The H atom bonded to N was located in a difference map and refined with restraint of N—H = 0.89 (3) Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ . Other H atoms were geometrically placed and were treated as riding, with C—H distances of 0.93, 0.96 and 0.98 Å for aromatic, methyl and methine H atoms, respectively, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{aromatic and methyne C})$  or  $1.5U_{\text{eq}}(\text{methyl C})$ .

Figures




Fig. 1. The molecular structure of (I), with atom labels and 30% probability displacement ellipsoids for non-H atoms.

***N*-(2-Chloro-2-nitro-1-phenylpropyl)-4-methylbenzenesulfonamide**

*Crystal data*

C<sub>16</sub>H<sub>17</sub>ClN<sub>2</sub>O<sub>4</sub>S

*M<sub>r</sub>* = 368.83

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

*a* = 7.8254 (8) Å

*b* = 19.610 (2) Å

*c* = 22.533 (3) Å

*V* = 3457.8 (7) Å<sup>3</sup>

*Z* = 8

*F*<sub>000</sub> = 1536

*D<sub>x</sub>* = 1.417 Mg m<sup>-3</sup>

Melting point: 423.2 K

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 7468 reflections

θ = 2.3–27.9°

μ = 0.36 mm<sup>-1</sup>

*T* = 291 (2) K

Block, colourless

0.30 × 0.26 × 0.24 mm

*Data collection*

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: sealed tube

Monochromator: graphite

*T* = 291(2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2000)

*T*<sub>min</sub> = 0.901, *T*<sub>max</sub> = 0.921

17527 measured reflections

3396 independent reflections

2467 reflections with *I* > 2σ(*I*)

*R*<sub>int</sub> = 0.070

θ<sub>max</sub> = 26.0°

θ<sub>min</sub> = 2.1°

*h* = -9→8

*k* = -24→24

*l* = -16→27

*Refinement*

Refinement on *F*<sup>2</sup>

Least-squares matrix: full

*R*[*F*<sup>2</sup> > 2σ(*F*<sup>2</sup>)] = 0.056

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$wR(F^2) = 0.125$	$w = 1/[\sigma^2(F_o^2) + (0.06P)^2 + 0.58P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
3396 reflections	$(\Delta/\sigma)_{\max} < 0.001$
222 parameters	$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7916 (4)	0.10320 (17)	0.61740 (13)	0.0374 (7)
H1A	0.7709	0.1030	0.6594	0.056*
H1B	0.8606	0.1420	0.6072	0.056*
H1C	0.8503	0.0621	0.6064	0.056*
C2	0.6185 (4)	0.10723 (16)	0.58376 (13)	0.0333 (7)
C3	0.5059 (4)	0.16923 (14)	0.59888 (13)	0.0292 (6)
H3	0.3995	0.1645	0.5763	0.035*
C4	0.5853 (3)	0.23727 (15)	0.58138 (13)	0.0293 (6)
C5	0.5405 (4)	0.26704 (17)	0.52790 (13)	0.0362 (7)
H5	0.4651	0.2445	0.5028	0.043*
C6	0.6055 (4)	0.32954 (17)	0.51103 (14)	0.0415 (8)
H6	0.5765	0.3484	0.4745	0.050*
C7	0.7156 (4)	0.36407 (18)	0.54968 (15)	0.0437 (8)
H7	0.7570	0.4070	0.5396	0.052*
C8	0.7623 (4)	0.33466 (18)	0.60226 (15)	0.0444 (8)
H8	0.8381	0.3572	0.6273	0.053*
C9	0.6972 (4)	0.27113 (17)	0.61859 (14)	0.0414 (8)
H9	0.7290	0.2516	0.6545	0.050*
C10	0.3185 (3)	0.28495 (15)	0.69070 (12)	0.0314 (6)
C11	0.2653 (4)	0.32776 (18)	0.64579 (14)	0.0402 (7)
H11	0.2085	0.3099	0.6131	0.048*
C12	0.2959 (5)	0.39679 (18)	0.64904 (15)	0.0465 (8)
H12	0.2601	0.4252	0.6184	0.056*
C13	0.3804 (4)	0.42454 (18)	0.69822 (14)	0.0419 (8)
C14	0.4350 (4)	0.38054 (17)	0.74268 (15)	0.0423 (8)

## supplementary materials

---

H14	0.4939	0.3980	0.7751	0.051*
C15	0.4037 (4)	0.31171 (16)	0.73966 (13)	0.0349 (7)
H15	0.4392	0.2831	0.7702	0.042*
C16	0.4064 (4)	0.49975 (18)	0.70326 (17)	0.0491 (9)
H16A	0.3241	0.5184	0.7304	0.074*
H16B	0.3922	0.5204	0.6650	0.074*
H16C	0.5196	0.5087	0.7177	0.074*
C11	0.65350 (9)	0.10276 (4)	0.50665 (3)	0.03580 (19)
N1	0.5083 (4)	0.04372 (13)	0.59707 (12)	0.0407 (6)
N2	0.4620 (3)	0.16217 (13)	0.66251 (11)	0.0334 (6)
H2A	0.547 (4)	0.1794 (17)	0.6841 (15)	0.040*
O1	0.1557 (3)	0.18367 (12)	0.64347 (10)	0.0439 (6)
O2	0.2635 (3)	0.17162 (12)	0.74582 (9)	0.0432 (6)
O3	0.3667 (3)	0.04155 (12)	0.57591 (12)	0.0547 (7)
O4	0.5674 (3)	-0.00057 (13)	0.62846 (12)	0.0595 (7)
S1	0.28433 (9)	0.19677 (4)	0.68662 (3)	0.03081 (19)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0406 (16)	0.0483 (18)	0.0233 (14)	0.0236 (14)	-0.0020 (12)	-0.0033 (14)
C2	0.0365 (16)	0.0373 (16)	0.0261 (14)	0.0020 (13)	0.0056 (12)	-0.0015 (13)
C3	0.0296 (15)	0.0309 (14)	0.0270 (14)	-0.0002 (12)	0.0055 (11)	-0.0002 (12)
C4	0.0267 (14)	0.0322 (15)	0.0290 (15)	-0.0027 (11)	0.0033 (12)	-0.0024 (12)
C5	0.0344 (15)	0.0448 (17)	0.0295 (15)	-0.0005 (14)	-0.0042 (12)	0.0001 (13)
C6	0.056 (2)	0.0446 (19)	0.0242 (16)	0.0022 (15)	0.0026 (14)	0.0134 (14)
C7	0.0534 (19)	0.0383 (18)	0.0393 (19)	-0.0083 (15)	0.0079 (15)	0.0100 (15)
C8	0.0513 (19)	0.0471 (19)	0.0349 (18)	-0.0214 (16)	0.0037 (15)	0.0031 (15)
C9	0.0496 (18)	0.0467 (19)	0.0279 (15)	-0.0154 (15)	-0.0060 (14)	0.0084 (14)
C10	0.0265 (14)	0.0402 (17)	0.0275 (15)	0.0008 (12)	0.0057 (11)	0.0011 (12)
C11	0.0452 (17)	0.0461 (19)	0.0293 (16)	0.0124 (14)	-0.0042 (14)	0.0062 (14)
C12	0.059 (2)	0.0433 (19)	0.0368 (18)	0.0120 (16)	-0.0016 (16)	0.0137 (16)
C13	0.0474 (19)	0.0432 (18)	0.0350 (17)	-0.0008 (14)	0.0099 (14)	0.0068 (15)
C14	0.0435 (18)	0.0471 (19)	0.0363 (17)	-0.0077 (14)	-0.0042 (14)	-0.0025 (15)
C15	0.0373 (15)	0.0392 (16)	0.0281 (15)	-0.0009 (13)	-0.0026 (12)	0.0021 (13)
C16	0.0394 (17)	0.050 (2)	0.058 (2)	-0.0181 (15)	0.0056 (17)	0.0087 (18)
C11	0.0403 (4)	0.0396 (4)	0.0275 (4)	0.0017 (3)	0.0128 (3)	-0.0108 (3)
N1	0.0507 (17)	0.0366 (15)	0.0347 (15)	-0.0014 (13)	0.0048 (13)	-0.0007 (12)
N2	0.0326 (13)	0.0325 (14)	0.0351 (15)	0.0000 (10)	0.0028 (11)	-0.0005 (11)
O1	0.0323 (10)	0.0607 (15)	0.0387 (12)	-0.0070 (10)	-0.0006 (10)	-0.0059 (11)
O2	0.0565 (14)	0.0451 (13)	0.0281 (11)	-0.0041 (10)	0.0182 (11)	0.0042 (10)
O3	0.0564 (15)	0.0482 (14)	0.0597 (17)	-0.0259 (12)	0.0007 (13)	-0.0081 (12)
O4	0.0644 (16)	0.0474 (15)	0.0667 (18)	0.0107 (12)	0.0152 (14)	0.0303 (14)
S1	0.0299 (3)	0.0364 (4)	0.0261 (4)	-0.0028 (3)	0.0059 (3)	0.0004 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C1—C2	1.555 (4)	C10—C11	1.379 (4)
C1—H1A	0.9600	C10—C15	1.392 (4)

C1—H1B	0.9600	C10—S1	1.752 (3)
C1—H1C	0.9600	C11—C12	1.377 (5)
C2—C3	1.539 (4)	C11—H11	0.9300
C2—N1	1.544 (4)	C12—C13	1.400 (5)
C2—C11	1.761 (3)	C12—H12	0.9300
C3—N2	1.481 (4)	C13—C14	1.390 (5)
C3—C4	1.524 (4)	C13—C16	1.493 (5)
C3—H3	0.9800	C14—C15	1.373 (4)
C4—C9	1.382 (4)	C14—H14	0.9300
C4—C5	1.384 (4)	C15—H15	0.9300
C5—C6	1.380 (4)	C16—H16A	0.9600
C5—H5	0.9300	C16—H16B	0.9600
C6—C7	1.400 (5)	C16—H16C	0.9600
C6—H6	0.9300	N1—O3	1.207 (3)
C7—C8	1.367 (5)	N1—O4	1.212 (3)
C7—H7	0.9300	N2—S1	1.639 (3)
C8—C9	1.395 (4)	N2—H2A	0.89 (3)
C8—H8	0.9300	O1—S1	1.423 (2)
C9—H9	0.9300	O2—S1	1.432 (2)
C2—C1—H1A	109.5	C11—C10—C15	119.8 (3)
C2—C1—H1B	109.5	C11—C10—S1	121.1 (2)
H1A—C1—H1B	109.5	C15—C10—S1	119.1 (2)
C2—C1—H1C	109.5	C12—C11—C10	120.5 (3)
H1A—C1—H1C	109.5	C12—C11—H11	119.8
H1B—C1—H1C	109.5	C10—C11—H11	119.8
C3—C2—N1	105.9 (2)	C11—C12—C13	120.4 (3)
C3—C2—C1	115.5 (2)	C11—C12—H12	119.8
N1—C2—C1	110.5 (2)	C13—C12—H12	119.8
C3—C2—C11	110.3 (2)	C14—C13—C12	118.3 (3)
N1—C2—C11	103.78 (19)	C14—C13—C16	121.1 (3)
C1—C2—C11	110.1 (2)	C12—C13—C16	120.6 (3)
N2—C3—C4	115.3 (2)	C15—C14—C13	121.3 (3)
N2—C3—C2	105.9 (2)	C15—C14—H14	119.3
C4—C3—C2	113.6 (2)	C13—C14—H14	119.3
N2—C3—H3	107.2	C14—C15—C10	119.7 (3)
C4—C3—H3	107.2	C14—C15—H15	120.2
C2—C3—H3	107.2	C10—C15—H15	120.2
C9—C4—C5	119.1 (3)	C13—C16—H16A	109.5
C9—C4—C3	121.5 (3)	C13—C16—H16B	109.5
C5—C4—C3	119.4 (3)	H16A—C16—H16B	109.5
C6—C5—C4	121.4 (3)	C13—C16—H16C	109.5
C6—C5—H5	119.3	H16A—C16—H16C	109.5
C4—C5—H5	119.3	H16B—C16—H16C	109.5
C5—C6—C7	119.0 (3)	O3—N1—O4	123.8 (3)
C5—C6—H6	120.5	O3—N1—C2	117.7 (3)
C7—C6—H6	120.5	O4—N1—C2	118.6 (3)
C8—C7—C6	120.0 (3)	C3—N2—S1	118.6 (2)
C8—C7—H7	120.0	C3—N2—H2A	109 (2)
C6—C7—H7	120.0	S1—N2—H2A	107 (2)

## supplementary materials

C7—C8—C9	120.5 (3)	O1—S1—O2	119.59 (14)
C7—C8—H8	119.8	O1—S1—N2	107.37 (14)
C9—C8—H8	119.8	O2—S1—N2	105.25 (14)
C4—C9—C8	120.0 (3)	O1—S1—C10	108.79 (14)
C4—C9—H9	120.0	O2—S1—C10	107.97 (14)
C8—C9—H9	120.0	N2—S1—C10	107.24 (13)
N1—C2—C3—N2	59.6 (3)	C12—C13—C14—C15	-1.7 (5)
C1—C2—C3—N2	-63.2 (3)	C16—C13—C14—C15	176.7 (3)
C11—C2—C3—N2	171.28 (19)	C13—C14—C15—C10	1.2 (5)
N1—C2—C3—C4	-172.9 (2)	C11—C10—C15—C14	-0.4 (4)
C1—C2—C3—C4	64.4 (3)	S1—C10—C15—C14	178.4 (2)
C11—C2—C3—C4	-61.2 (3)	C3—C2—N1—O3	50.9 (3)
N2—C3—C4—C9	36.7 (4)	C1—C2—N1—O3	176.7 (3)
C2—C3—C4—C9	-85.7 (3)	C11—C2—N1—O3	-65.3 (3)
N2—C3—C4—C5	-141.2 (3)	C3—C2—N1—O4	-128.4 (3)
C2—C3—C4—C5	96.3 (3)	C1—C2—N1—O4	-2.5 (4)
C9—C4—C5—C6	0.2 (5)	C11—C2—N1—O4	115.5 (3)
C3—C4—C5—C6	178.1 (3)	C4—C3—N2—S1	80.2 (3)
C4—C5—C6—C7	-1.6 (5)	C2—C3—N2—S1	-153.3 (2)
C5—C6—C7—C8	2.4 (5)	C3—N2—S1—O1	41.9 (3)
C6—C7—C8—C9	-1.8 (5)	C3—N2—S1—O2	170.3 (2)
C5—C4—C9—C8	0.5 (5)	C3—N2—S1—C10	-74.9 (2)
C3—C4—C9—C8	-177.4 (3)	C11—C10—S1—O1	-18.0 (3)
C7—C8—C9—C4	0.4 (5)	C15—C10—S1—O1	163.2 (2)
C15—C10—C11—C12	0.0 (5)	C11—C10—S1—O2	-149.2 (2)
S1—C10—C11—C12	-178.8 (3)	C15—C10—S1—O2	32.0 (3)
C10—C11—C12—C13	-0.4 (5)	C11—C10—S1—N2	97.8 (3)
C11—C12—C13—C14	1.2 (5)	C15—C10—S1—N2	-80.9 (2)
C11—C12—C13—C16	-177.1 (3)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3 $\cdots$ O1	0.98	2.46	2.933 (4)	109
C3—H3 $\cdots$ O3	0.98	2.42	2.779 (4)	101
C1—H1A $\cdots$ O2 <sup>i</sup>	0.96	2.52	3.369 (4)	147
C1—H1B $\cdots$ O1 <sup>ii</sup>	0.96	2.58	3.310 (4)	133
N2—H2A $\cdots$ O2 <sup>i</sup>	0.89 (2)	2.32 (3)	3.141 (3)	153

Symmetry codes: (i)  $x+1/2, y, -z+3/2$ ; (ii)  $x+1, y, z$ .



Fig. 1

