

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

trans-4-Nitrophenyl 4-(tosyloxymethyl)-cyclohexanecarboxylateQing-Rong Qi,^a Wen-Cai Huang,^b Zhi-Hua Mao^c and Hu Zheng^{a*}^aDepartment of Medicinal Chemistry, West China School of Pharmacy, Sichuan University, Chengdu 610041, People's Republic of China, ^bDepartment of Pharmaceutical and Bioengineering, School of Chemical Engineering, Sichuan University, Chengdu 610065, People's Republic of China, and ^cThe Center for Testing and Analysis, Sichuan University, Chengdu 610064, People's Republic of China

Correspondence e-mail: wcums416@yahoo.com.cn

Received 18 December 2007; accepted 2 January 2008

Key indicators: single-crystal X-ray study; $T = 292$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.060; wR factor = 0.191; data-to-parameter ratio = 13.3.

The title compound, $\text{C}_{21}\text{H}_{23}\text{NO}_7\text{S}$, is an important intermediate in the synthesis of poly(amidoamine) dendrimers. The cyclohexane ring adopts a chair conformation. The dihedral angle between the two aromatic rings is $69.5(2)^\circ$. The molecules are linked into a zigzag chain along the b axis by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For related literature, see: Bucourt & Hainaut (1965); Dunitz & Strickler (1966); Sewald & Jakubke (2002); Luger *et al.* (1972); van Koningsveld & Jansen (1984).



Experimental

Crystal data

 $\text{C}_{21}\text{H}_{23}\text{NO}_7\text{S}$
 $M_r = 433.46$

 Monoclinic, $P2_1/c$
 $a = 20.499(5)$ Å

 $b = 6.111(3)$ Å
 $c = 17.250(4)$ Å
 $\beta = 102.04(3)^\circ$
 $V = 2113.4(13)$ Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.20$ mm⁻¹
 $T = 292(2)$ K
 $0.40 \times 0.38 \times 0.30$ mm

Data collection

 Enraf-Nonius CAD-4
 diffractometer
 Absorption correction: none
 4094 measured reflections
 3627 independent reflections

 1850 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.004$
 3 standard reflections
 every 300 reflections
 intensity decay: 2.1%

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.191$
 $S = 1.06$
 3627 reflections

 272 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.40$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.53$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C5}-\text{H5}\cdots\text{O7}^i$	0.93	2.49	3.290 (6)	144

Symmetry code: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *DIFRAC* (Gabe & White, 1993); cell refinement: *DIFRAC*; data reduction: *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2549).

References

- Bucourt, R. & Hainaut, D. (1965). *Bull. Soc. Chim. Fr.* **5**, 1366–1378.
 Dunitz, J. D. & Strickler, P. (1966). *Helv. Chim. Acta*, **49**, 290–291.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
 Gabe, E. J. & White, P. S. (1993). *DIFRAC*. American Crystallographic Association Meeting, Pittsburgh, Abstract PA 104.
 Koningsveld, H. van & Jansen, J. C. (1984). *Acta Cryst.* **B40**, 420–424.
 Luger, P., Plieth, K. & Ruban, G. (1972). *Acta Cryst.* **B28**, 706–710.
 Sewald, N. & Jakubke, H. D. (2002). *Peptides: Chemistry and Biology*, 1st ed. Weinheim: Wiley.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o385 [doi:10.1107/S1600536808000068]

***trans*-4-Nitrophenyl 4-(tosyloxymethyl)cyclohexanecarboxylate**

Q.-R. Qi, W.-C. Huang, Z.-H. Mao and H. Zheng

Comment

Activated esters are commonly used in peptide synthesis to facilitate the reaction of carboxylic acid and amine to obtain amide. *trans*-Nitrophenyl ester of N-protected amino acid can react with amino group quickly at room temperature with fairly good yields (Sewald *et al.*, 2002). In our work, cyclohexane derivatives were designed to be linked to poly(amidoamine)dendrimer[PAMAM] through the amide bond which could be formed by the reaction of terminal amino groups of PAMAM dendrimer and cyclohexanecarboxylic acid. So the title compound, a *p*-nitrophenyl ester of *trans*-4-(tosyloxymethyl) cyclohexanecarboxylic acid was synthesized. As expected, the modification of periphery of PAMAM dendrimer effectively is realised under mild conditions. We report here the crystal structure of the title compound.

The cyclohexane ring of the title compound adopts a chair conformation. The average C—C bond length of the cyclohexane ring is 1.521 (5) Å, which is close to that of *trans*-1,4-cyclohexanedicarboxylic acid (1.523 (3) Å, Von Luger *et al.*, 1972). The mean endocyclic angle of the cyclohexane is 111.08 (3)°, which is close to that observed for cyclohexane rings (111.1°, Bucourt & Hainaut, 1965; 111.4 (4)°, Dunitz & Strickler, 1966; Von Luger *et al.*, 1972).

The molecules are linked into a zigzag chain along the *b* axis by C—H···O hydrogen bonds (Table 1).

Experimental

trans-4-(Tosyloxymethyl)cyclohexanecarboxylic acid (10 mmol) and *p*-nitrophenol (11 mmol) were dissolved in ethyl acetate (10 ml) and the solution was cooled to 273 K in an ice bath. Then a solution of DCC (11 mmol) in ethyl acetate (5 ml) was added dropwise with stirring. After the addition, the stirring was continued for 30 min at 273 K and for 24 h at room temperature. Then acetic acid (0.5 ml) was added and after 30 min the reaction solution was filtered and the filtrate was evaporated to leave a yellow solid. The title compound was recrystallized from acetone. Single crystals suitable for X-ray analysis were obtained by slow evaporation of a acetone-water solution at room temperature.

Refinement

H atoms were positioned geometrically (C—H = 0.93–0.98 Å) and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$.

Figures



Fig. 1. The molecular structure of (I), showing the atomic numbering scheme. Displacement ellipsoids drawn at the 30% probability level.

trans-4-Nitrophenyl 4-(tosyloxymethyl)cyclohexanecarboxylate

Crystal data

C₂₁H₂₃NO₇S

M_r = 433.46

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 20.499 (5) Å

b = 6.111 (3) Å

c = 17.250 (4) Å

β = 102.04 (3)°

V = 2113.4 (13) Å³

Z = 4

*F*₀₀₀ = 912

D_x = 1.362 Mg m⁻³

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 33 reflections

θ = 4.5–7.6°

μ = 0.20 mm⁻¹

T = 292 (2) K

Block, colourless

0.40 × 0.38 × 0.30 mm

Data collection

Enraf–Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 292(2) K

ω/2θ scans

Absorption correction: none

4094 measured reflections

3627 independent reflections

1850 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.004

θ_{max} = 25.0°

θ_{min} = 1.0°

h = -8→24

k = -7→0

l = -20→19

3 standard reflections

every 300 reflections

intensity decay: 2.1%

Refinement

Refinement on *F*²

Least-squares matrix: full

R [*F*² > 2σ(*F*²)] = 0.060

wR(*F*²) = 0.191

S = 1.06

3627 reflections

272 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F*_o²) + (0.097*P*)²]

where *P* = (*F*_o² + 2*F*_c²)/3

(Δ/σ)_{max} = 0.001

Δρ_{max} = 0.40 e Å⁻³

Δρ_{min} = -0.53 e Å⁻³

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.30112 (5)	-0.15510 (18)	0.17299 (7)	0.0556 (4)
O1	0.30040 (14)	-0.0770 (6)	0.09454 (16)	0.0702 (9)
O2	0.30099 (14)	-0.3830 (5)	0.1871 (2)	0.0763 (10)
O3	0.23652 (12)	-0.0742 (5)	0.19982 (18)	0.0632 (8)
O4	-0.09447 (15)	0.2718 (6)	0.1532 (2)	0.0885 (12)
O5	-0.08914 (13)	0.5252 (5)	0.06303 (18)	0.0686 (9)
O6	-0.39478 (15)	0.7233 (6)	-0.0126 (2)	0.0813 (11)
O7	-0.36075 (17)	0.9887 (7)	0.0674 (2)	0.0936 (12)
N1	-0.35100 (18)	0.8235 (8)	0.0323 (2)	0.0635 (10)
C1	0.39156 (18)	0.1686 (7)	0.2212 (2)	0.0500 (10)
H1	0.3740	0.2367	0.1732	0.060*
C2	0.44258 (19)	0.2665 (7)	0.2753 (2)	0.0536 (11)
H2	0.4589	0.4016	0.2633	0.064*
C3	0.46966 (18)	0.1673 (8)	0.3465 (2)	0.0526 (10)
C4	0.4444 (2)	-0.0304 (9)	0.3636 (3)	0.0650 (13)
H4	0.4618	-0.0973	0.4119	0.078*
C5	0.3931 (2)	-0.1337 (8)	0.3100 (2)	0.0593 (12)
H5	0.3769	-0.2690	0.3220	0.071*
C6	0.36691 (17)	-0.0320 (6)	0.2393 (2)	0.0440 (9)
C7	0.5248 (2)	0.2787 (10)	0.4050 (3)	0.0811 (16)
H7A	0.5125	0.4279	0.4121	0.122*
H7B	0.5652	0.2753	0.3852	0.122*
H7C	0.5316	0.2037	0.4550	0.122*
C8	0.21871 (19)	0.1584 (7)	0.1883 (3)	0.0657 (12)
H8A	0.2286	0.2104	0.1389	0.079*
H8B	0.2442	0.2448	0.2313	0.079*
C9	0.14546 (17)	0.1807 (7)	0.1864 (2)	0.0507 (10)
H9	0.1370	0.1230	0.2364	0.061*
C10	0.10207 (18)	0.0551 (7)	0.1192 (2)	0.0527 (10)
H10A	0.1131	-0.0992	0.1249	0.063*
H10B	0.1115	0.1046	0.0692	0.063*
C11	0.02817 (18)	0.0850 (8)	0.1176 (2)	0.0564 (11)
H11A	0.0178	0.0238	0.1655	0.068*

supplementary materials

H11B	0.0023	0.0067	0.0726	0.068*
C12	0.00917 (18)	0.3266 (7)	0.1115 (2)	0.0525 (11)
H12	0.0175	0.3836	0.0614	0.063*
C13	0.05248 (19)	0.4533 (7)	0.1792 (3)	0.0614 (12)
H13A	0.0426	0.4050	0.2291	0.074*
H13B	0.0418	0.6078	0.1732	0.074*
C14	0.1265 (2)	0.4210 (7)	0.1814 (3)	0.0648 (12)
H14A	0.1374	0.4839	0.1340	0.078*
H14B	0.1523	0.4975	0.2269	0.078*
C15	-0.0628 (2)	0.3620 (8)	0.1128 (2)	0.0545 (11)
C16	-0.1560 (2)	0.5894 (8)	0.0600 (2)	0.0574 (12)
C17	-0.2075 (2)	0.4605 (8)	0.0214 (3)	0.0662 (13)
H17	-0.1994	0.3249	0.0008	0.079*
C18	-0.2720 (2)	0.5386 (8)	0.0140 (3)	0.0631 (13)
H18	-0.3080	0.4554	-0.0120	0.076*
C19	-0.28246 (19)	0.7380 (8)	0.0449 (2)	0.0509 (10)
C20	-0.2310 (2)	0.8667 (8)	0.0847 (2)	0.0623 (12)
H20	-0.2392	1.0013	0.1059	0.075*
C21	-0.1658 (2)	0.7874 (9)	0.0920 (3)	0.0665 (13)
H21	-0.1297	0.8690	0.1185	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0444 (6)	0.0441 (7)	0.0734 (8)	0.0019 (5)	0.0014 (5)	-0.0017 (6)
O1	0.068 (2)	0.082 (2)	0.0556 (18)	0.0010 (16)	0.0017 (14)	-0.0056 (17)
O2	0.0633 (19)	0.041 (2)	0.114 (3)	-0.0006 (15)	-0.0060 (17)	0.0007 (19)
O3	0.0386 (15)	0.053 (2)	0.096 (2)	0.0042 (13)	0.0114 (14)	0.0135 (17)
O4	0.0562 (19)	0.106 (3)	0.108 (3)	0.0247 (18)	0.0286 (18)	0.052 (2)
O5	0.0447 (16)	0.079 (2)	0.082 (2)	0.0155 (15)	0.0136 (14)	0.0340 (19)
O6	0.0471 (18)	0.101 (3)	0.091 (2)	0.0097 (18)	0.0021 (17)	-0.007 (2)
O7	0.082 (2)	0.105 (3)	0.095 (3)	0.027 (2)	0.0196 (19)	-0.024 (3)
N1	0.060 (2)	0.073 (3)	0.059 (2)	0.018 (2)	0.0138 (19)	0.000 (2)
C1	0.050 (2)	0.047 (3)	0.054 (2)	0.0062 (19)	0.0105 (18)	0.002 (2)
C2	0.052 (2)	0.047 (3)	0.062 (3)	-0.0082 (19)	0.012 (2)	0.003 (2)
C3	0.043 (2)	0.063 (3)	0.054 (2)	-0.004 (2)	0.0146 (18)	-0.001 (2)
C4	0.064 (3)	0.078 (4)	0.050 (2)	-0.004 (2)	0.006 (2)	0.011 (2)
C5	0.061 (3)	0.057 (3)	0.062 (3)	-0.006 (2)	0.016 (2)	0.009 (2)
C6	0.043 (2)	0.034 (2)	0.056 (2)	0.0020 (16)	0.0105 (17)	0.0054 (19)
C7	0.063 (3)	0.106 (5)	0.070 (3)	-0.017 (3)	0.003 (2)	-0.013 (3)
C8	0.043 (2)	0.045 (3)	0.109 (4)	0.0012 (19)	0.015 (2)	0.001 (3)
C9	0.041 (2)	0.047 (3)	0.064 (3)	0.0016 (17)	0.0104 (18)	0.005 (2)
C10	0.050 (2)	0.041 (3)	0.068 (3)	0.0056 (18)	0.0142 (19)	-0.006 (2)
C11	0.047 (2)	0.059 (3)	0.061 (3)	-0.0058 (19)	0.0077 (19)	-0.011 (2)
C12	0.042 (2)	0.063 (3)	0.053 (2)	0.008 (2)	0.0124 (17)	0.010 (2)
C13	0.048 (2)	0.042 (3)	0.095 (3)	0.0063 (19)	0.016 (2)	-0.008 (2)
C14	0.050 (2)	0.046 (3)	0.098 (3)	-0.004 (2)	0.013 (2)	-0.006 (3)
C15	0.050 (2)	0.056 (3)	0.057 (2)	0.006 (2)	0.010 (2)	0.012 (2)

C16	0.047 (2)	0.067 (3)	0.057 (3)	0.008 (2)	0.0088 (19)	0.019 (2)
C17	0.059 (3)	0.056 (3)	0.079 (3)	0.014 (2)	0.005 (2)	-0.014 (3)
C18	0.045 (2)	0.071 (4)	0.067 (3)	-0.001 (2)	-0.001 (2)	-0.004 (3)
C19	0.048 (2)	0.054 (3)	0.049 (2)	0.005 (2)	0.0060 (18)	0.001 (2)
C20	0.062 (3)	0.059 (3)	0.063 (3)	0.007 (2)	0.007 (2)	-0.004 (2)
C21	0.056 (3)	0.069 (4)	0.068 (3)	-0.005 (2)	-0.003 (2)	0.000 (3)

Geometric parameters (Å, °)

S1—O2	1.414 (3)	C9—C10	1.515 (5)
S1—O1	1.432 (3)	C9—C14	1.517 (6)
S1—O3	1.571 (3)	C9—H9	0.98
S1—C6	1.747 (4)	C10—C11	1.520 (5)
O3—C8	1.471 (5)	C10—H10A	0.97
O4—C15	1.184 (5)	C10—H10B	0.97
O5—C15	1.353 (5)	C11—C12	1.525 (6)
O5—C16	1.416 (5)	C11—H11A	0.97
O6—N1	1.221 (5)	C11—H11B	0.97
O7—N1	1.214 (5)	C12—C15	1.495 (5)
N1—C19	1.472 (5)	C12—C13	1.523 (6)
C1—C2	1.383 (5)	C12—H12	0.98
C1—C6	1.386 (6)	C13—C14	1.523 (5)
C1—H1	0.93	C13—H13A	0.97
C2—C3	1.379 (6)	C13—H13B	0.97
C2—H2	0.93	C14—H14A	0.97
C3—C4	1.371 (6)	C14—H14B	0.97
C3—C7	1.511 (6)	C16—C21	1.362 (6)
C4—C5	1.399 (6)	C16—C17	1.374 (6)
C4—H4	0.93	C17—C18	1.386 (6)
C5—C6	1.373 (5)	C17—H17	0.93
C5—H5	0.93	C18—C19	1.365 (6)
C7—H7A	0.96	C18—H18	0.93
C7—H7B	0.96	C19—C20	1.379 (6)
C7—H7C	0.96	C20—C21	1.402 (6)
C8—C9	1.501 (5)	C20—H20	0.93
C8—H8A	0.97	C21—H21	0.93
C8—H8B	0.97		
O2—S1—O1	119.4 (2)	C11—C10—H10A	109.2
O2—S1—O3	103.13 (18)	C9—C10—H10B	109.2
O1—S1—O3	109.29 (18)	C11—C10—H10B	109.2
O2—S1—C6	109.87 (18)	H10A—C10—H10B	107.9
O1—S1—C6	109.28 (19)	C10—C11—C12	110.9 (3)
O3—S1—C6	104.74 (18)	C10—C11—H11A	109.5
C8—O3—S1	117.7 (3)	C12—C11—H11A	109.5
C15—O5—C16	118.9 (3)	C10—C11—H11B	109.5
O7—N1—O6	123.9 (4)	C12—C11—H11B	109.5
O7—N1—C19	118.2 (4)	H11A—C11—H11B	108.0
O6—N1—C19	117.9 (4)	C15—C12—C13	109.5 (3)
C2—C1—C6	119.4 (4)	C15—C12—C11	112.2 (4)

supplementary materials

C2—C1—H1	120.3	C13—C12—C11	109.9 (3)
C6—C1—H1	120.3	C15—C12—H12	108.4
C3—C2—C1	121.2 (4)	C13—C12—H12	108.4
C3—C2—H2	119.4	C11—C12—H12	108.4
C1—C2—H2	119.4	C12—C13—C14	111.8 (4)
C4—C3—C2	118.6 (4)	C12—C13—H13A	109.3
C4—C3—C7	121.1 (4)	C14—C13—H13A	109.3
C2—C3—C7	120.2 (4)	C12—C13—H13B	109.3
C3—C4—C5	121.4 (4)	C14—C13—H13B	109.3
C3—C4—H4	119.3	H13A—C13—H13B	107.9
C5—C4—H4	119.3	C9—C14—C13	111.6 (3)
C6—C5—C4	118.9 (4)	C9—C14—H14A	109.3
C6—C5—H5	120.5	C13—C14—H14A	109.3
C4—C5—H5	120.5	C9—C14—H14B	109.3
C5—C6—C1	120.4 (4)	C13—C14—H14B	109.3
C5—C6—S1	119.5 (3)	H14A—C14—H14B	108.0
C1—C6—S1	120.1 (3)	O4—C15—O5	121.4 (4)
C3—C7—H7A	109.5	O4—C15—C12	127.3 (4)
C3—C7—H7B	109.5	O5—C15—C12	111.2 (4)
H7A—C7—H7B	109.5	C21—C16—C17	122.9 (4)
C3—C7—H7C	109.5	C21—C16—O5	117.0 (4)
H7A—C7—H7C	109.5	C17—C16—O5	120.0 (4)
H7B—C7—H7C	109.5	C16—C17—C18	118.0 (5)
O3—C8—C9	108.0 (3)	C16—C17—H17	121.0
O3—C8—H8A	110.1	C18—C17—H17	121.0
C9—C8—H8A	110.1	C19—C18—C17	119.7 (4)
O3—C8—H8B	110.1	C19—C18—H18	120.1
C9—C8—H8B	110.1	C17—C18—H18	120.1
H8A—C8—H8B	108.4	C18—C19—C20	122.6 (4)
C8—C9—C10	113.3 (4)	C18—C19—N1	118.8 (4)
C8—C9—C14	109.4 (3)	C20—C19—N1	118.6 (4)
C10—C9—C14	110.1 (3)	C19—C20—C21	117.6 (5)
C8—C9—H9	107.9	C19—C20—H20	121.2
C10—C9—H9	107.9	C21—C20—H20	121.2
C14—C9—H9	107.9	C16—C21—C20	119.3 (4)
C9—C10—C11	112.2 (3)	C16—C21—H21	120.4
C9—C10—H10A	109.2	C20—C21—H21	120.4

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C5—H5 \cdots O7 ⁱ	0.93	2.49	3.290 (6)	144

Symmetry codes: (i) $-x, y-3/2, -z+1/2$.

Fig. 1

