

3,3-Dimethyl-1-[5-(1*H*-1,2,4-triazol-1-ylmethyl)-1,3,4-thiadiazol-2-ylsulfanyl]butan-2-one

Qing-Li Wei, Fu-Jin He, Fang Li and Sai Bi*

College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, 266042 Qingdao, Shandong, People's Republic of China
Correspondence e-mail: qustchemistry@126.com


Received 7 November 2007; accepted 24 December 2007

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.051; wR factor = 0.144; data-to-parameter ratio = 14.7.

In the molecule of the title compound, $\text{C}_{11}\text{H}_{15}\text{N}_5\text{OS}_2$, the thiadiazole and triazole rings are not coplanar, the dihedral angle formed by their mean planes being $59.9(2)^\circ$. The exocyclic S atom, and the methylene, carbonyl, *tert*-butyl and one methyl carbon form an approximately planar zigzag chain, which makes a dihedral angle of $74.6(1)^\circ$ with the thiadiazole ring.

Related literature

For the structure of the related compound, 1-(2,4-dichlorophenyl)-2-[5-(1*H*-1,2,4-triazol-1-ylmethyl)-1,3,4-thiadiazol-2-ylsulfanyl] ethanone, see: Wei *et al.* (2007). For the synthesis of the starting material 5-(1*H*-1,2,4-triazol-1-ylmethyl)-1,3,4-thiadiazole-2(3*H*)-thione, see: Hu *et al.* (2006), Xu *et al.* (2005, 2006).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{15}\text{N}_5\text{OS}_2$	$\gamma = 65.416(1)^\circ$
$M_r = 297.40$	$V = 731.3(1) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.9723(8) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.1103(8) \text{ \AA}$	$\mu = 0.36 \text{ mm}^{-1}$
$c = 10.1734(8) \text{ \AA}$	$T = 293(2) \text{ K}$
$\alpha = 60.728(1)^\circ$	$0.41 \times 0.22 \times 0.18 \text{ mm}$
$\beta = 80.340(1)^\circ$	

Data collection

Siemens SMART 1000 CCD area-detector diffractometer	3810 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	2529 independent reflections
$T_{\min} = 0.865$, $T_{\max} = 0.938$	2246 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.010$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	172 parameters
$wR(F^2) = 0.143$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\text{max}} = 0.51 \text{ e \AA}^{-3}$
2529 reflections	$\Delta\rho_{\text{min}} = -0.38 \text{ e \AA}^{-3}$

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT (Siemens, 1996); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 1997); software used to prepare material for publication: SHELXTL, PARST (Nardelli, 1995) and PLATON (Spek, 2003).

This project was supported by the Natural Science Foundation of Shandong Province (grant Nos. Z2006B01 and Y2006B07).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YA2060).

References

- Hu, Z.-Q., Yang, Y.-X., Shang, Y.-Q., Zhou, K. & Xu, L.-Z. (2006). *Acta Cryst.* **E62**, o3457–o3458.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). SHELXTL. Version 5.10. Bruker AXS, Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). SMART and SAINT. Siemens Analytical X-ray Instruments, Inc., Madison, Wisconsin, USA.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Wei, Q.-L., He, F.-J., Lin, Y.-S. & Bi, S. (2007). *Acta Cryst.* **E63**, o3649.
- Xu, L.-Z., Li, W.-H., Si, G.-D., Zhu, C.-Y., Li, K. & Hou, B.-R. (2005). *Chem. Res. Chin. Univ.* **21**, 444–446.
- Xu, L.-Z., Shang, Y.-Q., Yu, G.-P., Li, K. & Si, G.-D. (2006). *Chin. J. Struct. Chem.* **25**, 673–676.

supplementary materials

Acta Cryst. (2008). E64, o412 [doi:10.1107/S1600536807068286]

3,3-Dimethyl-1-[5-(1*H*-1,2,4-triazol-1-ylmethyl)-1,3,4-thiadiazol-2-ylsulfanyl]butan-2-one

Q.-L. Wei, F.-J. He, F. Li and S. Bi

Comment

Recently, we have reported the structure of 1-(2,4-dichlorophenyl)-2-[5-(1*H*-1,2,4-triazol-1-ylmethyl)-1,3,4-thiadiazol-2-ylsulfanyl] ethanone (Wei *et al.*, 2007). As part of our ongoing investigation of biological properties of 1,2,4-triazole and 1,3,4-thiadiazole derivatives, the title compound, (I), was synthesized; its crystal structure is reported here.

The bond lengths and angles are comparable with those of the above mentioned related compound, reported by Wei *et al.* (2007). The whole molecule is non-planar with a dihedral angle of 59.9 (2)° between the thiadiazole (C1/C2/N1/N2/S1) and triazole (N3—N5/C4/C5) rings. The S2—C6—C7—C8—C10 atoms form approximately planar zigzag chain, which makes a dihedral angle of 74.6 (1)° with the thiadiazole ring.

Experimental

8 mmol of 5-((1*H*-1,2,4-triazol-1-yl)methyl)-1,3,4-thiadiazole-2(3*H*)-thione (Hu *et al.*, 2006; Xu *et al.*, 2005; Xu *et al.*, 2006) was refluxed for 4 h with 8 mmol of 1-bromo-3,3-dimethylbutan-2-one in 50 ml of acetone in the presence of 8 mmol of triethylamine. The solid that precipitated was recrystallized from ethanol (1.21 g, yield 50.86%). Single crystals suitable for X-ray measurements were obtained by slow evaporation of ethylacetate solution at room temperature.

Refinement

After the H atoms were located in the difference map, they were fixed geometrically in the idealized positions and allowed to ride on the parent C atoms, with C—H distances of 0.96 Å (methyl), 0.97 Å (CH₂) or 0.93 Å (CH), and with $U_{\text{iso}}(\text{H})$ values of $1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{C})$ (for methyl H atoms).

Figures




Fig. 1. The structure of the title compound (I), showing 50% probability displacement ellipsoids and the atom numbering scheme.

3,3-Dimethyl-1-[5-(1*H*-1,2,4-triazol-1-ylmethyl)-1,3,4-thiadiazol-2-ylsulfanyl]butan-2-one

Crystal data

C₁₁H₁₅N₅OS₂

$M_r = 297.40$

Triclinic, $P\bar{1}$

$Z = 2$

$F(000) = 312$

$D_x = 1.351 \text{ Mg m}^{-3}$

supplementary materials

$a = 8.9723 (8) \text{ \AA}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$b = 10.1103 (8) \text{ \AA}$	Cell parameters from 2210 reflections
$c = 10.1734 (8) \text{ \AA}$	$\theta = 2.5\text{--}25.6^\circ$
$\alpha = 60.728 (1)^\circ$	$\mu = 0.36 \text{ mm}^{-1}$
$\beta = 80.340 (1)^\circ$	$T = 293 \text{ K}$
$\gamma = 65.416 (1)^\circ$	Block, colourless
$V = 731.3 (1) \text{ \AA}^3$	$0.41 \times 0.22 \times 0.18 \text{ mm}$

Data collection

Siemens SMART 1000 CCD area-detector diffractometer	2529 independent reflections
Radiation source: fine-focus sealed tube graphite	2246 reflections with $I > 2\sigma(I)$
Detector resolution: $8.33 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.010$
ω scans	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.3^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 10$
$T_{\text{min}} = 0.865$, $T_{\text{max}} = 0.938$	$k = -9 \rightarrow 12$
3810 measured reflections	$l = -12 \rightarrow 12$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.050$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.143$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0806P)^2 + 0.357P]$
2529 reflections	where $P = (F_o^2 + 2F_c^2)/3$
172 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.51 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.38 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.5556 (3)	0.4597 (3)	0.7067 (3)	0.0680 (7)
N2	0.6886 (3)	0.4050 (3)	0.7986 (3)	0.0670 (6)
N3	0.2627 (3)	0.8466 (3)	0.4764 (2)	0.0546 (5)
N4	0.2154 (4)	0.9710 (3)	0.5103 (3)	0.0718 (7)
N5	0.2371 (4)	1.0629 (3)	0.2618 (3)	0.0762 (7)
O1	1.0413 (3)	0.4947 (2)	0.7133 (2)	0.0746 (6)
S1	0.49906 (8)	0.68646 (9)	0.77341 (9)	0.0627 (3)
S2	0.81659 (9)	0.48127 (9)	0.96116 (8)	0.0609 (2)
C1	0.6733 (3)	0.5099 (3)	0.8425 (3)	0.0508 (6)
C2	0.4489 (3)	0.6004 (3)	0.6854 (3)	0.0548 (6)
C3	0.2921 (4)	0.6795 (4)	0.5949 (4)	0.0698 (8)
H3A	0.2954	0.6133	0.5496	0.084*
H3B	0.2013	0.6817	0.6618	0.084*
C4	0.2003 (4)	1.0979 (3)	0.3766 (3)	0.0657 (7)
H4	0.1665	1.2043	0.3630	0.079*
C5	0.2751 (4)	0.9038 (4)	0.3298 (3)	0.0695 (8)
H5	0.3066	0.8398	0.2807	0.083*
C6	0.9842 (3)	0.3090 (3)	0.9519 (3)	0.0543 (6)
H6A	1.0658	0.2611	1.0311	0.065*
H6B	0.9439	0.2261	0.9711	0.065*
C7	1.0658 (3)	0.3532 (3)	0.8017 (3)	0.0501 (6)
C8	1.1780 (4)	0.2127 (3)	0.7691 (3)	0.0629 (7)
C9	1.2994 (7)	0.0845 (7)	0.8962 (5)	0.199 (4)
H9A	1.3660	0.1311	0.9096	0.298*
H9B	1.3679	-0.0042	0.8744	0.298*
H9C	1.2430	0.0437	0.9868	0.298*
C10	1.2626 (5)	0.2749 (5)	0.6225 (4)	0.0910 (11)
H10A	1.3325	0.3208	0.6317	0.136*
H10B	1.1816	0.3581	0.5422	0.136*
H10C	1.3272	0.1853	0.6008	0.136*
C11	1.0686 (7)	0.1417 (8)	0.7445 (8)	0.154 (3)
H11A	1.1360	0.0522	0.7235	0.231*
H11B	0.9921	0.2261	0.6609	0.231*
H11C	1.0097	0.1028	0.8338	0.231*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0740 (15)	0.0541 (13)	0.0763 (16)	-0.0174 (12)	-0.0189 (12)	-0.0300 (12)
N2	0.0678 (14)	0.0555 (13)	0.0776 (16)	-0.0112 (11)	-0.0206 (12)	-0.0341 (12)
N3	0.0551 (12)	0.0475 (11)	0.0590 (13)	-0.0186 (9)	-0.0090 (10)	-0.0209 (10)
N4	0.0947 (18)	0.0605 (14)	0.0618 (14)	-0.0268 (13)	0.0001 (13)	-0.0315 (12)
N5	0.0926 (19)	0.0611 (15)	0.0560 (14)	-0.0159 (13)	-0.0140 (13)	-0.0200 (12)
O1	0.0788 (13)	0.0500 (11)	0.0763 (13)	-0.0282 (10)	0.0123 (11)	-0.0161 (10)

supplementary materials

S1	0.0575 (4)	0.0554 (4)	0.0780 (5)	-0.0157 (3)	-0.0043 (3)	-0.0359 (4)
S2	0.0656 (4)	0.0655 (5)	0.0606 (4)	-0.0234 (3)	-0.0027 (3)	-0.0359 (4)
C1	0.0536 (13)	0.0487 (13)	0.0494 (13)	-0.0200 (11)	0.0021 (11)	-0.0222 (11)
C2	0.0572 (14)	0.0470 (14)	0.0568 (14)	-0.0238 (12)	-0.0020 (11)	-0.0175 (11)
C3	0.0673 (17)	0.0545 (15)	0.0793 (19)	-0.0301 (14)	-0.0174 (15)	-0.0138 (14)
C4	0.0773 (18)	0.0486 (15)	0.0634 (17)	-0.0160 (13)	-0.0089 (14)	-0.0239 (13)
C5	0.0795 (19)	0.0606 (17)	0.0645 (18)	-0.0124 (14)	-0.0127 (14)	-0.0336 (15)
C6	0.0585 (14)	0.0493 (13)	0.0493 (14)	-0.0200 (11)	-0.0091 (11)	-0.0161 (11)
C7	0.0472 (12)	0.0462 (13)	0.0537 (14)	-0.0198 (10)	-0.0097 (10)	-0.0162 (11)
C8	0.0658 (16)	0.0536 (15)	0.0588 (16)	-0.0192 (13)	0.0017 (13)	-0.0216 (13)
C9	0.181 (6)	0.165 (5)	0.085 (3)	0.104 (5)	-0.051 (3)	-0.063 (3)
C10	0.095 (3)	0.095 (3)	0.082 (2)	-0.037 (2)	0.021 (2)	-0.046 (2)
C11	0.170 (5)	0.186 (5)	0.237 (7)	-0.129 (5)	0.117 (5)	-0.176 (6)

Geometric parameters (Å, °)

N1—C2	1.274 (3)	C5—H5	0.9300
N1—N2	1.387 (3)	C6—C7	1.516 (4)
N2—C1	1.290 (3)	C6—H6A	0.9700
N3—C5	1.313 (4)	C6—H6B	0.9700
N3—N4	1.347 (3)	C7—C8	1.517 (4)
N3—C3	1.456 (3)	C8—C9	1.482 (5)
N4—C4	1.314 (4)	C8—C10	1.515 (4)
N5—C5	1.310 (4)	C8—C11	1.544 (5)
N5—C4	1.332 (4)	C9—H9A	0.9600
O1—C7	1.205 (3)	C9—H9B	0.9600
S1—C1	1.715 (3)	C9—H9C	0.9600
S1—C2	1.728 (3)	C10—H10A	0.9600
S2—C1	1.751 (3)	C10—H10B	0.9600
S2—C6	1.799 (3)	C10—H10C	0.9600
C2—C3	1.498 (4)	C11—H11A	0.9600
C3—H3A	0.9700	C11—H11B	0.9600
C3—H3B	0.9700	C11—H11C	0.9600
C4—H4	0.9300		
C2—N1—N2	113.0 (2)	C7—C6—H6B	108.8
C1—N2—N1	111.5 (2)	S2—C6—H6B	108.8
C5—N3—N4	109.3 (2)	H6A—C6—H6B	107.7
C5—N3—C3	130.1 (3)	O1—C7—C6	120.5 (2)
N4—N3—C3	120.6 (2)	O1—C7—C8	122.2 (2)
C4—N4—N3	102.1 (2)	C6—C7—C8	117.3 (2)
C5—N5—C4	102.3 (2)	C9—C8—C10	111.0 (4)
C1—S1—C2	86.46 (12)	C9—C8—C7	110.6 (3)
C1—S2—C6	98.80 (12)	C10—C8—C7	110.9 (2)
N2—C1—S1	114.8 (2)	C9—C8—C11	110.8 (5)
N2—C1—S2	124.2 (2)	C10—C8—C11	106.0 (3)
S1—C1—S2	121.01 (15)	C7—C8—C11	107.3 (3)
N1—C2—C3	121.9 (3)	C8—C9—H9A	109.5
N1—C2—S1	114.2 (2)	C8—C9—H9B	109.5
C3—C2—S1	123.9 (2)	H9A—C9—H9B	109.5

N3—C3—C2	112.9 (2)	C8—C9—H9C	109.5
N3—C3—H3A	109.0	H9A—C9—H9C	109.5
C2—C3—H3A	109.0	H9B—C9—H9C	109.5
N3—C3—H3B	109.0	C8—C10—H10A	109.5
C2—C3—H3B	109.0	C8—C10—H10B	109.5
H3A—C3—H3B	107.8	H10A—C10—H10B	109.5
N4—C4—N5	115.0 (3)	C8—C10—H10C	109.5
N4—C4—H4	122.5	H10A—C10—H10C	109.5
N5—C4—H4	122.5	H10B—C10—H10C	109.5
N5—C5—N3	111.1 (3)	C8—C11—H11A	109.5
N5—C5—H5	124.4	C8—C11—H11B	109.5
N3—C5—H5	124.4	H11A—C11—H11B	109.5
C7—C6—S2	113.74 (17)	C8—C11—H11C	109.5
C7—C6—H6A	108.8	H11A—C11—H11C	109.5
S2—C6—H6A	108.8	H11B—C11—H11C	109.5
C2—N1—N2—C1	0.3 (4)	S1—C2—C3—N3	53.9 (4)
C5—N3—N4—C4	0.9 (3)	N3—N4—C4—N5	-1.1 (4)
C3—N3—N4—C4	-179.3 (2)	C5—N5—C4—N4	0.9 (4)
N1—N2—C1—S1	-1.4 (3)	C4—N5—C5—N3	-0.3 (4)
N1—N2—C1—S2	179.3 (2)	N4—N3—C5—N5	-0.4 (4)
C2—S1—C1—N2	1.6 (2)	C3—N3—C5—N5	179.8 (3)
C2—S1—C1—S2	-179.13 (18)	C1—S2—C6—C7	71.9 (2)
C6—S2—C1—N2	13.2 (3)	S2—C6—C7—O1	15.0 (3)
C6—S2—C1—S1	-166.03 (16)	S2—C6—C7—C8	-165.12 (19)
N2—N1—C2—C3	-177.4 (3)	O1—C7—C8—C9	129.1 (4)
N2—N1—C2—S1	0.9 (3)	C6—C7—C8—C9	-50.8 (5)
C1—S1—C2—N1	-1.4 (2)	O1—C7—C8—C10	5.4 (4)
C1—S1—C2—C3	176.9 (2)	C6—C7—C8—C10	-174.5 (2)
C5—N3—C3—C2	105.8 (3)	O1—C7—C8—C11	-109.9 (4)
N4—N3—C3—C2	-74.1 (4)	C6—C7—C8—C11	70.2 (4)
N1—C2—C3—N3	-127.9 (3)		

Fig. 1

