

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2-Butoxyphenyl)boronic acid

Marek Dąbrowski, Sergiusz Luliński and Janusz Serwatowski*

Warsaw University of Technology, Faculty of Chemistry, Noakowskiego 3, 00-664 Warsaw, Poland

Correspondence e-mail: marek@ch.pw.edu.pl


Received 27 December 2007; accepted 7 January 2008

Key indicators: single-crystal X-ray study; $T = 102$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.033; wR factor = 0.097; data-to-parameter ratio = 12.9.

The title compound, 2-($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{O}$) $\text{C}_6\text{H}_4\text{B}(\text{OH})_2$, exists as a centrosymmetric hydrogen-bonded dimer. Dimers are linked *via* $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ [with closest $\text{C}\cdots\text{C}$ contact of 3.540 (3) Å] interactions to produce a two-dimensional array.

Related literature

For related literature, see: Rettig & Trotter (1977). For the structures of related *ortho*-alkoxyarylboronic acids, see: Dabrowski *et al.* (2006); Serwatowski *et al.* (2006); Yang *et al.* (2005).



Experimental

Crystal data

$\text{C}_{10}\text{H}_{15}\text{BO}_3$ $V = 1062.98$ (9) Å³
 $M_r = 194.03$ $Z = 4$
 Monoclinic, $P2_1/c$ Mo $K\alpha$ radiation
 $a = 7.4809$ (4) Å $\mu = 0.09$ mm⁻¹
 $b = 15.3510$ (7) Å $T = 102$ (2) K
 $c = 9.2824$ (4) Å $0.74 \times 0.47 \times 0.32$ mm
 $\beta = 94.299$ (4)°

Data collection

Kuma KM4 CCD diffractometer 9363 measured reflections
 Absorption correction: multi-scan 2419 independent reflections
 (*CrysAlis RED*; Oxford 1924 reflections with $I > 2\sigma(I)$)
 Diffraction, 2005) $R_{\text{int}} = 0.012$
 $T_{\text{min}} = 0.92$, $T_{\text{max}} = 0.97$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$ 188 parameters
 $wR(F^2) = 0.096$ All H-atom parameters refined
 $S = 1.14$ $\Delta\rho_{\text{max}} = 0.35$ e Å⁻³
 2419 reflections $\Delta\rho_{\text{min}} = -0.16$ e Å⁻³

Table 1

Hydrogen-bond and $\text{C}-\text{H}\cdots\pi$ geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H}2\text{O}\cdots\text{O}1$	0.864 (16)	1.900 (16)	2.6547 (9)	145.1 (13)
$\text{O}3-\text{H}3\text{O}\cdots\text{O}2^{\text{ii}}$	0.909 (15)	1.870 (15)	2.7776 (9)	175.8 (13)
$\text{C}7-\text{H}7\text{A}\cdots\text{C}1^{\text{ii}}$	0.992 (10)	2.939 (10)	3.8346 (12)	150.7 (8)
$\text{C}7-\text{H}7\text{A}\cdots\text{C}2^{\text{ii}}$	0.992 (10)	3.103 (10)	4.0850 (12)	170.7 (8)
$\text{C}7-\text{H}7\text{A}\cdots\text{C}6^{\text{ii}}$	0.992 (10)	2.965 (10)	3.6436 (13)	126.5 (7)

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2005); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2005); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXL97*.

The X-ray measurements were undertaken in the Crystallographic Unit of the Physical Chemistry Laboratory at the Chemistry Department of the University of Warsaw. This work was supported by the Warsaw University of Technology and by the Polish Ministry of Science and Higher Education (grant No. N N205 055633). This work was supported by the Aldrich Chemical Company through the donation of chemicals.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2240).

References

- Brandenburg, K. (1999). *DIAMOND*. Version 2.1c. Crystal Impact GbR, Bonn, Germany.
 Dabrowski, M., Lulinski, S., Serwatowski, J. & Szczerbinska, M. (2006). *Acta Cryst. C* **62**, o702–o704.
 Oxford Diffraction (2005). *CrysAlis CCD* and *CrysAlis RED*. Versions 1.171. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
 Rettig, S. J. & Trotter, J. (1977). *Can. J. Chem.* **55**, 3071–3075.
 Serwatowski, J., Klis, T. & Kacprzak, K. (2006). *Acta Cryst. E* **62**, o1308–o1309.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
 Yang, Y., Escobedo, J. O., Wong, A., Schowalter, C. M., Touchy, M. C., Jiao, L., Crowe, W. E., Fronczek, F. R. & Strongin, R. M. (2005). *J. Org. Chem.* **70**, 6907–6912.

supplementary materials

Acta Cryst. (2008). E64, o437 [doi:10.1107/S1600536808000457]

(2-Butoxyphenyl)boronic acid

M. Dabrowski, S. Lulinski and J. Serwatowski

Comment

The boronic acid group is known to support supramolecular organization due to intermolecular hydrogen bonding. ortho-Substituents in the aryl ring may significantly influence the structural properties of arylboronic acids. There are a few structures of ortho-alkoxyarylboronic acids available in the literature, i.e. those reported by Yang et al. (2005), Serwatowski et al. (2006) and Dabrowski et al. (2006). We were interested in studying the effect of a longer alkoxy chain on the structural characteristics of the related compound, (I).

The molecular structure of (I) shows the entire molecule to be essentially planar, Fig. 1 & Table 1. The mean planes through the boronic and butoxy groups are approximately co-planar with the aromatic ring. The boronic group has an exo–endo conformation. The endo-oriented OH group forms an intramolecular O—H \cdots O bond with the butoxy-O atom, Table 2. As a result, a nearly planar six-membered ring is formed. This motif has been observed in the structures of related ortho-alkoxyarylboronic acids. Monomeric molecules form hydrogen-bonded centrosymmetric dimers typical of boronic acids (Rettig & Trotter, 1977). The crystal packing in (I) features a parallel arrangement of hydrogen-bonded dimers (Fig. 2). It is stabilized in terms of CH- π interactions between the H7a atom of the butoxy group and the aromatic ring of the adjacent molecule: the distance of H7A from the ring centre is 2.777 (11) Å [symmetry code (ii): 1 - x, 1 - y, 1 - z]. As a result, a centrosymmetric dimeric motif can be distinguished. In addition, weak π - π interactions between a pair of aromatic rings lead to their face-to-face center-to-edge stacking with the shortest contact between two C atoms C1—C1ⁱⁱⁱ = 3.540 (3) Å [symmetry code (iii): -x, 1 - y, 1 - z]. The other two π - π interactions are C1 \cdots C2ⁱⁱⁱ = 3.594 (5) Å and C1 \cdots C6ⁱⁱⁱ = 3.819 (3) Å. Thus, alternate CH- π and π - π interactions result in formation of a two-dimensional array. In conclusion, the hydrogen-bonded dimeric structure of (I) is typical of boronic acids whereas the unique secondary supramolecular assembly is achieved due to weaker CH- π and π - π interactions.

Experimental

Crystals suitable for the X-ray diffraction analysis were grown by slow evaporation of a solution of the acid (0.2 g) in acetone/water (20 ml, 1:1).

Refinement

All H atoms were located in difference syntheses and refined freely. The range of O—H distances = 0.864 (16) to 0.909 (15) Å and range of C—H distances = 0.954 (11) to 1.030 (10) Å.

Figures




Fig. 1. The molecular structure of (I). Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. The crystal packing in (I) showing hydrogen-bonding, C—H...π and π-π interactions as dashed lines.

(2-Butoxyphenyl)boronic acid

Crystal data

$C_{10}H_{15}BO_3$

$M_r = 194.03$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 7.4809$ (4) Å

$b = 15.3510$ (7) Å

$c = 9.2824$ (4) Å

$\beta = 94.299$ (4)°

$V = 1062.98$ (9) Å³

$Z = 4$

$F_{000} = 416$

$D_x = 1.212$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 7109 reflections

$\theta = 2.3$ – 30.0 °

$\mu = 0.09$ mm⁻¹

$T = 102$ (2) K

Prismatic, colourless

$0.74 \times 0.47 \times 0.32$ mm

Data collection

Kuma KM4 CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 8.6479 pixels mm⁻¹

$T = 102$ (2) K

ω scans

Absorption correction: multi-scan
(CrysAlis RED; Oxford Diffraction, 2005)

$T_{\min} = 0.92$, $T_{\max} = 0.97$

9363 measured reflections

2419 independent reflections

1924 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$

$\theta_{\text{max}} = 27.5$ °

$\theta_{\text{min}} = 2.7$ °

$h = -9 \rightarrow 9$

$k = -19 \rightarrow 19$

$l = -12 \rightarrow 11$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	All H-atom parameters refined
$R[F^2 > 2\sigma(F^2)] = 0.032$	$w = 1/[\sigma^2(F_o^2) + (0.0603P)^2]$
$wR(F^2) = 0.096$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.14$	$(\Delta/\sigma)_{\max} < 0.001$
2419 reflections	$\Delta\rho_{\max} = 0.35 \text{ e } \text{\AA}^{-3}$
188 parameters	$\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1+0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.015 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.22865 (11)	0.48929 (6)	0.46605 (10)	0.0187 (2)
C2	0.15354 (11)	0.42983 (6)	0.36299 (10)	0.0181 (2)
C3	0.12666 (12)	0.34464 (6)	0.41017 (11)	0.0213 (2)
C4	0.16761 (12)	0.31891 (6)	0.55234 (11)	0.0242 (2)
C5	0.23872 (12)	0.37970 (7)	0.65050 (10)	0.0244 (2)
C6	0.27140 (12)	0.46477 (6)	0.60865 (10)	0.0219 (2)
C7	0.32860 (12)	0.63751 (6)	0.51594 (10)	0.0196 (2)
C8	0.34388 (13)	0.72079 (6)	0.43222 (10)	0.0205 (2)
C9	0.42380 (14)	0.79405 (6)	0.52702 (11)	0.0236 (2)
C10	0.46998 (16)	0.87395 (7)	0.44072 (13)	0.0324 (3)
B1	0.09189 (13)	0.45600 (7)	0.20334 (11)	0.0189 (2)
O1	0.25505 (9)	0.57244 (4)	0.41655 (7)	0.02163 (19)
O2	0.10984 (9)	0.53922 (4)	0.15252 (7)	0.02530 (19)
O3	0.01624 (9)	0.39432 (4)	0.11431 (8)	0.0266 (2)
H2O	0.157 (2)	0.5714 (9)	0.2219 (17)	0.056 (4)*
H3O	-0.0232 (18)	0.4136 (9)	0.0250 (16)	0.055 (4)*

supplementary materials

H3	0.0761 (15)	0.3037 (7)	0.3414 (12)	0.027 (3)*
H4	0.1452 (14)	0.2583 (7)	0.5786 (12)	0.028 (3)*
H5	0.2692 (15)	0.3638 (8)	0.7519 (13)	0.034 (3)*
H6	0.3221 (14)	0.5074 (7)	0.6758 (12)	0.029 (3)*
H7A	0.4484 (14)	0.6175 (7)	0.5556 (10)	0.022 (3)*
H7B	0.2485 (13)	0.6452 (7)	0.5943 (11)	0.019 (2)*
H8A	0.4243 (13)	0.7100 (6)	0.3536 (11)	0.022 (3)*
H8B	0.2185 (14)	0.7378 (7)	0.3876 (12)	0.026 (3)*
H9A	0.5375 (14)	0.7721 (7)	0.5801 (12)	0.027 (3)*
H9B	0.3361 (16)	0.8101 (7)	0.5990 (12)	0.030 (3)*
H10A	0.5591 (18)	0.8578 (8)	0.3676 (15)	0.052 (4)*
H10B	0.3595 (18)	0.8990 (8)	0.3852 (14)	0.045 (3)*
H10C	0.5225 (16)	0.9206 (8)	0.5042 (13)	0.045 (4)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0164 (4)	0.0187 (5)	0.0210 (5)	0.0035 (3)	0.0007 (3)	0.0005 (4)
C2	0.0152 (4)	0.0192 (5)	0.0200 (4)	0.0025 (3)	0.0009 (3)	0.0001 (4)
C3	0.0193 (5)	0.0195 (5)	0.0251 (5)	0.0011 (4)	0.0010 (4)	-0.0008 (4)
C4	0.0241 (5)	0.0203 (5)	0.0284 (5)	0.0027 (4)	0.0032 (4)	0.0056 (4)
C5	0.0242 (5)	0.0281 (5)	0.0209 (5)	0.0060 (4)	0.0012 (4)	0.0053 (4)
C6	0.0229 (5)	0.0233 (5)	0.0190 (5)	0.0035 (4)	-0.0016 (4)	-0.0015 (4)
C7	0.0212 (5)	0.0204 (5)	0.0167 (4)	0.0003 (3)	-0.0026 (4)	-0.0041 (4)
C8	0.0223 (5)	0.0216 (5)	0.0171 (5)	0.0011 (4)	-0.0020 (4)	-0.0013 (4)
C9	0.0306 (5)	0.0193 (5)	0.0200 (5)	0.0018 (4)	-0.0031 (4)	-0.0021 (4)
C10	0.0397 (6)	0.0234 (5)	0.0332 (6)	-0.0038 (5)	-0.0032 (5)	0.0012 (5)
B1	0.0179 (5)	0.0184 (5)	0.0205 (5)	0.0009 (4)	0.0011 (4)	-0.0009 (4)
O1	0.0289 (4)	0.0173 (4)	0.0176 (3)	-0.0030 (3)	-0.0047 (3)	-0.0005 (3)
O2	0.0369 (4)	0.0203 (4)	0.0176 (4)	-0.0063 (3)	-0.0057 (3)	-0.0002 (3)
O3	0.0363 (4)	0.0194 (4)	0.0225 (4)	-0.0020 (3)	-0.0090 (3)	-0.0008 (3)

Geometric parameters (\AA , $^\circ$)

C1—O1	1.3758 (11)	C7—H7B	0.983 (10)
C1—C6	1.3902 (13)	C8—C9	1.5223 (13)
C1—C2	1.4083 (13)	C8—H8A	0.994 (10)
C2—C3	1.3984 (13)	C8—H8B	1.030 (10)
C2—B1	1.5710 (13)	C9—C10	1.5189 (14)
C3—C4	1.3895 (13)	C9—H9A	1.009 (11)
C3—H3	0.954 (11)	C9—H9B	1.001 (12)
C4—C5	1.3825 (14)	C10—H10A	1.017 (14)
C4—H4	0.980 (10)	C10—H10B	1.017 (13)
C5—C6	1.3895 (14)	C10—H10C	0.990 (13)
C5—H5	0.982 (12)	B1—O3	1.3526 (12)
C6—H6	0.962 (12)	B1—O2	1.3718 (12)
C7—O1	1.4408 (10)	O2—H2O	0.864 (16)
C7—C8	1.5050 (13)	O3—H3O	0.909 (15)
C7—H7A	0.992 (10)		

O1—C1—C6	122.72 (9)	C7—C8—C9	111.76 (7)
O1—C1—C2	115.75 (8)	C7—C8—H8A	108.1 (6)
C6—C1—C2	121.53 (9)	C9—C8—H8A	108.3 (6)
C3—C2—C1	116.92 (8)	C7—C8—H8B	108.7 (6)
C3—C2—B1	119.71 (8)	C9—C8—H8B	110.6 (6)
C1—C2—B1	123.30 (8)	H8A—C8—H8B	109.3 (8)
C4—C3—C2	122.46 (9)	C10—C9—C8	112.75 (8)
C4—C3—H3	119.8 (6)	C10—C9—H9A	108.1 (6)
C2—C3—H3	117.8 (6)	C8—C9—H9A	108.4 (6)
C5—C4—C3	118.72 (9)	C10—C9—H9B	109.8 (6)
C5—C4—H4	122.8 (6)	C8—C9—H9B	108.6 (6)
C3—C4—H4	118.5 (6)	H9A—C9—H9B	109.1 (9)
C4—C5—C6	121.15 (9)	C9—C10—H10A	110.0 (7)
C4—C5—H5	120.9 (7)	C9—C10—H10B	111.4 (7)
C6—C5—H5	118.0 (7)	H10A—C10—H10B	107.6 (11)
C5—C6—C1	119.20 (9)	C9—C10—H10C	111.5 (7)
C5—C6—H6	121.8 (7)	H10A—C10—H10C	108.8 (10)
C1—C6—H6	119.0 (7)	H10B—C10—H10C	107.4 (10)
O1—C7—C8	107.37 (7)	O3—B1—O2	119.28 (8)
O1—C7—H7A	108.3 (6)	O3—B1—C2	118.50 (8)
C8—C7—H7A	110.7 (6)	O2—B1—C2	122.22 (8)
O1—C7—H7B	109.4 (6)	C1—O1—C7	119.14 (7)
C8—C7—H7B	110.7 (6)	B1—O2—H2O	108.7 (10)
H7A—C7—H7B	110.3 (8)	B1—O3—H3O	114.9 (9)
O1—C1—C2—C3	179.49 (7)	C2—C1—C6—C5	-0.21 (14)
C6—C1—C2—C3	-1.00 (13)	O1—C7—C8—C9	-179.00 (7)
O1—C1—C2—B1	-3.50 (13)	C7—C8—C9—C10	170.24 (9)
C6—C1—C2—B1	176.01 (8)	C3—C2—B1—O3	-1.04 (13)
C1—C2—C3—C4	1.34 (13)	C1—C2—B1—O3	-177.97 (8)
B1—C2—C3—C4	-175.78 (8)	C3—C2—B1—O2	177.95 (8)
C2—C3—C4—C5	-0.45 (14)	C1—C2—B1—O2	1.02 (14)
C3—C4—C5—C6	-0.84 (14)	C6—C1—O1—C7	-0.53 (12)
C4—C5—C6—C1	1.16 (14)	C2—C1—O1—C7	178.98 (8)
O1—C1—C6—C5	179.27 (8)	C8—C7—O1—C1	179.46 (7)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O2—H2O...O1	0.864 (16)	1.900 (16)	2.6547 (9)	145.1 (13)
O3—H3O...O2 ⁱ	0.909 (15)	1.870 (15)	2.7776 (9)	175.8 (13)
C7—H7A...C1 ⁱⁱ	0.992 (10)	2.939 (10)	3.8346 (12)	150.7 (8)
C7—H7A...C2 ⁱⁱ	0.992 (10)	3.103 (10)	4.0850 (12)	170.7 (8)
C7—H7A...C6 ⁱⁱ	0.992 (10)	2.965 (10)	3.6436 (13)	126.5 (7)

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

