

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5-(4-Bromobenzylidene)-5H-dibenzo-
[a,d][7]annulene

Ren-Hua Zheng

School of Pharmaceutical and Chemical Engineering, Taizhou University, Linhai 317000, People's Republic of China
Correspondence e-mail: zrh1977@126.com


Received 15 January 2008; accepted 3 February 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.054; wR factor = 0.141; data-to-parameter ratio = 18.1.

The tricyclic system of the title compound, $\text{C}_{22}\text{H}_{15}\text{Br}$, has a concave shape, with a dihedral angle between the benzene ring planes of 48.68 (1)°.

Related literature

For related literature, see: Allen *et al.* (1987); Bergmann & Solomonovici (1970); Larson (1970).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{15}\text{Br}$
 $M_r = 359.26$
Monoclinic, $P2_1/c$

$a = 8.4857$ (5) Å
 $b = 19.0479$ (8) Å
 $c = 10.6808$ (5) Å

$\beta = 104.6802$ (16)°
 $V = 1670.03$ (14) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 2.46$ mm⁻¹
 $T = 296$ (1) K
 $0.57 \times 0.46 \times 0.29$ mm

Data collection

Rigaku R-Axis RAPID diffractometer
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.227$, $T_{\max} = 0.489$

8896 measured reflections
3792 independent reflections
2462 reflections with $F^2 > 2\sigma(F^2)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.140$
 $S = 1.00$
3792 reflections

209 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.83$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.93$ e Å⁻³

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2004); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *CrystalStructure*.

The author acknowledges the help of Professor Jian-Ming Gu of Zhejiang University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2131).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
Bergmann, D. E. & Solomonovici, A. (1970). *Synthesis*, **2**, 183–189.
Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Larson, A. C. (1970). *Crystallographic Computing*, edited by F. R. Ahmed, S. R. Hall & C. P. Huber, pp. 291–294. Copenhagen: Munksgaard.
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MS (2004). *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.

supplementary materials

Acta Cryst. (2008). E64, o589 [doi:10.1107/S1600536808003711]

5-(4-Bromobenzylidene)-5*H*-dibenzo[*a,d*][7]annulene

R.-H. Zheng

Comment

The title compound was synthesized through Wittig–Horner reaction (Bergmann & Solomonovici, 1970).

The molecular structure is shown in Fig. 1. The bond lengths and angles are generally within normal ranges (Allen *et al.*, 1987). Packing diagram is given in Fig. 2. The seven-membered ring is in a boat conformation. The dihedral angle between the benzene A (C1–C6) and the plane defined by the atoms C9/C14/C17/C22 is 29.5 (1)°. Benzene C (C9–C14) and benzene D (C17–C22) form with the plane defined by C9/C14/C17/C22 dihedral angles of 24.1 (1)° and 26.4 (1)°, respectively, while the dihedral angle between them is 48.68 (1)°. The crystal packing (Fig. 2) is mainly stabilized by van der Waals forces.

Experimental

The title compound was synthesized by treating solution of (4-bromo-benzyl)-phosphonic acid diethyl ester (1.53 g, 5 mmol) and dibenzo[*a,d*]cyclohepten-5-one (1.03 g, 5 mmol) in 100 ml anhydrous THF under nitrogen with solid potassium *tert*-butoxide (1.68 g, 15 mmol) which was added in one portion. The mixture was refluxed with stirring for 10 h. Solvent was removed by rotary evaporation. The residue was purified by column chromatography (silica gel) using *n*-hexane as eluent. Colorless crystals were obtained by slow evaporation of a dichloromethane solution.

Refinement

The H atoms were positioned geometrically and allowed to ride on their parent atoms, with C—H = 0.93 Å and $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Molecular structure of of the title compound with displacement ellipsoids shown at the 50% probability level.




Fig. 2. Crystal packing diagram of the title compound.

5-(4-Bromobenzylidene)-5*H*-dibenzo[*a,d*][7]annulene

Crystal data

$C_{22}H_{15}Br$	$F_{000} = 728.00$
$M_r = 359.26$	$D_x = 1.429 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71075 \text{ \AA}$
$a = 8.4857 (5) \text{ \AA}$	Cell parameters from 8966 reflections
$b = 19.0479 (8) \text{ \AA}$	$\theta = 3.2\text{--}27.4^\circ$
$c = 10.6808 (5) \text{ \AA}$	$\mu = 2.47 \text{ mm}^{-1}$
$\beta = 104.6802 (16)^\circ$	$T = 296 (1) \text{ K}$
$V = 1670.03 (14) \text{ \AA}^3$	Chunk, colourless
$Z = 4$	$0.57 \times 0.46 \times 0.29 \text{ mm}$

Data collection

Rigaku R-AXIS RAPID diffractometer	2462 reflections with $F^2 > 2\sigma(F^2)$
Detector resolution: $10.00 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.051$
ω scans	$\theta_{\text{max}} = 27.4^\circ$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$h = 0 \rightarrow 10$
$T_{\text{min}} = 0.227$, $T_{\text{max}} = 0.489$	$k = 0 \rightarrow 24$
8896 measured reflections	$l = -13 \rightarrow 13$
3792 independent reflections	

Refinement

Refinement on F^2	$w = 1/[0.001F_o^2 + 6.2\sigma(F_o^2)]/(4F_o^2)$
$R[F^2 > 2\sigma(F^2)] = 0.053$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$wR(F^2) = 0.140$	$\Delta\rho_{\text{max}} = 0.83 \text{ e \AA}^{-3}$
$S = 1.01$	$\Delta\rho_{\text{min}} = -0.93 \text{ e \AA}^{-3}$
3792 reflections	Extinction correction: Larson (1970)
209 parameters	Extinction coefficient: 351 (32)
H-atom parameters constrained	

Special details

Refinement. Refinement using all reflections. The weighted R -factor (wR) and goodness of fit (S) are based on F^2 . R -factor (gt) are based on F . The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating R -factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
-----	-----	-----	----------------------------------

Br1	0.09761 (6)	0.48750 (2)	-0.21051 (4)	0.08526 (18)
C1	0.2486 (4)	0.54224 (18)	-0.0874 (3)	0.0574 (10)
C2	0.2994 (4)	0.51963 (18)	0.0385 (3)	0.0618 (11)
C3	0.4098 (4)	0.55959 (17)	0.1270 (3)	0.0586 (11)
C4	0.4709 (4)	0.62248 (17)	0.0924 (2)	0.0482 (9)
C5	0.4180 (4)	0.64341 (19)	-0.0356 (3)	0.0588 (11)
C6	0.3069 (4)	0.60356 (19)	-0.1258 (3)	0.0616 (11)
C7	0.5914 (4)	0.66577 (17)	0.1840 (3)	0.0516 (10)
C8	0.6033 (4)	0.67805 (17)	0.3102 (2)	0.0481 (9)
C9	0.7365 (3)	0.72298 (16)	0.3866 (2)	0.0450 (9)
C10	0.7624 (4)	0.78939 (18)	0.3398 (3)	0.0561 (10)
C11	0.8893 (4)	0.83153 (19)	0.4062 (3)	0.0612 (11)
C12	0.9902 (4)	0.8091 (2)	0.5201 (3)	0.0654 (12)
C13	0.9669 (4)	0.7436 (2)	0.5676 (3)	0.0618 (11)
C14	0.8397 (4)	0.69947 (17)	0.5039 (3)	0.0502 (9)
C15	0.8235 (4)	0.63084 (18)	0.5605 (3)	0.0560 (10)
C16	0.6901 (4)	0.59282 (18)	0.5526 (3)	0.0573 (10)
C17	0.5221 (4)	0.61096 (17)	0.4876 (2)	0.0493 (9)
C18	0.3957 (5)	0.58558 (19)	0.5374 (3)	0.0643 (12)
C19	0.2344 (5)	0.6034 (2)	0.4844 (4)	0.0720 (14)
C20	0.1960 (5)	0.6485 (2)	0.3806 (3)	0.0673 (13)
C21	0.3170 (4)	0.67289 (18)	0.3276 (3)	0.0550 (10)
C22	0.4800 (4)	0.65368 (14)	0.3770 (2)	0.0459 (9)
H2	0.2599	0.4778	0.0637	0.074*
H3	0.4444	0.5441	0.2121	0.070*
H5	0.4576	0.6850	-0.0618	0.071*
H6	0.2725	0.6183	-0.2114	0.074*
H7	0.6696	0.6875	0.1501	0.062*
H10	0.6935	0.8055	0.2630	0.067*
H11	0.9060	0.8753	0.3731	0.073*
H12	1.0740	0.8378	0.5654	0.078*
H13	1.0377	0.7283	0.6441	0.074*
H15	0.9200	0.6110	0.6087	0.067*
H16	0.7057	0.5493	0.5935	0.069*
H18	0.4209	0.5557	0.6086	0.077*
H19	0.1528	0.5850	0.5187	0.086*
H20	0.0886	0.6624	0.3465	0.081*
H21	0.2898	0.7030	0.2568	0.066*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0863 (4)	0.0690 (3)	0.0828 (3)	-0.0041 (2)	-0.0113 (2)	-0.0165 (2)
C1	0.061 (2)	0.0547 (19)	0.057 (2)	0.0027 (17)	0.0150 (18)	-0.0078 (16)
C2	0.075 (2)	0.0516 (19)	0.059 (2)	-0.0067 (18)	0.0163 (19)	-0.0031 (16)
C3	0.072 (2)	0.058 (2)	0.0442 (19)	-0.0005 (18)	0.0113 (18)	0.0038 (15)
C4	0.047 (2)	0.0511 (17)	0.0476 (18)	-0.0003 (15)	0.0138 (15)	-0.0006 (14)
C5	0.065 (2)	0.060 (2)	0.050 (2)	-0.0063 (18)	0.0130 (18)	0.0094 (15)

supplementary materials

C6	0.068 (2)	0.067 (2)	0.046 (2)	0.0006 (19)	0.0055 (18)	-0.0010 (16)
C7	0.047 (2)	0.0579 (19)	0.0488 (19)	-0.0013 (15)	0.0106 (16)	0.0042 (15)
C8	0.045 (2)	0.0513 (18)	0.0462 (19)	0.0033 (14)	0.0092 (15)	0.0039 (14)
C9	0.0390 (18)	0.0527 (17)	0.0451 (17)	-0.0027 (14)	0.0137 (14)	-0.0061 (14)
C10	0.054 (2)	0.062 (2)	0.053 (2)	-0.0040 (17)	0.0153 (17)	0.0009 (16)
C11	0.059 (2)	0.059 (2)	0.067 (2)	-0.0109 (18)	0.0178 (19)	-0.0059 (17)
C12	0.054 (2)	0.072 (2)	0.069 (2)	-0.0145 (18)	0.012 (2)	-0.0151 (19)
C13	0.046 (2)	0.076 (2)	0.057 (2)	0.0030 (18)	0.0017 (17)	-0.0032 (18)
C14	0.0395 (19)	0.0592 (19)	0.0508 (19)	0.0018 (15)	0.0095 (15)	-0.0089 (15)
C15	0.052 (2)	0.058 (2)	0.054 (2)	0.0050 (17)	0.0062 (17)	0.0034 (16)
C16	0.068 (2)	0.0480 (18)	0.053 (2)	0.0076 (17)	0.0106 (18)	0.0045 (14)
C17	0.056 (2)	0.0462 (17)	0.0474 (19)	-0.0015 (15)	0.0160 (17)	-0.0048 (13)
C18	0.074 (2)	0.065 (2)	0.060 (2)	-0.005 (2)	0.028 (2)	0.0046 (17)
C19	0.065 (2)	0.086 (2)	0.075 (2)	-0.017 (2)	0.036 (2)	-0.012 (2)
C20	0.051 (2)	0.084 (2)	0.069 (2)	0.002 (2)	0.021 (2)	-0.016 (2)
C21	0.050 (2)	0.062 (2)	0.053 (2)	0.0002 (17)	0.0117 (17)	-0.0066 (16)
C22	0.046 (2)	0.0461 (17)	0.0466 (17)	-0.0028 (14)	0.0123 (15)	-0.0064 (13)

Geometric parameters (Å, °)

Br1—C1	1.899 (3)	C17—C22	1.404 (4)
C1—C2	1.372 (4)	C18—C19	1.384 (5)
C1—C6	1.371 (5)	C19—C20	1.375 (5)
C2—C3	1.379 (4)	C20—C21	1.374 (6)
C3—C4	1.392 (4)	C21—C22	1.399 (4)
C4—C5	1.386 (4)	C2—H2	0.930
C4—C7	1.475 (4)	C3—H3	0.930
C5—C6	1.390 (4)	C5—H5	0.930
C7—C8	1.346 (4)	C6—H6	0.930
C8—C9	1.486 (4)	C7—H7	0.930
C8—C22	1.482 (5)	C10—H10	0.930
C9—C10	1.398 (4)	C11—H11	0.930
C9—C14	1.408 (4)	C12—H12	0.930
C10—C11	1.386 (4)	C13—H13	0.930
C11—C12	1.367 (4)	C15—H15	0.930
C12—C13	1.378 (5)	C16—H16	0.930
C13—C14	1.401 (4)	C18—H18	0.930
C14—C15	1.461 (4)	C19—H19	0.930
C15—C16	1.328 (5)	C20—H20	0.930
C16—C17	1.461 (4)	C21—H21	0.930
C17—C18	1.398 (6)		
Br1—C1—C2	119.7 (2)	C8—C22—C17	121.8 (2)
Br1—C1—C6	119.4 (2)	C8—C22—C21	119.2 (2)
C2—C1—C6	120.9 (3)	C17—C22—C21	119.1 (3)
C1—C2—C3	119.2 (3)	C1—C2—H2	120.4
C2—C3—C4	121.8 (3)	C3—C2—H2	120.4
C3—C4—C5	117.4 (2)	C2—C3—H3	119.1
C3—C4—C7	123.2 (2)	C4—C3—H3	119.1
C5—C4—C7	119.4 (3)	C4—C5—H5	119.3

C4—C5—C6	121.3 (3)	C6—C5—H5	119.3
C1—C6—C5	119.3 (3)	C1—C6—H6	120.3
C4—C7—C8	128.7 (3)	C5—C6—H6	120.3
C7—C8—C9	120.2 (3)	C4—C7—H7	115.7
C7—C8—C22	123.0 (2)	C8—C7—H7	115.7
C9—C8—C22	116.6 (2)	C9—C10—H10	119.6
C8—C9—C10	119.4 (2)	C11—C10—H10	119.6
C8—C9—C14	121.4 (2)	C10—C11—H11	119.9
C10—C9—C14	119.2 (2)	C12—C11—H11	119.9
C9—C10—C11	120.9 (2)	C11—C12—H12	120.2
C10—C11—C12	120.3 (3)	C13—C12—H12	120.2
C11—C12—C13	119.7 (3)	C12—C13—H13	119.0
C12—C13—C14	121.9 (3)	C14—C13—H13	119.0
C9—C14—C13	118.1 (3)	C14—C15—H15	115.5
C9—C14—C15	123.4 (2)	C16—C15—H15	115.5
C13—C14—C15	118.5 (2)	C15—C16—H16	116.0
C14—C15—C16	129.0 (3)	C17—C16—H16	116.0
C15—C16—C17	127.9 (3)	C17—C18—H18	118.9
C16—C17—C18	119.2 (3)	C19—C18—H18	118.9
C16—C17—C22	123.1 (3)	C18—C19—H19	120.3
C18—C17—C22	117.7 (3)	C20—C19—H19	120.3
C17—C18—C19	122.2 (3)	C19—C20—H20	120.2
C18—C19—C20	119.4 (4)	C21—C20—H20	120.2
C19—C20—C21	119.7 (3)	C20—C21—H21	119.1
C20—C21—C22	121.8 (3)	C22—C21—H21	119.1
Br1—C1—C2—C3	179.7 (3)	C10—C9—C14—C13	-1.3 (5)
Br1—C1—C6—C5	-179.7 (3)	C10—C9—C14—C15	-179.6 (3)
C2—C1—C6—C5	-0.5 (6)	C14—C9—C10—C11	1.0 (5)
C6—C1—C2—C3	0.4 (6)	C9—C10—C11—C12	-1.0 (6)
C1—C2—C3—C4	0.2 (5)	C10—C11—C12—C13	1.2 (6)
C2—C3—C4—C5	-0.7 (5)	C11—C12—C13—C14	-1.5 (6)
C2—C3—C4—C7	-178.8 (3)	C12—C13—C14—C9	1.5 (5)
C3—C4—C5—C6	0.7 (5)	C12—C13—C14—C15	180.0 (2)
C3—C4—C7—C8	-37.4 (5)	C9—C14—C15—C16	-30.8 (6)
C5—C4—C7—C8	144.6 (3)	C13—C14—C15—C16	150.9 (4)
C7—C4—C5—C6	178.8 (3)	C14—C15—C16—C17	-2.0 (6)
C4—C5—C6—C1	-0.1 (4)	C15—C16—C17—C18	-148.6 (3)
C4—C7—C8—C9	179.2 (3)	C15—C16—C17—C22	30.7 (5)
C4—C7—C8—C22	-6.3 (5)	C16—C17—C18—C19	176.9 (3)
C7—C8—C9—C10	51.4 (4)	C16—C17—C22—C8	6.1 (4)
C7—C8—C9—C14	-127.2 (3)	C16—C17—C22—C21	-174.9 (3)
C7—C8—C22—C17	123.5 (3)	C18—C17—C22—C8	-174.6 (3)
C7—C8—C22—C21	-55.6 (4)	C18—C17—C22—C21	4.5 (4)
C9—C8—C22—C17	-61.8 (4)	C22—C17—C18—C19	-2.5 (5)
C9—C8—C22—C21	119.2 (3)	C17—C18—C19—C20	-1.2 (6)
C22—C8—C9—C10	-123.5 (3)	C18—C19—C20—C21	2.8 (6)
C22—C8—C9—C14	57.9 (4)	C19—C20—C21—C22	-0.7 (5)
C8—C9—C10—C11	-177.6 (3)	C20—C21—C22—C8	176.0 (3)
C8—C9—C14—C13	177.3 (3)	C20—C21—C22—C17	-3.1 (4)

supplementary materials

C8—C9—C14—C15

-1.0 (5)

Fig. 1




Fig. 2

