

## 2,3-Bis(2-chlorobenzyl)naphthalene-1,4-dione

Shiliang Huang,<sup>a,\*</sup> Zhi-Shu Huang,<sup>a</sup> Lian-Quan Gu,<sup>a</sup>  
Zhongyuan Zhou<sup>b</sup> and Seik Weng Ng<sup>c</sup>

<sup>a</sup>School of Pharmaceutical Science, Sun Yat-Sen University, Guangzhou, People's Republic of China, <sup>b</sup>Department of Applied Biology and Chemical Technology, Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong, and <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: lsshsl@mail.sysu.edu.cn


Received 21 November 2007; accepted 21 December 2007

Key indicators: single-crystal X-ray study;  $T = 295\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  
 $R$  factor = 0.045;  $wR$  factor = 0.131; data-to-parameter ratio = 17.3.

The title disubstituted naphthalene-1,4-dione,  $C_{24}H_{16}Cl_2O_2$ , has the two chlorobenzyl substituents related by a non-crystallographic twofold rotation axis, generating a chiral conformation; both enantiomers are present. The two chlorobenzene rings are nearly perpendicular to the fused ring system, making angles of  $88.8(1)$  and  $77.5(1)^\circ$  with it.

### Related literature

2,3-Bis(2-chlorobenzyl)naphthalene-1,4-dione was the unexpected product in the attempted synthesis of 2-(2-chlorobenzyl)naphthalene-1,4-dione by a free-radical alkylation (see Boehm *et al.*, 1981; Chen *et al.*, 2005; Tsai *et al.*, 2001). Although the title dichloro compound has not been reported, 2,3-dibenzylnaphthalene-1,4-dione has been known for a long time (Baxter & Sanders, 1974; Chen *et al.*, 2005; Oettmeier *et al.*, 1986; Sharma & Torsell, 1978; Yamago *et al.*, 2002). This class of compounds exhibits antitumour activity (Driscoll, 1974; Driscoll *et al.*, 1974).



### Experimental

#### Crystal data

$C_{24}H_{16}Cl_2O_2$	$V = 1907.1(3)\text{ \AA}^3$
$M_r = 407.27$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 9.998(1)\text{ \AA}$	$\mu = 0.36\text{ mm}^{-1}$
$b = 10.272(1)\text{ \AA}$	$T = 295(2)\text{ K}$
$c = 18.804(2)\text{ \AA}$	$0.5 \times 0.5 \times 0.5\text{ mm}$
$\beta = 99.071(1)^\circ$	

#### Data collection

Bruker SMART area-detector diffractometer	4366 independent reflections
Absorption correction: none	2921 reflections with $I > 2\sigma(I)$
12440 measured reflections	$R_{\text{int}} = 0.035$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$	253 parameters
$wR(F^2) = 0.131$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.18\text{ e \AA}^{-3}$
4366 reflections	$\Delta\rho_{\text{min}} = -0.42\text{ e \AA}^{-3}$

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2008).

The authors are indebted to the National Natural Science Foundation of China (grant No. 30701050), the Guangdong Provincial Science Foundation (grant No. 7301303) and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OB2105).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–201.
- Baxter, I. & Sanders, J. K. M. (1974). *J. Chem. Soc. Chem. Commun.* pp. 143–144.
- Boehm, P., Cooper, K., Hudson, A. T., Elphick, J. P. & McHardy, N. (1981). *J. Med. Chem.* **24**, 295–299.
- Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, H.-L., Lin, C.-Y., Cheng, Y.-C., Tsai, A. I. & Chuang, C.-P. (2005). *Synthesis*, pp. 977–985.
- Driscoll, J. S. (1974). *Cancer Chemother. Rep.* **4**, 3–4.
- Driscoll, J. S., Hazard, G. F., Wood, H. B. & Goldin, A. (1974). *Cancer Chemother. Rep.* **4**, 1–27.
- Oettmeier, W., Dierig, C. & Masson, K. (1986). *Quant. Struct.-Act. Rel.* **5**, 50–54.
- Sharma, S. C. & Torsell, K. (1978). *Acta Chem. Scand.* **B32**, 347–353.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Tsai, A.-I., Wu, Y.-L. & Chuang, C.-P. (2001). *Tetrahedron*, **57**, 7829–7837.
- Westrip, S. P. (2008). *publCIF*. In preparation.
- Yamago, S., Hashidume, M. & Yoshida, J.-I. (2002). *Tetrahedron*, **58**, 6805–6813.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o593 [doi:10.1107/S1600536807068006]

## 2,3-Bis(2-chlorobenzyl)naphthalene-1,4-dione

S. Huang, Z.-S. Huang, L.-Q. Gu, Z. Zhou and S. W. Ng

### Experimental

The title compound (**I**) was synthesized by using a modification of the procedure by Boehm *et al.* (1981). 1,4-Naphthoquinone (4 g, 25.3 mmol), 2-(2-chlorophenyl)acetic acid (8.60 g, 50.6 mmol), and silver nitrate (2.13 g, 12.6 mmol) in acetonitrile (60 ml) were heated to 353 K. A solution of ammonium peroxydisulfate in 30 ml water was added dropwise within 30 min. The mixture was then refluxed for another 2 h. The solution was concentrated to give a brown solid. The crude product was purified by column chromatography (silica gel, 1:2 dichloromethane:petroleum ether) to give yellow solid in 40% yield. Single crystals of (**I**) were obtained by slow evaporation of a 1:2 dichloromethane:petroleum ether (50 ml) solution.

### Figures




Fig. 1. The molecular structure of (**I**) with 50% probability ellipsoids for the non-H atoms.

## 2,3-Bis(2-chlorobenzyl)naphthalene-1,4-dione

### Crystal data

C <sub>24</sub> H <sub>16</sub> Cl <sub>2</sub> O <sub>2</sub>	<i>F</i> <sub>000</sub> = 840
<i>M<sub>r</sub></i> = 407.27	<i>D<sub>x</sub></i> = 1.418 Mg m <sup>-3</sup>
Monoclinic, <i>P</i> 2 <sub>1</sub> /n	Mo <i>K</i> α radiation
Hall symbol: -P 2yn	$\lambda$ = 0.71073 Å
<i>a</i> = 9.998 (1) Å	Cell parameters from 6278 reflections
<i>b</i> = 10.272 (1) Å	$\theta$ = 2.2–27.5°
<i>c</i> = 18.804 (2) Å	$\mu$ = 0.36 mm <sup>-1</sup>
$\beta$ = 99.071 (1)°	<i>T</i> = 295 (2) K
<i>V</i> = 1907.1 (3) Å <sup>3</sup>	Block, yellow
<i>Z</i> = 4	0.5 × 0.5 × 0.5 mm

### Data collection

Bruker SMART area-detector 2921 reflections with  $I > 2\sigma(I)$

# supplementary materials

---

diffractometer

Radiation source: medium-focus sealed tube  $R_{\text{int}} = 0.035$

Monochromator: graphite  $\theta_{\text{max}} = 27.5^\circ$

$T = 295(2)$  K  $\theta_{\text{min}} = 2.2^\circ$

$\varphi$  and  $\omega$  scans  $h = -12 \rightarrow 12$

Absorption correction: none  $k = -13 \rightarrow 13$

12440 measured reflections  $l = -24 \rightarrow 11$

4366 independent reflections

## Refinement

Refinement on  $F^2$

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.044$

H-atom parameters constrained

$wR(F^2) = 0.131$

$$w = 1/\sigma^2(F_o^2) + (0.0628P)^2 + 0.3578P \\ \text{where } P = (F_o^2 + 2F_c^2)/3$$

$S = 1.02$

$$(\Delta/\sigma)_{\text{max}} = 0.001$$

4366 reflections

$$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$$

253 parameters

$$\Delta\rho_{\text{min}} = -0.42 \text{ e \AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.53007 (6)	0.55702 (6)	0.66580 (4)	0.0779 (2)
Cl2	-0.07641 (7)	0.55387 (6)	0.39047 (4)	0.0856 (2)
O1	0.48173 (15)	1.00324 (15)	0.60292 (8)	0.0693 (4)
O2	0.03492 (16)	1.00716 (16)	0.40921 (9)	0.0763 (5)
C1	0.39509 (19)	0.64277 (17)	0.69086 (10)	0.0508 (5)
C2	0.3610 (3)	0.6185 (2)	0.75798 (11)	0.0707 (7)
H2	0.4108	0.5589	0.7887	0.085*
C3	0.2548 (3)	0.6819 (2)	0.77899 (13)	0.0821 (8)
H3	0.2317	0.6658	0.8242	0.099*
C4	0.1817 (3)	0.7694 (2)	0.73391 (13)	0.0743 (7)
H4	0.1081	0.8118	0.7481	0.089*
C5	0.2169 (2)	0.79478 (19)	0.66737 (10)	0.0550 (5)
H5	0.1667	0.8550	0.6373	0.066*
C6	0.32481 (18)	0.73306 (16)	0.64433 (9)	0.0438 (4)
C7	0.3679 (2)	0.76083 (18)	0.57207 (10)	0.0524 (5)
H7A	0.4658	0.7688	0.5789	0.063*
H7B	0.3431	0.6871	0.5405	0.063*
C8	0.30674 (17)	0.88202 (17)	0.53547 (9)	0.0426 (4)
C9	0.37930 (17)	1.00544 (18)	0.55815 (9)	0.0438 (4)
C10	0.32835 (17)	1.12940 (16)	0.52474 (9)	0.0422 (4)
C11	0.39225 (19)	1.24608 (19)	0.54606 (11)	0.0555 (5)
H11	0.4650	1.2469	0.5836	0.067*

C12	0.3487 (2)	1.3603 (2)	0.51213 (14)	0.0698 (6)
H12	0.3920	1.4382	0.5266	0.084*
C13	0.2408 (2)	1.3596 (2)	0.45657 (15)	0.0745 (7)
H13	0.2131	1.4368	0.4330	0.089*
C14	0.1739 (2)	1.2458 (2)	0.43582 (12)	0.0618 (5)
H14	0.1002	1.2465	0.3988	0.074*
C15	0.21608 (18)	1.12966 (17)	0.47002 (9)	0.0447 (4)
C16	0.14175 (18)	1.00724 (18)	0.45086 (9)	0.0476 (4)
C17	0.19512 (18)	0.88289 (17)	0.48466 (9)	0.0443 (4)
C18	0.1139 (2)	0.76407 (19)	0.45892 (11)	0.0566 (5)
H18A	0.0185	0.7835	0.4574	0.068*
H18B	0.1372	0.6946	0.4936	0.068*
C19	0.13550 (17)	0.71606 (17)	0.38532 (10)	0.0466 (4)
C20	0.23605 (19)	0.76496 (19)	0.35008 (10)	0.0542 (5)
H20	0.2938	0.8290	0.3724	0.065*
C21	0.2529 (2)	0.7209 (2)	0.28236 (12)	0.0675 (6)
H21	0.3211	0.7555	0.2597	0.081*
C22	0.1689 (3)	0.6263 (3)	0.24877 (13)	0.0762 (7)
H22	0.1804	0.5965	0.2034	0.091*
C23	0.0682 (2)	0.5759 (2)	0.28197 (13)	0.0694 (6)
H23	0.0108	0.5121	0.2592	0.083*
C24	0.05248 (19)	0.62039 (18)	0.34967 (11)	0.0553 (5)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0746 (4)	0.0631 (4)	0.0901 (5)	0.0227 (3)	-0.0056 (3)	0.0124 (3)
Cl2	0.0785 (4)	0.0736 (4)	0.1006 (5)	-0.0274 (3)	0.0012 (3)	0.0022 (3)
O1	0.0628 (9)	0.0722 (10)	0.0629 (9)	0.0011 (7)	-0.0211 (7)	0.0104 (7)
O2	0.0696 (9)	0.0745 (10)	0.0715 (10)	0.0136 (8)	-0.0305 (8)	-0.0104 (8)
C1	0.0583 (11)	0.0404 (9)	0.0490 (10)	-0.0050 (8)	-0.0064 (9)	0.0029 (8)
C2	0.1079 (19)	0.0510 (12)	0.0466 (12)	-0.0113 (12)	-0.0080 (12)	0.0110 (10)
C3	0.134 (2)	0.0653 (15)	0.0519 (13)	-0.0148 (16)	0.0309 (14)	0.0074 (11)
C4	0.0997 (18)	0.0636 (14)	0.0682 (14)	-0.0003 (13)	0.0402 (13)	0.0038 (11)
C5	0.0640 (12)	0.0510 (11)	0.0520 (11)	0.0054 (9)	0.0157 (9)	0.0062 (9)
C6	0.0519 (10)	0.0376 (8)	0.0399 (9)	-0.0032 (7)	0.0012 (7)	0.0009 (7)
C7	0.0638 (11)	0.0465 (10)	0.0479 (10)	0.0142 (9)	0.0118 (9)	0.0056 (8)
C8	0.0507 (10)	0.0447 (10)	0.0333 (8)	0.0093 (8)	0.0100 (7)	0.0031 (7)
C9	0.0442 (9)	0.0532 (10)	0.0332 (8)	0.0052 (8)	0.0037 (7)	0.0011 (7)
C10	0.0453 (9)	0.0439 (9)	0.0385 (9)	0.0045 (7)	0.0102 (7)	-0.0013 (7)
C11	0.0516 (10)	0.0549 (11)	0.0603 (12)	-0.0011 (9)	0.0097 (9)	-0.0046 (9)
C12	0.0709 (14)	0.0443 (11)	0.0967 (18)	-0.0021 (10)	0.0214 (13)	-0.0018 (11)
C13	0.0813 (15)	0.0472 (12)	0.0973 (18)	0.0131 (11)	0.0211 (14)	0.0212 (12)
C14	0.0621 (12)	0.0606 (13)	0.0610 (12)	0.0173 (10)	0.0050 (10)	0.0161 (10)
C15	0.0478 (9)	0.0467 (10)	0.0397 (9)	0.0097 (8)	0.0073 (7)	0.0034 (7)
C16	0.0505 (10)	0.0539 (11)	0.0359 (9)	0.0089 (8)	-0.0009 (8)	-0.0028 (8)
C17	0.0509 (10)	0.0465 (10)	0.0365 (9)	0.0017 (8)	0.0097 (7)	-0.0027 (7)
C18	0.0641 (12)	0.0530 (11)	0.0531 (11)	-0.0076 (9)	0.0101 (9)	-0.0038 (9)

## supplementary materials

---

C19	0.0468 (9)	0.0418 (9)	0.0476 (10)	0.0059 (8)	-0.0036 (8)	-0.0018 (8)
C20	0.0497 (10)	0.0586 (11)	0.0514 (11)	0.0030 (9)	-0.0007 (9)	-0.0095 (9)
C21	0.0657 (13)	0.0817 (16)	0.0549 (12)	0.0107 (12)	0.0092 (10)	-0.0092 (11)
C22	0.0836 (16)	0.0850 (17)	0.0555 (13)	0.0203 (14)	-0.0027 (12)	-0.0212 (12)
C23	0.0766 (15)	0.0565 (12)	0.0658 (14)	0.0061 (11)	-0.0176 (12)	-0.0163 (11)
C24	0.0538 (11)	0.0436 (10)	0.0626 (12)	0.0045 (8)	-0.0089 (9)	-0.0003 (9)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C11—C1	1.738 (2)	C11—H11	0.9300
C12—C24	1.740 (2)	C12—C13	1.379 (3)
O1—C9	1.219 (2)	C12—H12	0.9300
O2—C16	1.221 (2)	C13—C14	1.372 (3)
C1—C2	1.381 (3)	C13—H13	0.9300
C1—C6	1.388 (2)	C14—C15	1.389 (3)
C2—C3	1.357 (3)	C14—H14	0.9300
C2—H2	0.9300	C15—C16	1.476 (3)
C3—C4	1.365 (3)	C16—C17	1.488 (2)
C3—H3	0.9300	C17—C18	1.503 (3)
C4—C5	1.377 (3)	C18—C19	1.516 (3)
C4—H4	0.9300	C18—H18A	0.9700
C5—C6	1.379 (3)	C18—H18B	0.9700
C5—H5	0.9300	C19—C20	1.383 (3)
C6—C7	1.516 (2)	C19—C24	1.389 (2)
C7—C8	1.505 (2)	C20—C21	1.386 (3)
C7—H7A	0.9700	C20—H20	0.9300
C7—H7B	0.9700	C21—C22	1.371 (3)
C8—C17	1.350 (2)	C21—H21	0.9300
C8—C9	1.489 (3)	C22—C23	1.367 (3)
C9—C10	1.475 (2)	C22—H22	0.9300
C10—C11	1.387 (3)	C23—C24	1.385 (3)
C10—C15	1.398 (2)	C23—H23	0.9300
C11—C12	1.373 (3)		
C2—C1—C6	121.6 (2)	C14—C13—C12	120.6 (2)
C2—C1—Cl1	118.05 (16)	C14—C13—H13	119.7
C6—C1—Cl1	120.36 (15)	C12—C13—H13	119.7
C3—C2—C1	119.8 (2)	C13—C14—C15	120.1 (2)
C3—C2—H2	120.1	C13—C14—H14	120.0
C1—C2—H2	120.1	C15—C14—H14	120.0
C2—C3—C4	120.1 (2)	C14—C15—C10	119.49 (18)
C2—C3—H3	119.9	C14—C15—C16	120.95 (17)
C4—C3—H3	119.9	C10—C15—C16	119.53 (15)
C3—C4—C5	120.0 (2)	O2—C16—C15	120.86 (17)
C3—C4—H4	120.0	O2—C16—C17	119.48 (18)
C5—C4—H4	120.0	C15—C16—C17	119.62 (15)
C4—C5—C6	121.55 (19)	C8—C17—C16	120.59 (16)
C4—C5—H5	119.2	C8—C17—C18	124.53 (16)
C6—C5—H5	119.2	C16—C17—C18	114.86 (16)
C5—C6—C1	116.92 (17)	C17—C18—C19	114.23 (16)

C5—C6—C7	122.83 (16)	C17—C18—H18A	108.7
C1—C6—C7	120.25 (16)	C19—C18—H18A	108.7
C8—C7—C6	114.41 (15)	C17—C18—H18B	108.7
C8—C7—H7A	108.7	C19—C18—H18B	108.7
C6—C7—H7A	108.7	H18A—C18—H18B	107.6
C8—C7—H7B	108.7	C20—C19—C24	116.75 (18)
C6—C7—H7B	108.7	C20—C19—C18	122.56 (16)
H7A—C7—H7B	107.6	C24—C19—C18	120.68 (18)
C17—C8—C9	120.54 (15)	C19—C20—C21	121.61 (19)
C17—C8—C7	124.07 (17)	C19—C20—H20	119.2
C9—C8—C7	115.39 (15)	C21—C20—H20	119.2
O1—C9—C10	120.42 (17)	C22—C21—C20	120.0 (2)
O1—C9—C8	120.02 (17)	C22—C21—H21	120.0
C10—C9—C8	119.54 (15)	C20—C21—H21	120.0
C11—C10—C15	119.39 (16)	C23—C22—C21	120.0 (2)
C11—C10—C9	120.69 (16)	C23—C22—H22	120.0
C15—C10—C9	119.90 (15)	C21—C22—H22	120.0
C12—C11—C10	120.40 (19)	C22—C23—C24	119.5 (2)
C12—C11—H11	119.8	C22—C23—H23	120.2
C10—C11—H11	119.8	C24—C23—H23	120.2
C11—C12—C13	120.0 (2)	C19—C24—C23	122.1 (2)
C11—C12—H12	120.0	C19—C24—Cl2	119.53 (16)
C13—C12—H12	120.0	C23—C24—Cl2	118.37 (16)
C6—C1—C2—C3	1.4 (3)	C9—C10—C15—C14	-176.07 (17)
Cl1—C1—C2—C3	-178.88 (18)	C11—C10—C15—C16	-175.53 (16)
C1—C2—C3—C4	0.0 (4)	C9—C10—C15—C16	5.7 (2)
C2—C3—C4—C5	-0.9 (4)	C14—C15—C16—O2	-6.5 (3)
C3—C4—C5—C6	0.4 (4)	C10—C15—C16—O2	171.68 (18)
C4—C5—C6—C1	0.9 (3)	C14—C15—C16—C17	175.61 (17)
C4—C5—C6—C7	-178.7 (2)	C10—C15—C16—C17	-6.2 (3)
C2—C1—C6—C5	-1.8 (3)	C9—C8—C17—C16	-0.6 (2)
Cl1—C1—C6—C5	178.44 (14)	C7—C8—C17—C16	-179.65 (16)
C2—C1—C6—C7	177.85 (18)	C9—C8—C17—C18	-179.10 (16)
Cl1—C1—C6—C7	-1.9 (2)	C7—C8—C17—C18	1.8 (3)
C5—C6—C7—C8	14.0 (3)	O2—C16—C17—C8	-174.32 (18)
C1—C6—C7—C8	-165.61 (16)	C15—C16—C17—C8	3.6 (3)
C6—C7—C8—C17	-96.4 (2)	O2—C16—C17—C18	4.4 (3)
C6—C7—C8—C9	84.5 (2)	C15—C16—C17—C18	-177.76 (16)
C17—C8—C9—O1	-178.37 (17)	C8—C17—C18—C19	-104.7 (2)
C7—C8—C9—O1	0.8 (2)	C16—C17—C18—C19	76.7 (2)
C17—C8—C9—C10	0.1 (2)	C17—C18—C19—C20	9.4 (3)
C7—C8—C9—C10	179.22 (15)	C17—C18—C19—C24	-169.87 (16)
O1—C9—C10—C11	-3.0 (3)	C24—C19—C20—C21	0.2 (3)
C8—C9—C10—C11	178.54 (16)	C18—C19—C20—C21	-179.10 (18)
O1—C9—C10—C15	175.71 (17)	C19—C20—C21—C22	-0.2 (3)
C8—C9—C10—C15	-2.7 (2)	C20—C21—C22—C23	0.3 (3)
C15—C10—C11—C12	-2.2 (3)	C21—C22—C23—C24	-0.4 (3)
C9—C10—C11—C12	176.55 (18)	C20—C19—C24—C23	-0.3 (3)
C10—C11—C12—C13	0.1 (3)	C18—C19—C24—C23	179.01 (18)

## **supplementary materials**

---

C11—C12—C13—C14	1.6 (4)	C20—C19—C24—Cl2	179.97 (14)
C12—C13—C14—C15	-1.1 (3)	C18—C19—C24—Cl2	-0.7 (2)
C13—C14—C15—C10	-1.1 (3)	C22—C23—C24—C19	0.4 (3)
C13—C14—C15—C16	177.1 (2)	C22—C23—C24—Cl2	-179.87 (17)
C11—C10—C15—C14	2.7 (3)		

Fig. 1

