

N,4-Dimethyl-N-(4-nitrobenzyl)benzenesulfonamide

Hong-Yuan Zhu,* Zhou Wu and Shende Jiang

School of Pharmaceutical Science and Technology, Tianjin University, Tianjin 300072, People's Republic of China

Correspondence e-mail: zhhy93@163.com


Received 5 December 2007; accepted 5 January 2008

Key indicators: single-crystal X-ray study; $T = 113\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.057; wR factor = 0.124; data-to-parameter ratio = 13.2.

In the title compound, $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_4\text{S}$, there is a dihedral angle of $63.30(8)^\circ$ between the nitrobenzyl and benzene rings, which are separated by a sulfonamide unit. The crystal packing is stabilized by a $\text{C}-\text{H}\cdots\text{O}$ interaction.

Related literature

For the use of aromatic nitro and amine compounds as precursors in dye synthesis, see: Lauwiner *et al.* (1998); Yang *et al.* (2004). For the preparation of the title compound, see: Andersen *et al.* (1988). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_4\text{S}$	$V = 1510.0(5)\text{ \AA}^3$
$M_r = 320.36$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 9.5694(19)\text{ \AA}$	$\mu = 0.23\text{ mm}^{-1}$
$b = 6.1335(12)\text{ \AA}$	$T = 113(2)\text{ K}$
$c = 26.126(5)\text{ \AA}$	$0.20 \times 0.18 \times 0.10\text{ mm}$
$\beta = 100.03(3)^\circ$	

Data collection

Rigaku Saturn diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2003)
 $T_{\min} = 0.955$, $T_{\max} = 0.977$

8935 measured reflections
2661 independent reflections
2188 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.057$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.123$
 $S = 1.10$
2661 reflections

201 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.36\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C9—H9A…O1 ⁱ	0.99	2.32	3.287 (4)	166

Symmetry code: (i) $x, y - 1, z$.

Data collection: *CrystalClear* (Rigaku, 2003); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Bruker, 1997); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2454).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Andersen, K. K., Chumpradit, S. & McIntyre, D. J. (1988). *J. Org. Chem.* **53**, 4667–4675.
Bruker (1997). *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.
Lauwiner, M., Rys, P. & Wissmann, J. (1998). *Appl. Catal. A: Gen.* **172**, 141–148.
Rigaku (2003). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Yang, X. W. & Yang, W. (2004). *Shanghai Dyestuffs*, **32**, 20–30.

supplementary materials

Acta Cryst. (2008). E64, o596 [doi:10.1107/S1600536808000330]

N,4-Dimethyl-N-(4-nitrobenzyl)benzenesulfonamide

H.-Y. Zhu, Z. Wu and S. Jiang

Comment

Aromatic amines are widely employed for organic synthesis, especially as dye intermediates. One method for preparing aromatic amines is reduction of the corresponding aromatic nitro compound (Lauwiner *et al.*, 1998). In our recent work, the title compound (I), Fig. 1, was reduced to the corresponding aromatic amine for potential use as an intermediate in the synthesis of Acid Blue 264 dye according to Yang *et al.* (2004).

In (I), all bonds lengths and angles are normal (Allen *et al.*, 1987). The dihedral angle between the two aryl rings is 63.30 (8) Å. The distances of S1—C1 and S1—N1 are 1.762 (3) and 1.637 (2) Å respectively. The neighboring molecules are linked together by weak C—H···O hydrogen bonds, Table 1.

Experimental

The title compound (I) was synthesized according to the procedure of Andersen *et al.* (1988). Colorless single crystals (m.p. 403–404 K) were obtained by slow evaporation of a solution in absolute alcohol.

Refinement

C-bound H atoms were positioned geometrically and refined in the riding-model approximation, with $d(C—H) = 0.93$ Å, $U_{iso} = 1.2U_{eq}$ (C) for aromatic, 0.96 Å, $U_{iso} = 1.5U_{eq}$ (C) for CH_3 atoms and 0.97 Å, $U_{iso} = 1.2U_{eq}$ (C) for CH_2 atoms.

Figures




Fig. 1. The structure of (I) with displacement ellipsoids drawn at the 30% probability level and H atoms shown as small spheres of arbitrary radii.




Fig. 2. The crystal packing of (I) with hydrogen bonds drawn as dashed lines.

N,4-Dimethyl-N-(4-nitrobenzyl)benzenesulfonamide

Crystal data

$\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_4\text{S}$

$F_{000} = 672$

$M_r = 320.36$

$D_x = 1.409 \text{ Mg m}^{-3}$

supplementary materials

Monoclinic, $P2_1/c$	Melting point: 403-404 K
Hall symbol: -P 2ybc	Mo $K\alpha$ radiation
$a = 9.5694 (19) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 6.1335 (12) \text{ \AA}$	Cell parameters from 3609 reflections
$c = 26.126 (5) \text{ \AA}$	$\theta = 2.2\text{--}27.9^\circ$
$\beta = 100.03 (3)^\circ$	$\mu = 0.23 \text{ mm}^{-1}$
$V = 1510.0 (5) \text{ \AA}^3$	$T = 113 (2) \text{ K}$
$Z = 4$	Block, colorless
	$0.20 \times 0.18 \times 0.10 \text{ mm}$

Data collection

Rigaku Saturn diffractometer	2661 independent reflections
Radiation source: fine-focus sealed tube	2188 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.057$
Detector resolution: 7.31 pixels mm^{-1}	$\theta_{\text{max}} = 25.0^\circ$
$T = 113(2) \text{ K}$	$\theta_{\text{min}} = 2.2^\circ$
ω scans	$h = -11 \rightarrow 11$
Absorption correction: multi-scan (CrystalClear; Rigaku, 2003)	$k = -6 \rightarrow 7$
$T_{\text{min}} = 0.955, T_{\text{max}} = 0.977$	$l = -31 \rightarrow 26$
8935 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.057$	H-atom parameters constrained
$wR(F^2) = 0.123$	$w = 1/[\sigma^2(F_o^2) + (0.0454P)^2 + 1.1085P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.10$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2661 reflections	$\Delta\rho_{\text{max}} = 0.36 \text{ e \AA}^{-3}$
201 parameters	$\Delta\rho_{\text{min}} = -0.45 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculat-

ing R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.12405 (8)	0.46188 (11)	0.13266 (3)	0.0293 (2)
O1	0.0775 (3)	0.6688 (3)	0.14746 (9)	0.0531 (7)
O2	0.2167 (2)	0.4497 (4)	0.09498 (8)	0.0445 (6)
O3	-0.5173 (2)	-0.0352 (3)	-0.10469 (8)	0.0334 (5)
O4	-0.5383 (2)	-0.3199 (3)	-0.05751 (8)	0.0329 (5)
N1	-0.0187 (2)	0.3232 (3)	0.10887 (9)	0.0221 (5)
N2	-0.4801 (2)	-0.1489 (4)	-0.06589 (9)	0.0259 (5)
C1	0.2079 (3)	0.3328 (4)	0.19015 (11)	0.0210 (6)
C2	0.2046 (3)	0.4289 (5)	0.23762 (12)	0.0316 (7)
H2	0.1530	0.5601	0.2395	0.038*
C3	0.2765 (3)	0.3336 (5)	0.28224 (12)	0.0311 (7)
H3	0.2738	0.4004	0.3149	0.037*
C4	0.3529 (3)	0.1419 (4)	0.28053 (11)	0.0251 (6)
C5	0.3516 (3)	0.0458 (4)	0.23231 (12)	0.0285 (7)
H5	0.4007	-0.0878	0.2304	0.034*
C6	0.2809 (3)	0.1393 (4)	0.18703 (11)	0.0264 (6)
H6	0.2823	0.0723	0.1543	0.032*
C7	0.4382 (3)	0.0457 (5)	0.32894 (12)	0.0364 (8)
H7A	0.5367	0.0959	0.3327	0.055*
H7B	0.4356	-0.1137	0.3265	0.055*
H7C	0.3981	0.0922	0.3592	0.055*
C8	-0.1208 (3)	0.2999 (7)	0.14427 (14)	0.0496 (10)
H8A	-0.2139	0.2607	0.1244	0.074*
H8B	-0.1282	0.4381	0.1624	0.074*
H8C	-0.0886	0.1852	0.1697	0.074*
C9	0.0035 (3)	0.1211 (4)	0.08077 (12)	0.0282 (7)
H9A	0.0301	0.0020	0.1062	0.034*
H9B	0.0831	0.1428	0.0616	0.034*
C10	-0.1269 (3)	0.0558 (4)	0.04315 (11)	0.0222 (6)
C11	-0.1885 (3)	-0.1471 (4)	0.04786 (11)	0.0259 (6)
H11	-0.1499	-0.2403	0.0758	0.031*
C12	-0.3050 (3)	-0.2153 (4)	0.01256 (11)	0.0235 (6)
H12	-0.3473	-0.3534	0.0159	0.028*
C13	-0.3577 (3)	-0.0763 (4)	-0.02765 (11)	0.0215 (6)
C14	-0.2999 (3)	0.1272 (4)	-0.03363 (11)	0.0231 (6)
H14	-0.3386	0.2195	-0.0617	0.028*
C15	-0.1842 (3)	0.1921 (4)	0.00249 (11)	0.0223 (6)
H15	-0.1435	0.3317	-0.0006	0.027*

Atomic displacement parameters (\AA^2)

$$U^{11} \quad U^{22} \quad U^{33} \quad U^{12} \quad U^{13} \quad U^{23}$$

supplementary materials

S1	0.0358 (4)	0.0223 (4)	0.0238 (4)	-0.0080 (3)	-0.0117 (3)	0.0051 (3)
O1	0.0816 (18)	0.0182 (11)	0.0437 (16)	0.0046 (10)	-0.0330 (13)	-0.0008 (9)
O2	0.0393 (12)	0.0675 (16)	0.0242 (13)	-0.0274 (11)	-0.0015 (10)	0.0159 (11)
O3	0.0316 (11)	0.0426 (12)	0.0224 (12)	-0.0005 (9)	-0.0054 (9)	0.0013 (9)
O4	0.0290 (11)	0.0425 (12)	0.0269 (12)	-0.0163 (9)	0.0041 (9)	-0.0049 (9)
N1	0.0191 (11)	0.0236 (12)	0.0210 (14)	-0.0015 (9)	-0.0036 (10)	-0.0023 (9)
N2	0.0232 (12)	0.0347 (14)	0.0199 (14)	-0.0022 (10)	0.0039 (10)	-0.0082 (11)
C1	0.0172 (13)	0.0232 (14)	0.0192 (15)	-0.0034 (10)	-0.0060 (11)	0.0046 (11)
C2	0.0344 (16)	0.0269 (15)	0.0291 (18)	0.0094 (12)	-0.0067 (14)	-0.0033 (12)
C3	0.0319 (16)	0.0401 (17)	0.0188 (17)	0.0110 (13)	-0.0027 (13)	-0.0033 (13)
C4	0.0195 (13)	0.0292 (15)	0.0257 (17)	-0.0019 (11)	0.0018 (12)	0.0074 (12)
C5	0.0241 (14)	0.0258 (15)	0.0344 (19)	0.0062 (11)	0.0023 (13)	0.0022 (13)
C6	0.0255 (14)	0.0321 (15)	0.0212 (16)	0.0008 (12)	0.0034 (12)	-0.0032 (12)
C7	0.0321 (16)	0.0421 (18)	0.032 (2)	0.0049 (13)	-0.0044 (14)	0.0127 (14)
C8	0.0212 (16)	0.093 (3)	0.034 (2)	-0.0083 (17)	0.0030 (15)	-0.0152 (19)
C9	0.0248 (15)	0.0227 (14)	0.0340 (19)	0.0016 (11)	-0.0039 (13)	-0.0031 (12)
C10	0.0193 (13)	0.0219 (14)	0.0238 (16)	0.0008 (11)	-0.0006 (12)	-0.0048 (11)
C11	0.0250 (14)	0.0254 (14)	0.0252 (17)	0.0027 (11)	-0.0015 (12)	0.0026 (12)
C12	0.0225 (14)	0.0229 (14)	0.0255 (17)	-0.0025 (11)	0.0051 (12)	-0.0025 (11)
C13	0.0172 (13)	0.0271 (14)	0.0197 (16)	-0.0013 (10)	0.0016 (11)	-0.0069 (11)
C14	0.0219 (14)	0.0267 (14)	0.0202 (16)	0.0007 (11)	0.0025 (12)	0.0023 (11)
C15	0.0212 (14)	0.0194 (13)	0.0259 (17)	-0.0024 (10)	0.0033 (12)	-0.0015 (11)

Geometric parameters (\AA , $^\circ$)

S1—O1	1.421 (2)	C7—H7A	0.9800
S1—O2	1.437 (2)	C7—H7B	0.9800
S1—N1	1.637 (2)	C7—H7C	0.9800
S1—C1	1.762 (3)	C8—H8A	0.9800
O3—N2	1.231 (3)	C8—H8B	0.9800
O4—N2	1.225 (3)	C8—H8C	0.9800
N1—C8	1.464 (4)	C9—C10	1.502 (4)
N1—C9	1.475 (3)	C9—H9A	0.9900
N2—C13	1.470 (3)	C9—H9B	0.9900
C1—C2	1.378 (4)	C10—C15	1.387 (4)
C1—C6	1.387 (4)	C10—C11	1.392 (4)
C2—C3	1.376 (4)	C11—C12	1.382 (4)
C2—H2	0.9500	C11—H11	0.9500
C3—C4	1.390 (4)	C12—C13	1.379 (4)
C3—H3	0.9500	C12—H12	0.9500
C4—C5	1.389 (4)	C13—C14	1.385 (4)
C4—C7	1.502 (4)	C14—C15	1.382 (4)
C5—C6	1.381 (4)	C14—H14	0.9500
C5—H5	0.9500	C15—H15	0.9500
C6—H6	0.9500		
O1—S1—O2	119.53 (15)	C4—C7—H7C	109.5
O1—S1—N1	106.68 (13)	H7A—C7—H7C	109.5
O2—S1—N1	106.66 (13)	H7B—C7—H7C	109.5
O1—S1—C1	106.68 (14)	N1—C8—H8A	109.5

O2—S1—C1	108.47 (13)	N1—C8—H8B	109.5
N1—S1—C1	108.43 (12)	H8A—C8—H8B	109.5
C8—N1—C9	113.7 (2)	N1—C8—H8C	109.5
C8—N1—S1	114.5 (2)	H8A—C8—H8C	109.5
C9—N1—S1	116.25 (17)	H8B—C8—H8C	109.5
O4—N2—O3	123.8 (2)	N1—C9—C10	112.0 (2)
O4—N2—C13	118.1 (2)	N1—C9—H9A	109.2
O3—N2—C13	118.0 (2)	C10—C9—H9A	109.2
C2—C1—C6	120.6 (3)	N1—C9—H9B	109.2
C2—C1—S1	119.8 (2)	C10—C9—H9B	109.2
C6—C1—S1	119.6 (2)	H9A—C9—H9B	107.9
C3—C2—C1	119.6 (3)	C15—C10—C11	119.3 (2)
C3—C2—H2	120.2	C15—C10—C9	120.9 (2)
C1—C2—H2	120.2	C11—C10—C9	119.8 (2)
C2—C3—C4	121.3 (3)	C12—C11—C10	121.1 (3)
C2—C3—H3	119.3	C12—C11—H11	119.5
C4—C3—H3	119.3	C10—C11—H11	119.5
C5—C4—C3	117.8 (3)	C13—C12—C11	117.9 (2)
C5—C4—C7	121.0 (3)	C13—C12—H12	121.1
C3—C4—C7	121.2 (3)	C11—C12—H12	121.1
C6—C5—C4	121.7 (3)	C12—C13—C14	122.9 (2)
C6—C5—H5	119.1	C12—C13—N2	118.3 (2)
C4—C5—H5	119.1	C14—C13—N2	118.9 (2)
C5—C6—C1	118.8 (3)	C15—C14—C13	118.1 (3)
C5—C6—H6	120.6	C15—C14—H14	121.0
C1—C6—H6	120.6	C13—C14—H14	121.0
C4—C7—H7A	109.5	C14—C15—C10	120.8 (2)
C4—C7—H7B	109.5	C14—C15—H15	119.6
H7A—C7—H7B	109.5	C10—C15—H15	119.6
O1—S1—N1—C8	-56.7 (3)	C2—C1—C6—C5	-0.7 (4)
O2—S1—N1—C8	174.5 (2)	S1—C1—C6—C5	177.0 (2)
C1—S1—N1—C8	57.9 (2)	C8—N1—C9—C10	66.9 (3)
O1—S1—N1—C9	167.4 (2)	S1—N1—C9—C10	-156.8 (2)
O2—S1—N1—C9	38.6 (2)	N1—C9—C10—C15	60.7 (3)
C1—S1—N1—C9	-78.0 (2)	N1—C9—C10—C11	-121.6 (3)
O1—S1—C1—C2	6.3 (3)	C15—C10—C11—C12	0.5 (4)
O2—S1—C1—C2	136.2 (2)	C9—C10—C11—C12	-177.3 (3)
N1—S1—C1—C2	-108.3 (2)	C10—C11—C12—C13	0.3 (4)
O1—S1—C1—C6	-171.4 (2)	C11—C12—C13—C14	-0.6 (4)
O2—S1—C1—C6	-41.4 (2)	C11—C12—C13—N2	179.0 (2)
N1—S1—C1—C6	74.0 (2)	O4—N2—C13—C12	7.1 (4)
C6—C1—C2—C3	1.1 (4)	O3—N2—C13—C12	-172.3 (2)
S1—C1—C2—C3	-176.6 (2)	O4—N2—C13—C14	-173.3 (2)
C1—C2—C3—C4	0.1 (5)	O3—N2—C13—C14	7.3 (4)
C2—C3—C4—C5	-1.6 (4)	C12—C13—C14—C15	0.0 (4)
C2—C3—C4—C7	176.3 (3)	N2—C13—C14—C15	-179.5 (2)
C3—C4—C5—C6	2.0 (4)	C13—C14—C15—C10	0.8 (4)
C7—C4—C5—C6	-175.9 (3)	C11—C10—C15—C14	-1.0 (4)
C4—C5—C6—C1	-0.9 (4)	C9—C10—C15—C14	176.7 (2)


supplementary materials

Hydrogen-bond geometry (Å, °)

$D\text{---H}\cdots A$	$D\text{---H}$	$H\cdots A$	$D\cdots A$	$D\text{---H}\cdots A$
C9—H9A···O1 ⁱ	0.99	2.32	3.287 (4)	166

Symmetry codes: (i) $x, y-1, z$.

Fig. 1



supplementary materials

Fig. 2

