

(Heptanedioato-κ²O,O')bis(1,10-phenanthroline-κ²N,N')zinc(II) hexahydrate

Jian-Li Lin* and Yuan-Yuan Wang

State Key Laboratory, Base of Novel Functional Materials and Preparation Science, Institute of Solid Materials Chemistry, Faculty of Materials Science and Chemical Engineering, Ningbo University, Ningbo 315211, People's Republic of China
Correspondence e-mail: leikeweipublic@hotmail.com


Received 28 December 2007; accepted 4 March 2008

Key indicators: single-crystal X-ray study; T = 296 K; mean σ(C–C) = 0.005 Å; R factor = 0.060; wR factor = 0.117; data-to-parameter ratio = 16.3.

In the crystal structure of the title compound, [Zn(C₇H₁₀O₄)(C₁₂H₈N₂)₂].6H₂O, the Zn^{II} atom is coordinated by two carboxylate O atoms of a mono-bidentate chelating pimelate anion (pimelic acid is heptanedioic acid) and four N atoms of two phenanthroline ligands, forming a considerably distorted octahedral ZnN₄O₂ coordination geometry. The complexes are assembled into a three-dimensional network via C–H...O, C–H...π and π–π interactions. The mean interplanar distance between adjacent phenanthroline ligands is 3.399 (2) Å.

Related literature

For related literature, see: Ge & Zheng (2005); Wei *et al.* (2002); Zheng (2004); Zheng, Kong & Chen (2003); Zheng, Lin & Kong (2003); Zheng *et al.* (2001, 2002); Zheng & Ying (2004).



Experimental

Crystal data

[Zn(C₇H₁₀O₄)(C₁₂H₈N₂)₂].6H₂O
M_r = 692.02
Monoclinic, P2₁/n
a = 9.2050 (18) Å

b = 21.241 (4) Å
c = 16.598 (3) Å
β = 96.48 (3)°
V = 3224.6 (11) Å³
Z = 4

Mo Kα radiation
μ = 0.82 mm⁻¹
T = 296 (2) K
0.43 × 0.26 × 0.22 mm

Data collection

Bruker P4 diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
T_{min} = 0.697, T_{max} = 0.834
9226 measured reflections
7393 independent reflections

3956 reflections with I > 2σ(I)
R_{int} = 0.043
3 standard reflections
every 97 reflections
intensity decay: no

Refinement

R[F² > 2σ(F²)] = 0.059
wR(F²) = 0.117
S = 1.01
7393 reflections
453 parameters
18 restraints

H atoms treated by a mixture of independent and constrained refinement
Δρ_{max} = 0.34 e Å⁻³
Δρ_{min} = -0.29 e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Zn–N4	2.128 (3)	Zn–N1	2.167 (3)
Zn–N3	2.129 (3)	Zn–O1	2.178 (3)
Zn–N2	2.148 (3)	Zn–O2	2.224 (3)
N4–Zn–N3	78.26 (11)	N2–Zn–O1	87.11 (10)
N4–Zn–N2	99.28 (11)	N1–Zn–O1	146.42 (10)
N3–Zn–N2	168.21 (11)	N4–Zn–O2	156.42 (10)
N4–Zn–N1	109.01 (11)	N3–Zn–O2	90.89 (10)
N3–Zn–N1	92.26 (11)	N2–Zn–O2	95.35 (10)
N2–Zn–N1	77.55 (11)	N1–Zn–O2	92.10 (10)
N4–Zn–O1	102.80 (10)	O1–Zn–O2	59.45 (9)
N3–Zn–O1	104.68 (10)		

Table 2

Hydrogen-bond geometry (Å, °).

D–H...A	D–H	H...A	D...A	D–H...A
O5–H5A...O8 ⁱ	0.84 (2)	1.95 (2)	2.776 (5)	165 (5)
O5–H5B...O10	0.84 (4)	1.90 (4)	2.733 (6)	170 (4)
O6–H6A...O4 ⁱⁱ	0.85 (4)	2.02 (2)	2.861 (5)	176 (3)
O6–H6B...O7	0.83 (3)	2.16 (3)	2.985 (5)	170 (3)
O7–H7A...O5	0.85 (3)	1.89 (3)	2.734 (5)	176 (5)
O7–H7B...O2 ⁱⁱⁱ	0.86 (2)	1.97 (2)	2.828 (4)	174 (5)
O8–H8A...O9	0.85 (3)	1.96 (3)	2.804 (4)	172 (4)
O8–H8B...O4 ⁱⁱ	0.85 (3)	1.99 (3)	2.832 (4)	173 (3)
O9–H9A...O7	0.84 (4)	1.95 (2)	2.789 (4)	175 (4)
O9–H9B...O3 ^{iv}	0.85 (3)	1.89 (3)	2.736 (4)	173 (4)
O10–H10A...O3 ^v	0.86 (2)	1.88 (4)	2.732 (4)	171 (4)
O10–H10B...O1	0.85 (3)	2.11 (3)	2.957 (4)	176 (4)
C2–H2...O9 ^v	0.93	2.53	3.429 (5)	162
C5–H5...O2 ^{vi}	0.93	2.55	3.381 (4)	149
C17–H17...O1 ^{vii}	0.93	2.59	3.263 (4)	129
C18–H18...O6 ^{iv}	0.93	2.50	3.344 (5)	151
C26–H26B...Cg1 ⁱ	0.97	2.99	3.791 (4)	140
C27–H27A...Cg2 ⁱ	0.97	2.82	3.375 (4)	117

Symmetry codes: (i) x – 1, y, z; (ii) x + 1, y, z; (iii) -x + 1/2, y + 1/2, -z + 3/2; (iv) x + 1/2, -y + 1/2, z - 1/2; (v) -x + 3/2, y - 1/2, -z + 3/2; (vi) -x + 1, -y, -z + 2; (vii) -x + 1, -y, -z + 1. Cg1 and Cg2 are the centroids of the C16–C19/C23/C24 and C23/C19–C22/N4 rings, respectively.

Data collection: XSCANS (Siemens, 1996); cell refinement: XSCANS; data reduction: XSCANS; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine

structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This project was supported by the Zhejiang Provincial Fund for Analysis and Measurements (grant No. 04058), the Scientific Research Fund of Ningbo University (grant No. XK200457), the Expert Project of Key Basic Research of the Ministry of Science and Technology of China (grant No. 2003CCA00800), the Zhejiang Provincial Natural Science Foundation (grant No. Z203067) and the Ningbo Municipal Natural Science Foundation (grant No. 2003 A62026).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2271).

References

- Ge, C.-X. & Zheng, Y.-Q. (2005). *J. Coord. Chem.* **58**, 1199–1208.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). *XSCANS*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Wei, D.-Y., Zheng, Y.-Q. & Lin, J.-L. (2002). *Z. Anorg. Allg. Chem.* **628**, 2005–2012.
- Zheng, Y.-Q. (2004). *Z. Kristallogr. NCS*, **219**, 427–428.
- Zheng, Y.-Q., Kong, Z.-P. & Chen, K. (2003). *Z. Kristallogr. New Cryst. Struct.* **218**, 225–226.
- Zheng, Y.-Q., Lin, J.-L. & Kong, Z.-P. (2003). *Z. Anorg. Allg. Chem.* **629**, 357–361.
- Zheng, Y.-Q., Lin, J.-L. & Sun, J. (2001). *Z. Anorg. Allg. Chem.* **627**, 1997–2001.
- Zheng, Y.-Q., Liu, W.-H. & Lin, J.-L. (2002). *Z. Anorg. Allg. Chem.* **628**, 1401–1405.
- Zheng, Y.-Q. & Ying, E.-B. (2004). *Z. Kristallogr. New Cryst. Struct.* **219**, 423–424.

supplementary materials

Acta Cryst. (2008). E64, m527-m528 [doi:10.1107/S1600536808006004]

(Heptanedioato- κ^2O,O')bis(1,10-phenanthroline- κ^2N,N')zinc(II) hexahydrate

J.-L. Lin and Y.-Y. Wang

Comment

Previous investigation on self-assembly of metal ions, heteroaromatic N-donor ligands and pimelate anions exhibits various coordinating modes of pimelate anions. For example, pimelate anion bridges two metal ions in bis-monodentate fashion (Zheng, Lin *et al.*, 2003; Ge & Zheng, 2005; Zheng & Ying, 2004), in chelating/monodentate fashion (Zheng, 2004) and in bis-chelating fashion (Zheng, Kong *et al.*, 2003). When bridging three metal ions, pimelate anion can offer one carboxylate monoatomically to bridge two metal ions and the other end monodentately to coordinate one metal ion (Zheng *et al.*, 2001; Ge & Zheng, 2005). Furthermore, pimelate anion can bichelate two Cd atoms with one oxygen bonded to additional Cd atom to bridge three metal atoms to form polymeric chains (Zheng *et al.*, 2002). To the best of our knowledge, the title Zn compound represents a new example with pimelate anion coordinating one metal atom in a chelating fashion.

The title compound consists of $[\text{Zn}(\text{phen})_2(\text{C}_7\text{H}_{10}\text{O}_4)]$ complex and hydrogen bonded H_2O molecules. As demonstrated in Fig. 1, the Zn atom in the complex cation is coordinated by two carboxylate oxygen atoms of one mono-chelating pimelate $(\text{C}_7\text{H}_{10}\text{O}_4)^{2-}$ anion and four nitrogen atoms of two phenanthroline (phen) ligands to define a considerably distorted octahedral ZnN_4O_2 chromophore. Two phen ligands chelating the central Zn atom form V-shaped cleft and the mono-chelating pimelate ligand is twisted at the carbon atom next to the chelating carboxylate end to adopt a *gauche* conformation around the C26—C27 bond. The present complex looks very like the monovalent $[\text{Zn}(\text{phen})_2(\text{C}_8\text{H}_{13}\text{O}_4)]^+$ complex cation found in the earlier-reported $[\text{Zn}(\text{phen})_2(\text{C}_8\text{H}_{13}\text{O}_4)](\text{NO}_3)\cdot\text{H}_2\text{O}$, where the Zn atoms are coordinated by two phen ligands and one hydrogen suberate $(\text{C}_8\text{H}_{13}\text{O}_4)^-$ anion (Wei *et al.*, 2002).

Along [001] direction, the complex are arranged with the clefts orientating alternatively up- and downwards and the symmetry-related phen ligands orientate anti-parallelly to each other and the mean interplanar distance of 3.399 (2) Å suggests that the N-donor ligands are engaged in intercationic π - π stacking interactions. In this sense, the complex cations are, *via* π - π stacking interactions, assembled into one-dimensional chains extending parallel to [001] and careful inspection indicates that the resulting chains are stabilized by intercationic C5—H5 \cdots O2 hydrogen bonds. In the (011) plane, the chains are so arranged that the twisted pimelate ligands are located in the clefts of the adjacent chains and the alkyl C—H bonds are directed to the phen plane (C1 to C12) to form C—H \cdots π interactions. According to the above description, supramolecular assembly of the complex cations into two-dimensional layers (Fig. 2) is achieved due to intercationic π - π , C—H \cdots O and C—H \cdots π interactions. The lattice H_2O molecules are sandwiched between the cationic layers and form hydrogen bonded anionic chains propagating along [100]. The water molecules except the O5 one are hydrogen-bonded to the carboxylate O atoms.

Experimental

NaOH (2.0 ml, 1 M) was dropwise added to a stirred solution of $\text{Zn}(\text{NO}_3)_2\cdot 6\text{H}_2\text{O}$ (0.295 g, 0.99 mmol) in H_2O (5.0 ml) to produce white precipitate, which was separated by centrifugation and washed with deionized water for several times. The

supplementary materials

fresh precipitate was moved to a solution of phenanthroline monohydrate (0.200 g, 1.0 mmol) and pimelic acid (0.162 g, 1.0 mmol) in CH₃OH/H₂O (1:1 v/v; 20 ml), and then NaOH (0.5 ml, 1 M) was dropwise added. The resulting suspension was filtered out and the colorless filtrate (pH = 8.57) was allowed to stand at room temperature and slow evaporation for several weeks afforded a mixture of prismatic colorless crystals of [Zn(C₇H₁₀O₄)(C₁₂H₈N₂)₂] \cdot 6H₂O and plate-like colorless crystals. The former crystals are stable, and the latter are found to easily deteriorate upon isolation from the mother liquor.

Refinement

H atoms of water molecules were located in a difference Fourier map, and were refined with distance restraints of O—H = 0.85 (2) and H \cdots H = 1.38 (2) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$. Other H atoms were placed in geometrically idealized positions (C—H = 0.93–0.97 Å) and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the Zn^{II} complex, showing 30% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. A view of a single layer of (I). H atoms and water molecules have been omitted.

(Heptanedioato- $\kappa^2\text{O},\text{O}'$)bis(1,10-phenanthroline- $\kappa^2\text{N},\text{N}'$)zinc(II) hexahydrate

Crystal data

[Zn(C₇H₁₀O₄)(C₁₂H₈N₂)₂] \cdot 6H₂O

$M_r = 692.02$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.2050$ (18) Å

$b = 21.241$ (4) Å

$c = 16.598$ (3) Å

$\beta = 96.48$ (3)°

$V = 3224.6$ (11) Å³

$Z = 4$

$F_{000} = 1448$

$D_x = 1.425$ Mg m⁻³

Melting point: 163 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 5\text{--}12.5^\circ$

$\mu = 0.82$ mm⁻¹

$T = 296$ (2) K

Block, colourless

$0.43 \times 0.26 \times 0.22$ mm

Data collection

Bruker P4 diffractometer	$R_{\text{int}} = 0.043$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 27.5^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 1.6^\circ$
$T = 296(2)$ K	$h = -11 \rightarrow 1$
$\theta/2\theta$ scans	$k = -1 \rightarrow 27$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = -21 \rightarrow 21$
$T_{\text{min}} = 0.697$, $T_{\text{max}} = 0.834$	3 standard reflections
9226 measured reflections	every 97 reflections
7393 independent reflections	intensity decay: no
3956 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.059$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.117$	$w = 1/[\sigma^2(F_o^2) + (0.0346P)^2 + 1.0334P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
7393 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
453 parameters	$\Delta\rho_{\text{max}} = 0.34 \text{ e } \text{\AA}^{-3}$
18 restraints	$\Delta\rho_{\text{min}} = -0.29 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.00132 (18)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
-----	-----	-----	----------------------------------

supplementary materials

Zn	0.45796 (4)	0.04742 (2)	0.72867 (2)	0.03662 (14)
N1	0.6065 (3)	0.00835 (14)	0.82659 (17)	0.0385 (7)
N2	0.4468 (3)	0.11526 (14)	0.82373 (17)	0.0396 (7)
C1	0.6859 (4)	-0.04345 (19)	0.8282 (2)	0.0484 (9)
H1	0.6865	-0.0662	0.7804	0.058*
C2	0.7686 (5)	-0.0660 (2)	0.8973 (3)	0.0619 (12)
H2	0.8243	-0.1024	0.8952	0.074*
C3	0.7668 (5)	-0.0338 (2)	0.9682 (3)	0.0629 (12)
H3	0.8202	-0.0486	1.0153	0.075*
C4	0.6848 (4)	0.02125 (19)	0.9703 (2)	0.0486 (10)
C5	0.6771 (5)	0.0579 (2)	1.0425 (2)	0.0644 (13)
H5	0.7280	0.0448	1.0911	0.077*
C6	0.5972 (5)	0.1109 (2)	1.0404 (2)	0.0654 (13)
H6	0.5950	0.1339	1.0878	0.079*
C7	0.5155 (4)	0.13299 (19)	0.9674 (2)	0.0478 (10)
C8	0.4325 (5)	0.1880 (2)	0.9616 (3)	0.0614 (12)
H8	0.4282	0.2131	1.0072	0.074*
C9	0.3580 (5)	0.20512 (19)	0.8897 (3)	0.0575 (11)
H9	0.3010	0.2414	0.8859	0.069*
C10	0.3680 (4)	0.16756 (19)	0.8216 (2)	0.0513 (10)
H10	0.3171	0.1798	0.7725	0.062*
C11	0.5207 (4)	0.09748 (17)	0.8962 (2)	0.0405 (9)
C12	0.6062 (4)	0.04080 (17)	0.8974 (2)	0.0394 (9)
N3	0.5138 (3)	-0.02261 (14)	0.64572 (17)	0.0359 (7)
N4	0.5801 (3)	0.10061 (13)	0.65101 (16)	0.0339 (7)
C13	0.4767 (4)	-0.08320 (19)	0.6419 (2)	0.0479 (10)
H13	0.4083	-0.0977	0.6747	0.058*
C14	0.5360 (4)	-0.12543 (18)	0.5911 (2)	0.0474 (10)
H14	0.5051	-0.1671	0.5891	0.057*
C15	0.6391 (4)	-0.10603 (18)	0.5445 (2)	0.0455 (10)
H15	0.6815	-0.1345	0.5115	0.055*
C16	0.6810 (4)	-0.04252 (17)	0.54645 (18)	0.0357 (8)
C17	0.7875 (4)	-0.01760 (19)	0.4982 (2)	0.0458 (10)
H17	0.8350	-0.0444	0.4654	0.055*
C18	0.8192 (4)	0.0444 (2)	0.4999 (2)	0.0451 (9)
H18	0.8873	0.0599	0.4676	0.054*
C19	0.7503 (4)	0.08676 (17)	0.5505 (2)	0.0366 (8)
C20	0.7791 (4)	0.15174 (18)	0.5536 (2)	0.0471 (10)
H20	0.8472	0.1691	0.5227	0.057*
C21	0.7055 (5)	0.18921 (19)	0.6029 (2)	0.0542 (11)
H21	0.7200	0.2326	0.6036	0.065*
C22	0.6091 (4)	0.16208 (17)	0.6518 (2)	0.0444 (9)
H22	0.5632	0.1879	0.6865	0.053*
C23	0.6491 (3)	0.06355 (16)	0.60015 (19)	0.0320 (8)
C24	0.6144 (3)	-0.00242 (16)	0.59791 (18)	0.0293 (8)
O1	0.2329 (3)	0.07270 (12)	0.68624 (14)	0.0445 (6)
O2	0.2732 (3)	-0.01112 (12)	0.76159 (15)	0.0468 (7)
C25	0.1853 (4)	0.02656 (18)	0.7233 (2)	0.0386 (9)
C26	0.0227 (4)	0.01661 (18)	0.7220 (2)	0.0451 (10)

H26A	0.0056	-0.0170	0.7594	0.054*
H26B	-0.0172	0.0034	0.6680	0.054*
C27	-0.0573 (4)	0.07532 (18)	0.7449 (2)	0.0463 (10)
H27A	-0.1616	0.0672	0.7362	0.056*
H27B	-0.0365	0.1091	0.7086	0.056*
C28	-0.0188 (4)	0.09761 (17)	0.8318 (2)	0.0427 (9)
H28A	-0.0387	0.0641	0.8687	0.051*
H28B	0.0849	0.1070	0.8408	0.051*
C29	-0.1048 (4)	0.15551 (18)	0.8501 (2)	0.0474 (10)
H29A	-0.0795	0.1893	0.8149	0.057*
H29B	-0.2079	0.1466	0.8365	0.057*
C30	-0.0816 (4)	0.17858 (18)	0.9365 (2)	0.0497 (10)
H30A	0.0187	0.1929	0.9476	0.060*
H30B	-0.0934	0.1431	0.9720	0.060*
C31	-0.1816 (4)	0.23129 (17)	0.9587 (2)	0.0424 (9)
O3	-0.1667 (3)	0.24772 (12)	1.03240 (15)	0.0561 (7)
O4	-0.2700 (3)	0.25491 (12)	0.90506 (16)	0.0518 (7)
O5	0.0621 (4)	0.29022 (19)	0.7220 (3)	0.0979 (13)
H5A	-0.029 (2)	0.294 (2)	0.722 (3)	0.117*
H5B	0.082 (5)	0.2586 (18)	0.695 (3)	0.117*
O6	0.5105 (4)	0.35021 (17)	0.9056 (2)	0.0760 (10)
H6A	0.577 (4)	0.3225 (18)	0.908 (2)	0.091*
H6B	0.463 (4)	0.351 (2)	0.8597 (16)	0.091*
O7	0.3106 (3)	0.36089 (15)	0.7516 (2)	0.0664 (9)
H7A	0.234 (3)	0.3389 (15)	0.740 (3)	0.080*
H7B	0.286 (4)	0.3996 (9)	0.744 (3)	0.080*
O8	0.7737 (3)	0.30601 (17)	0.75204 (18)	0.0672 (9)
H8A	0.694 (3)	0.312 (2)	0.7220 (19)	0.081*
H8B	0.754 (4)	0.289 (2)	0.7958 (15)	0.081*
O9	0.5085 (3)	0.31376 (16)	0.65131 (18)	0.0663 (9)
H9A	0.453 (4)	0.3297 (19)	0.683 (2)	0.080*
H9B	0.456 (4)	0.2967 (19)	0.6116 (18)	0.080*
O10	0.1348 (4)	0.19689 (15)	0.6205 (2)	0.0749 (10)
H10A	0.196 (4)	0.2109 (18)	0.589 (2)	0.090*
H10B	0.166 (5)	0.1614 (13)	0.638 (3)	0.090*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn	0.0340 (2)	0.0411 (3)	0.0360 (2)	0.0035 (2)	0.00918 (16)	-0.0029 (2)
N1	0.0361 (17)	0.0382 (18)	0.0412 (18)	-0.0007 (15)	0.0050 (14)	-0.0012 (15)
N2	0.0360 (17)	0.0397 (18)	0.0446 (19)	-0.0014 (15)	0.0117 (15)	-0.0034 (15)
C1	0.050 (2)	0.044 (2)	0.050 (2)	0.005 (2)	0.0023 (19)	0.003 (2)
C2	0.055 (3)	0.059 (3)	0.070 (3)	0.017 (2)	-0.001 (2)	0.016 (2)
C3	0.062 (3)	0.069 (3)	0.055 (3)	-0.002 (3)	-0.008 (2)	0.016 (2)
C4	0.049 (2)	0.052 (3)	0.043 (2)	-0.011 (2)	-0.0036 (19)	0.006 (2)
C5	0.071 (3)	0.080 (4)	0.041 (2)	-0.012 (3)	-0.001 (2)	0.003 (2)
C6	0.081 (3)	0.076 (3)	0.040 (3)	-0.023 (3)	0.010 (2)	-0.019 (2)

supplementary materials

C7	0.053 (2)	0.048 (3)	0.044 (2)	-0.014 (2)	0.012 (2)	-0.0103 (19)
C8	0.072 (3)	0.050 (3)	0.066 (3)	-0.006 (2)	0.020 (3)	-0.029 (2)
C9	0.056 (3)	0.041 (2)	0.078 (3)	0.006 (2)	0.018 (2)	-0.015 (2)
C10	0.047 (2)	0.048 (3)	0.060 (3)	0.002 (2)	0.010 (2)	-0.006 (2)
C11	0.038 (2)	0.043 (2)	0.041 (2)	-0.0112 (19)	0.0085 (18)	-0.0016 (18)
C12	0.041 (2)	0.040 (2)	0.038 (2)	-0.0066 (19)	0.0083 (17)	0.0002 (18)
N3	0.0355 (17)	0.0338 (17)	0.0385 (17)	0.0038 (14)	0.0044 (14)	-0.0020 (13)
N4	0.0336 (16)	0.0360 (18)	0.0329 (16)	0.0057 (14)	0.0078 (13)	-0.0007 (14)
C13	0.043 (2)	0.049 (3)	0.053 (3)	-0.002 (2)	0.010 (2)	0.004 (2)
C14	0.050 (2)	0.037 (2)	0.054 (2)	0.003 (2)	0.000 (2)	-0.005 (2)
C15	0.051 (2)	0.045 (2)	0.039 (2)	0.010 (2)	0.0009 (19)	-0.0086 (18)
C16	0.0365 (19)	0.044 (2)	0.0256 (17)	0.0073 (19)	-0.0020 (15)	-0.0029 (17)
C17	0.054 (2)	0.053 (3)	0.033 (2)	0.011 (2)	0.0116 (18)	-0.0075 (18)
C18	0.047 (2)	0.058 (3)	0.0324 (19)	0.006 (2)	0.0126 (17)	0.003 (2)
C19	0.035 (2)	0.045 (2)	0.0300 (19)	0.0011 (18)	0.0048 (16)	0.0051 (17)
C20	0.051 (2)	0.047 (2)	0.045 (2)	0.000 (2)	0.0145 (19)	0.005 (2)
C21	0.066 (3)	0.036 (2)	0.063 (3)	-0.006 (2)	0.020 (2)	0.002 (2)
C22	0.050 (2)	0.037 (2)	0.047 (2)	0.0023 (19)	0.0128 (19)	-0.0064 (18)
C23	0.0281 (18)	0.040 (2)	0.0274 (18)	0.0062 (16)	0.0009 (15)	0.0012 (15)
C24	0.0280 (18)	0.036 (2)	0.0229 (17)	0.0040 (16)	0.0006 (14)	-0.0008 (15)
O1	0.0456 (15)	0.0491 (16)	0.0402 (14)	-0.0004 (13)	0.0104 (12)	-0.0003 (13)
O2	0.0367 (14)	0.0506 (17)	0.0530 (16)	0.0073 (13)	0.0044 (12)	-0.0009 (13)
C25	0.038 (2)	0.047 (2)	0.033 (2)	-0.0004 (19)	0.0100 (17)	-0.0141 (18)
C26	0.033 (2)	0.053 (2)	0.050 (2)	-0.0004 (19)	0.0042 (17)	-0.013 (2)
C27	0.037 (2)	0.055 (2)	0.046 (2)	0.0071 (19)	0.0028 (18)	-0.0063 (19)
C28	0.034 (2)	0.051 (2)	0.043 (2)	0.0076 (19)	0.0055 (17)	-0.0003 (19)
C29	0.045 (2)	0.045 (2)	0.053 (2)	0.0041 (19)	0.0102 (19)	-0.003 (2)
C30	0.049 (2)	0.051 (3)	0.049 (2)	0.009 (2)	0.0070 (19)	-0.004 (2)
C31	0.046 (2)	0.032 (2)	0.052 (3)	0.0022 (19)	0.018 (2)	-0.0001 (19)
O3	0.075 (2)	0.0500 (17)	0.0447 (17)	0.0128 (16)	0.0137 (15)	-0.0042 (14)
O4	0.0510 (17)	0.0512 (17)	0.0543 (17)	0.0130 (14)	0.0110 (14)	0.0012 (14)
O5	0.076 (2)	0.083 (3)	0.145 (4)	-0.020 (2)	0.054 (3)	-0.036 (2)
O6	0.073 (2)	0.078 (2)	0.077 (2)	0.0225 (19)	0.0106 (18)	-0.023 (2)
O7	0.0578 (19)	0.060 (2)	0.083 (2)	0.0079 (16)	0.0120 (18)	-0.0120 (19)
O8	0.055 (2)	0.095 (3)	0.0522 (18)	0.0039 (19)	0.0068 (15)	0.0066 (18)
O9	0.0517 (19)	0.083 (2)	0.064 (2)	0.0020 (17)	0.0033 (15)	-0.0208 (17)
O10	0.087 (2)	0.059 (2)	0.085 (2)	0.0016 (19)	0.042 (2)	0.0066 (18)

Geometric parameters (Å, °)

Zn—N4	2.128 (3)	C17—H17	0.9300
Zn—N3	2.129 (3)	C18—C19	1.427 (5)
Zn—N2	2.148 (3)	C18—H18	0.9300
Zn—N1	2.167 (3)	C19—C23	1.401 (5)
Zn—O1	2.178 (3)	C19—C20	1.405 (5)
Zn—O2	2.224 (3)	C20—C21	1.374 (5)
Zn—C25	2.540 (4)	C20—H20	0.9300
N1—C1	1.320 (4)	C21—C22	1.393 (5)
N1—C12	1.363 (4)	C21—H21	0.9300

N2—C10	1.325 (5)	C22—H22	0.9300
N2—C11	1.367 (4)	C23—C24	1.437 (5)
C1—C2	1.388 (5)	O1—C25	1.262 (4)
C1—H1	0.9300	O2—C25	1.259 (4)
C2—C3	1.363 (6)	C25—C26	1.509 (5)
C2—H2	0.9300	C26—C27	1.519 (5)
C3—C4	1.395 (6)	C26—H26A	0.9700
C3—H3	0.9300	C26—H26B	0.9700
C4—C12	1.400 (5)	C27—C28	1.521 (5)
C4—C5	1.438 (6)	C27—H27A	0.9700
C5—C6	1.341 (6)	C27—H27B	0.9700
C5—H5	0.9300	C28—C29	1.512 (5)
C6—C7	1.431 (6)	C28—H28A	0.9700
C6—H6	0.9300	C28—H28B	0.9700
C7—C8	1.393 (6)	C29—C30	1.507 (5)
C7—C11	1.408 (5)	C29—H29A	0.9700
C8—C9	1.357 (6)	C29—H29B	0.9700
C8—H8	0.9300	C30—C31	1.521 (5)
C9—C10	1.395 (5)	C30—H30A	0.9700
C9—H9	0.9300	C30—H30B	0.9700
C10—H10	0.9300	C31—O4	1.242 (4)
C11—C12	1.437 (5)	C31—O3	1.264 (4)
N3—C13	1.331 (4)	O5—H5A	0.84 (2)
N3—C24	1.355 (4)	O5—H5B	0.84 (4)
N4—C22	1.333 (4)	O6—H6A	0.85 (4)
N4—C23	1.363 (4)	O6—H6B	0.83 (3)
C13—C14	1.385 (5)	O7—H7A	0.85 (3)
C13—H13	0.9300	O7—H7B	0.86 (2)
C14—C15	1.355 (5)	O8—H8A	0.85 (3)
C14—H14	0.9300	O8—H8B	0.85 (3)
C15—C16	1.403 (5)	O9—H9A	0.84 (4)
C15—H15	0.9300	O9—H9B	0.85 (3)
C16—C24	1.396 (4)	O10—H10A	0.86 (4)
C16—C17	1.436 (5)	O10—H10B	0.85 (3)
C17—C18	1.349 (5)		
N4—Zn—N3	78.26 (11)	C14—C15—H15	120.4
N4—Zn—N2	99.28 (11)	C16—C15—H15	120.4
N3—Zn—N2	168.21 (11)	C24—C16—C15	117.5 (3)
N4—Zn—N1	109.01 (11)	C24—C16—C17	119.4 (3)
N3—Zn—N1	92.26 (11)	C15—C16—C17	123.0 (3)
N2—Zn—N1	77.55 (11)	C18—C17—C16	120.5 (3)
N4—Zn—O1	102.80 (10)	C18—C17—H17	119.8
N3—Zn—O1	104.68 (10)	C16—C17—H17	119.8
N2—Zn—O1	87.11 (10)	C17—C18—C19	121.3 (3)
N1—Zn—O1	146.42 (10)	C17—C18—H18	119.4
N4—Zn—O2	156.42 (10)	C19—C18—H18	119.4
N3—Zn—O2	90.89 (10)	C23—C19—C20	117.3 (3)
N2—Zn—O2	95.35 (10)	C23—C19—C18	119.6 (3)
N1—Zn—O2	92.10 (10)	C20—C19—C18	123.1 (3)

supplementary materials

O1—Zn—O2	59.45 (9)	C21—C20—C19	119.2 (4)
N4—Zn—C25	131.21 (12)	C21—C20—H20	120.4
N3—Zn—C25	99.51 (11)	C19—C20—H20	120.4
N2—Zn—C25	90.80 (11)	C20—C21—C22	119.8 (4)
N1—Zn—C25	119.78 (12)	C20—C21—H21	120.1
O1—Zn—C25	29.77 (10)	C22—C21—H21	120.1
O2—Zn—C25	29.69 (10)	N4—C22—C21	122.6 (3)
C1—N1—C12	117.3 (3)	N4—C22—H22	118.7
C1—N1—Zn	129.5 (3)	C21—C22—H22	118.7
C12—N1—Zn	113.0 (2)	N4—C23—C19	123.3 (3)
C10—N2—C11	118.2 (3)	N4—C23—C24	117.5 (3)
C10—N2—Zn	127.9 (3)	C19—C23—C24	119.2 (3)
C11—N2—Zn	113.6 (2)	N3—C24—C16	122.8 (3)
N1—C1—C2	123.8 (4)	N3—C24—C23	117.3 (3)
N1—C1—H1	118.1	C16—C24—C23	119.9 (3)
C2—C1—H1	118.1	C25—O1—Zn	91.2 (2)
C3—C2—C1	118.8 (4)	C25—O2—Zn	89.2 (2)
C3—C2—H2	120.6	O2—C25—O1	120.1 (3)
C1—C2—H2	120.6	O2—C25—C26	119.9 (3)
C2—C3—C4	120.1 (4)	O1—C25—C26	120.0 (4)
C2—C3—H3	120.0	O2—C25—Zn	61.09 (19)
C4—C3—H3	120.0	O1—C25—Zn	59.01 (18)
C3—C4—C12	117.1 (4)	C26—C25—Zn	177.7 (3)
C3—C4—C5	123.5 (4)	C25—C26—C27	112.9 (3)
C12—C4—C5	119.3 (4)	C25—C26—H26A	109.0
C6—C5—C4	120.7 (4)	C27—C26—H26A	109.0
C6—C5—H5	119.6	C25—C26—H26B	109.0
C4—C5—H5	119.6	C27—C26—H26B	109.0
C5—C6—C7	122.1 (4)	H26A—C26—H26B	107.8
C5—C6—H6	118.9	C26—C27—C28	115.2 (3)
C7—C6—H6	118.9	C26—C27—H27A	108.5
C8—C7—C11	117.5 (4)	C28—C27—H27A	108.5
C8—C7—C6	124.5 (4)	C26—C27—H27B	108.5
C11—C7—C6	118.0 (4)	C28—C27—H27B	108.5
C9—C8—C7	120.3 (4)	H27A—C27—H27B	107.5
C9—C8—H8	119.8	C29—C28—C27	111.8 (3)
C7—C8—H8	119.8	C29—C28—H28A	109.2
C8—C9—C10	119.1 (4)	C27—C28—H28A	109.2
C8—C9—H9	120.5	C29—C28—H28B	109.2
C10—C9—H9	120.5	C27—C28—H28B	109.2
N2—C10—C9	123.0 (4)	H28A—C28—H28B	107.9
N2—C10—H10	118.5	C30—C29—C28	115.8 (3)
C9—C10—H10	118.5	C30—C29—H29A	108.3
N2—C11—C7	122.0 (4)	C28—C29—H29A	108.3
N2—C11—C12	117.5 (3)	C30—C29—H29B	108.3
C7—C11—C12	120.5 (3)	C28—C29—H29B	108.3
N1—C12—C4	122.9 (4)	H29A—C29—H29B	107.4
N1—C12—C11	117.7 (3)	C29—C30—C31	116.5 (3)
C4—C12—C11	119.3 (3)	C29—C30—H30A	108.2

C13—N3—C24	117.8 (3)	C31—C30—H30A	108.2
C13—N3—Zn	128.8 (3)	C29—C30—H30B	108.2
C24—N3—Zn	112.9 (2)	C31—C30—H30B	108.2
C22—N4—C23	117.7 (3)	H30A—C30—H30B	107.3
C22—N4—Zn	129.3 (2)	O4—C31—O3	124.9 (3)
C23—N4—Zn	112.6 (2)	O4—C31—C30	119.2 (3)
N3—C13—C14	122.6 (4)	O3—C31—C30	115.8 (4)
N3—C13—H13	118.7	H5A—O5—H5B	111 (3)
C14—C13—H13	118.7	H6A—O6—H6B	111 (3)
C15—C14—C13	120.0 (4)	H7A—O7—H7B	107 (3)
C15—C14—H14	120.0	H8A—O8—H8B	108 (3)
C13—C14—H14	120.0	H9A—O9—H9B	108 (3)
C14—C15—C16	119.2 (3)	H10A—O10—H10B	107 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O5—H5A \cdots O8 ⁱ	0.84 (2)	1.95 (2)	2.776 (5)	165 (5)
O5—H5B \cdots O10	0.84 (4)	1.90 (4)	2.733 (6)	170 (4)
O6—H6A \cdots O4 ⁱⁱ	0.85 (4)	2.02 (2)	2.861 (5)	176 (3)
O6—H6B \cdots O7	0.83 (3)	2.16 (3)	2.985 (5)	170 (3)
O7—H7A \cdots O5	0.85 (3)	1.89 (3)	2.734 (5)	176 (5)
O7—H7B \cdots O2 ⁱⁱⁱ	0.86 (2)	1.97 (2)	2.828 (4)	174 (5)
O8—H8A \cdots O9	0.85 (3)	1.96 (3)	2.804 (4)	172 (4)
O8—H8B \cdots O4 ⁱⁱ	0.85 (3)	1.99 (3)	2.832 (4)	173 (3)
O9—H9A \cdots O7	0.84 (4)	1.95 (2)	2.789 (4)	175 (4)
O9—H9B \cdots O3 ^{iv}	0.85 (3)	1.89 (3)	2.736 (4)	173 (4)
O10—H10A \cdots O3 ^{iv}	0.86 (2)	1.88 (4)	2.732 (4)	171 (4)
O10—H10B \cdots O1	0.85 (3)	2.11 (3)	2.957 (4)	176 (4)
C2—H2 \cdots O9 ^v	0.93	2.53	3.429 (5)	162
C5—H5 \cdots O2 ^{vi}	0.93	2.55	3.381 (4)	149
C17—H17 \cdots O1 ^{vii}	0.93	2.59	3.263 (4)	129
C18—H18 \cdots O6 ^{iv}	0.93	2.50	3.344 (5)	151
C26—H26B \cdots Cg1 ⁱ	0.97	2.99	3.791 (4)	140
C27—H27A \cdots Cg2 ⁱ	0.97	2.82	3.375 (4)	117

Symmetry codes: (i) $x-1, y, z$; (ii) $x+1, y, z$; (iii) $-x+1/2, y+1/2, -z+3/2$; (iv) $x+1/2, -y+1/2, z-1/2$; (v) $-x+3/2, y-1/2, -z+3/2$; (vi) $-x+1, -y, -z+2$; (vii) $-x+1, -y, -z+1$.

Fig. 1




Fig. 2

