

**(2,2'-Bipyridine- κ^2N,N') $\{[(3\text{-methoxy-}2\text{-oxidobenzylidene-}\kappa O^2\text{)hydrazono]-}$
methanolato- $\kappa^2N^2,O\}$ dimethyltin(IV)**

Shaukat Shuja,^a Saqib Ali,^a M. Nawaz Tahir,^b Nasir Khalid^c and Islam Ullah Khan^{d*}

^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan,

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan, ^cChemistry

Division, Pakistan Institute of Nuclear Science and Technology, PO Nilore,

Islamabad, Pakistan, and ^dDepartment of Chemistry, Government College University, Lahore, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 28 February 2008; accepted 5 March 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.004$ Å;
 R factor = 0.025; wR factor = 0.067; data-to-parameter ratio = 19.7.

In the crystal structure of the title compound, $[Sn(CH_3)_2(C_9H_8N_2O_3)(C_{10}H_8N_2)]$, the Sn atom exhibits a pentagonal bipyramidal coordination geometry defined by two C, three N and two O atoms. The bond distances for Sn—C, Sn—N and Sn—O are in the ranges 2.097 (3)–2.098 (3), 2.298 (2)–2.623 (2) and 2.157 (2)–2.266 (2) Å, respectively. The molecular structure of the monomeric compound is stabilized by three intramolecular C—H···O hydrogen bonds, all involving bipyridine C—H groups.

Related literature

For related literature, see: Chen *et al.* (2006); Diouf *et al.* (2004); Shuja *et al.* (2007a,b, 2007c). For bond-length data, see: Allen (2002).



Experimental

Crystal data

$[Sn(CH_3)_2(C_9H_8N_2O_3)(C_{10}H_8N_2)]$

$M_r = 497.12$

Monoclinic, $P2_1/c$

$a = 12.3834$ (3) Å

$b = 9.9094$ (2) Å

$c = 17.1730$ (4) Å

$\beta = 103.302$ (1)°

$V = 2050.80$ (8) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 1.28$ mm⁻¹

$T = 296$ (2) K

$0.25 \times 0.18 \times 0.15$ mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.749$, $T_{\max} = 0.820$

24401 measured reflections

5524 independent reflections

4407 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.066$

$S = 1.04$

5524 reflections

280 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 1.12$ e Å⁻³

$\Delta\rho_{\min} = -0.41$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Sn1—C9	2.097 (3)	Sn1—N1	2.2980 (18)
Sn1—C10	2.098 (3)	Sn1—N3	2.5825 (18)
Sn1—O1	2.1572 (14)	Sn1—N4	2.6231 (19)
Sn1—O2	2.2658 (15)		
C9—Sn1—C10	169.76 (11)	C10—Sn1—N3	90.33 (9)
C9—Sn1—O1	93.97 (9)	O1—Sn1—N3	138.83 (6)
C10—Sn1—O1	90.20 (9)	O2—Sn1—N3	75.90 (6)
C9—Sn1—O2	94.23 (9)	N1—Sn1—N3	143.65 (6)
C10—Sn1—O2	87.60 (11)	O1—Sn1—N4	76.91 (6)
O1—Sn1—O2	145.23 (6)	O2—Sn1—N4	136.43 (6)
C9—Sn1—N1	94.39 (9)	N1—Sn1—N4	153.65 (6)
C10—Sn1—N1	95.65 (10)	N3—Sn1—N4	62.68 (6)
O1—Sn1—N1	77.12 (6)	N4—Sn1—C9	91.77 (9)
O2—Sn1—N1	68.60 (6)	N4—Sn1—C10	80.06 (10)
C9—Sn1—N3	80.37 (9)		

Table 2

Hydrogen-bond geometry (Å, °).

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C12—H12···O2	0.93	2.40	2.993 (3)	121
C21—H21···O1	0.93	2.36	2.968 (3)	123
C21—H21···O3	0.93	2.50	3.390 (3)	161

Data collection: *APEX2* (Bruker, 2007); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the Higher Education Commission, Islamabad, Pakistan, for funding the purchase of the diffractometer and for financial support to SS for PhD under the Indigenous Scholarship Scheme (PIN Code: 042–111889).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2056).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Bruker (2005). SADABS. Bruker AXS Inc. Madison, Wisconsin, USA.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc. Madison, Wisconsin, USA.
- Chen, S.-W., Yin, H.-D. & Wang, D.-Q. (2006). *Acta Cryst.* **E62**, m1654–m1655.
- Diouf, O., Gaye, M., Sall, A. S. & Slebodnick, C. (2004). *Z. Kristallogr.* **219**, 435–436.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shuja, S., Ali, S., Khalid, N., Broker, G. A. & Tiekink, E. R. T. (2007a). *Acta Cryst.* **E63**, m1025–m1026.
- Shuja, S., Ali, S., Khalid, N. & Parvez, M. (2007c). *Acta Cryst.* **E63**, o879–o880.
- Shuja, S., Ali, S., Meetsma, A., Broker, G. A. & Tiekink, E. R. T. (2007b). *Acta Cryst.* **E63**, m1130–m1132.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, m531-m532 [doi:10.1107/S1600536808006090]

(2,2'-Bipyridine- κ^2N,N') $\{[(3\text{-methoxy-}2\text{-oxidobenzylidene-}\kappa O^2)\text{hydrazone]methanolato-}$ $\kappa^2N^2,O\}$ dimethyltin(IV)

S. Shuja, S. Ali, M. N. Tahir, N. Khalid and I. U. Khan

Comment

Diorganotin(IV) complexes of Schiff base ligands derived from 3-methoxysalicylaldehyde and hydrazine derivatives are limited in number (Chen *et al.*, 2006, Diouf *et al.*, 2004). In continuation of our efforts to synthesize various Schiff base ligands of substituted salicylaldehydes with hydrazines or amino acids as well as the corresponding organotin derivatives (Shuja *et al.*, 2007a, 2007b, 2007c), we herein report the structure of the title compound (I).

In the monomeric structure of the title compound (I), the coordination around Sn consists of two O-atoms and one N-atom of the Schiff base ligand [*N*-Formyl-*N'*-(3-methoxy-2-oxidobenzylidene)hydrazine], two N-atoms of 2,2'-bipyridine and two C-atoms of methyl groups. The shortest bond of Sn is realised with the methyl C-atoms showing nearly equal values (2.097 (3) and 2.098 (3) Å) corresponding very well with the bond lengths observed in [(3-Methoxy-2-oxidobenzaldehyde benzoylhydrazone)dimethyltin(IV)] (Chen *et al.*, 2006). The Sn1—O1 bond distance of 2.157 (1) Å is greater than the values reported for [diphenyl(methoxy-*N*-salicylideneacetylhydrazone)tin(IV)] (2.068 (2) Å, Diouf *et al.*, 2004 and 2.131 (3) Å, Chen *et al.*, 2006). The same is true for the bonds Sn1—O2 (2.266 (2) Å) and Sn1—N1 (2.298 (2) Å) also being longer compared to those previously reported (Diouf *et al.*, 2004, Chen *et al.*, 2006). These observations are most probably due to the additional coordination of bipyridine to tin. A CCDC search (Allen, 2002) showed that the title compound is indeed the first structurally characterized tin organyl with a bipyridine ligand attached to tin. The bond distances of N-atoms of bipyridine measure to 2.583 (2) Å (Sn1—N3) and 2.623 (2) Å (Sn1—N4), respectively. The bond distances in the hydrazine ligand are comparable with those reported for {[*N*-Formyl-*N'*-(2-oxidobenzylidene)hydrazine- κ^3O,N,O']diphenyl tin(IV)} (Shuja *et al.*, 2007b). The bond angles around Sn1 are in the range between 68.60 (6)° and 169.8 (1)°. The dihedral angle between (O1/C1/C2/C3/C4 /C5/C6/C7) and (N1/N2/C8/O2) is 36.0 (1)° while the angle between the rings (N3/C12/C13/C14/C15/C16) and (N4/C17/C18/C19/C20/C21) is 17.0 (1)°. The molecular structure of the title compound as well as the observed conformation are stabilized by three intramolecular H-bonds of C—H···O type (Fig. 1) all involving bipyridine C—H functions (Table 2). The closest intermolecular contact of molecules is at a distance of 3.208 (3) Å between O1···C15ⁱ [symmetry code: $i = x, -y + 1/2, z + 1/2$]. A positive electron peak corresponding to 1.12 \AA^{-3} remains at a distance of 0.93 Å near Sn1.

Experimental

N-(2-hydroxy-3-methoxy-benzylidene)formylhydrazine (0.58 g, 3 mmol) and Et₃N (0.86 ml, 6 mmol) were added to anhydrous toluene (100 ml) in a round bottom flask equipped with a reflux condenser. Dimethyltin(IV) dichloride (0.66 g, 3 mmol) dissolved in anhydrous toluene (20 ml) and 2,2'-bipyridine (0.47 g, 3 mmol) were then added. The reaction mixture was stirred at room temperature for 5 hr and allowed to stand overnight. The Et₃NHCl salt formed during the reaction was filtered off and the resulting clear yellow solution was evaporated with a rotary evaporator under reduced pressure. Recrystallization from chloroform yielded crystals suitable for X-ray diffraction.

supplementary materials

Refinement

The coordinates of H-atoms of methyl carbons attached to tin were refined freely. The remaining H atoms were positioned geometrically, with C—H = 0.93 and 0.97 Å for aromatic and methyl H, and constrained to ride on their parent atoms. The $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H, and $x=1.2$ for aromatic H atoms.

Figures




Fig. 1. *ORTEP-3 for Windows* (Farrugia, 1997) drawing of the title compound, $\text{C}_{21}\text{H}_{22}\text{N}_4\text{O}_3\text{Sn}$ with the atom numbering scheme. The thermal ellipsoids are drawn at the 30% probability level. H-atoms are shown by small circles of arbitrary radii. The intramolecular H-bonding is shown by dashed lines.




Fig. 2. The unit cell packing of (I) (Spek, 2003), showing that there is no intermolecular hydrogen bonding.

(I)

Crystal data

$[\text{Sn}(\text{CH}_3)_2(\text{C}_9\text{H}_8\text{N}_2\text{O}_3)(\text{C}_{10}\text{H}_8\text{N}_2)]$	$F_{000} = 1000$
$M_r = 497.12$	$D_x = 1.610 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 12.3834 (3) \text{ \AA}$	Cell parameters from 4407 reflections
$b = 9.9094 (2) \text{ \AA}$	$\theta = 1.7\text{--}29.2^\circ$
$c = 17.1730 (4) \text{ \AA}$	$\mu = 1.28 \text{ mm}^{-1}$
$\beta = 103.302 (1)^\circ$	$T = 296 (2) \text{ K}$
$V = 2050.80 (8) \text{ \AA}^3$	Prismatic, yellow
$Z = 4$	$0.25 \times 0.18 \times 0.15 \text{ mm}$

Data collection

Bruker KappaAPEXII CCD diffractometer	5524 independent reflections
Radiation source: fine-focus sealed tube	4407 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.028$
Detector resolution: 7.3 pixels mm^{-1}	$\theta_{\text{max}} = 29.2^\circ$

$T = 296(2)$ K $\theta_{\min} = 1.7^\circ$
 ω scans $h = -16 \rightarrow 16$
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005) $k = -13 \rightarrow 12$
 $T_{\min} = 0.749$, $T_{\max} = 0.820$ $l = -23 \rightarrow 23$
 24401 measured reflections

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.025$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.066$ $w = 1/[\sigma^2(F_o^2) + (0.0299P)^2 + 0.9401P]$
 $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.04$ $(\Delta/\sigma)_{\max} = 0.003$
 5524 reflections $\Delta\rho_{\max} = 1.12 \text{ e } \text{\AA}^{-3}$
 280 parameters $\Delta\rho_{\min} = -0.41 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.268225 (12)	0.389388 (13)	1.008997 (8)	0.03310 (5)
O1	0.22028 (13)	0.22099 (15)	1.07319 (9)	0.0415 (4)
O2	0.31515 (18)	0.61079 (15)	1.01551 (11)	0.0514 (4)
O3	0.11847 (15)	-0.01098 (16)	1.07144 (11)	0.0528 (4)
N1	0.23627 (17)	0.48884 (18)	1.12248 (11)	0.0425 (4)
N2	0.2877 (2)	0.6145 (2)	1.14445 (14)	0.0565 (6)
N3	0.31169 (16)	0.43881 (18)	0.87174 (11)	0.0383 (4)
N4	0.29433 (16)	0.18153 (18)	0.92055 (11)	0.0408 (4)
C1	0.13973 (19)	0.2149 (2)	1.11147 (13)	0.0387 (5)
C2	0.0823 (2)	0.0913 (2)	1.11224 (14)	0.0442 (5)
C3	-0.0043 (2)	0.0808 (3)	1.15032 (17)	0.0594 (7)

supplementary materials

H3	-0.0432	0.0001	1.1482	0.071*
C4	-0.0336 (3)	0.1904 (3)	1.19174 (18)	0.0671 (8)
H4	-0.0921	0.1829	1.2171	0.081*
C5	0.0230 (2)	0.3077 (3)	1.19522 (16)	0.0572 (7)
H5	0.0041	0.3795	1.2244	0.069*
C6	0.1101 (2)	0.3234 (2)	1.15545 (13)	0.0425 (5)
C7	0.1685 (2)	0.4502 (3)	1.16414 (14)	0.0467 (6)
H7	0.1559	0.5090	1.2033	0.056*
C8	0.3242 (2)	0.6617 (3)	1.08479 (16)	0.0556 (7)
H8	0.3620	0.7434	1.0939	0.067*
C9	0.1069 (2)	0.4231 (3)	0.94090 (17)	0.0460 (6)
H9A	0.056 (3)	0.407 (3)	0.969 (2)	0.069*
H9B	0.093 (3)	0.370 (3)	0.896 (2)	0.069*
H9C	0.099 (2)	0.506 (3)	0.9240 (18)	0.069*
C10	0.4361 (2)	0.3463 (3)	1.05755 (18)	0.0523 (6)
H10A	0.477 (3)	0.366 (3)	1.027 (2)	0.078*
H10B	0.444 (3)	0.252 (3)	1.0710 (18)	0.078*
H10C	0.463 (3)	0.399 (3)	1.104 (2)	0.078*
C11	0.0699 (3)	-0.1410 (3)	1.07460 (19)	0.0639 (8)
H11A	0.1023	-0.2040	1.0441	0.096*
H11B	-0.0086	-0.1358	1.0527	0.096*
H11C	0.0835	-0.1707	1.1292	0.096*
C12	0.2997 (2)	0.5636 (2)	0.84114 (15)	0.0469 (6)
H12	0.2818	0.6325	0.8727	0.056*
C13	0.3121 (2)	0.5960 (2)	0.76579 (15)	0.0510 (6)
H13	0.3021	0.6840	0.7468	0.061*
C14	0.3398 (2)	0.4948 (3)	0.71961 (15)	0.0557 (7)
H14	0.3498	0.5134	0.6687	0.067*
C15	0.3526 (2)	0.3653 (3)	0.74940 (14)	0.0504 (6)
H15	0.3717	0.2957	0.7188	0.060*
C16	0.33668 (17)	0.3393 (2)	0.82525 (12)	0.0363 (4)
C17	0.34066 (18)	0.2002 (2)	0.85795 (13)	0.0378 (5)
C18	0.3865 (2)	0.0940 (3)	0.82431 (16)	0.0530 (6)
H18	0.4185	0.1088	0.7811	0.064*
C19	0.3843 (3)	-0.0340 (3)	0.85548 (18)	0.0630 (8)
H19	0.4156	-0.1060	0.8338	0.076*
C20	0.3361 (3)	-0.0541 (3)	0.91812 (17)	0.0580 (7)
H20	0.3328	-0.1399	0.9393	0.070*
C21	0.2919 (2)	0.0563 (2)	0.94976 (16)	0.0506 (6)
H21	0.2593	0.0429	0.9928	0.061*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.03738 (9)	0.02873 (8)	0.03378 (8)	-0.00102 (6)	0.00941 (6)	-0.00056 (5)
O1	0.0526 (9)	0.0335 (8)	0.0437 (9)	0.0000 (7)	0.0223 (7)	0.0018 (6)
O2	0.0729 (12)	0.0353 (8)	0.0494 (10)	-0.0122 (8)	0.0214 (9)	-0.0058 (7)
O3	0.0625 (11)	0.0398 (9)	0.0580 (11)	-0.0116 (8)	0.0180 (9)	0.0027 (8)

N1	0.0546 (12)	0.0355 (9)	0.0383 (10)	-0.0012 (8)	0.0128 (9)	-0.0050 (8)
N2	0.0823 (17)	0.0419 (11)	0.0469 (12)	-0.0131 (10)	0.0181 (12)	-0.0141 (9)
N3	0.0462 (11)	0.0341 (9)	0.0345 (9)	-0.0023 (8)	0.0091 (8)	0.0016 (7)
N4	0.0515 (11)	0.0322 (9)	0.0417 (10)	0.0015 (8)	0.0171 (9)	0.0008 (8)
C1	0.0434 (12)	0.0430 (12)	0.0301 (10)	-0.0003 (9)	0.0093 (9)	0.0070 (9)
C2	0.0502 (14)	0.0445 (13)	0.0379 (12)	-0.0018 (10)	0.0101 (11)	0.0088 (9)
C3	0.0587 (17)	0.0640 (17)	0.0588 (17)	-0.0135 (13)	0.0204 (14)	0.0175 (13)
C4	0.0664 (19)	0.076 (2)	0.0713 (19)	0.0012 (15)	0.0403 (16)	0.0179 (16)
C5	0.0649 (17)	0.0664 (18)	0.0490 (15)	0.0104 (14)	0.0311 (13)	0.0086 (12)
C6	0.0502 (13)	0.0465 (13)	0.0336 (11)	0.0046 (10)	0.0154 (10)	0.0057 (9)
C7	0.0622 (16)	0.0455 (13)	0.0354 (12)	0.0060 (11)	0.0176 (11)	-0.0026 (10)
C8	0.0755 (19)	0.0366 (12)	0.0558 (16)	-0.0149 (12)	0.0172 (14)	-0.0122 (11)
C9	0.0414 (13)	0.0469 (13)	0.0488 (15)	0.0023 (11)	0.0085 (11)	0.0030 (11)
C10	0.0409 (14)	0.0609 (16)	0.0540 (16)	0.0022 (12)	0.0089 (12)	0.0071 (13)
C11	0.075 (2)	0.0450 (14)	0.0666 (18)	-0.0154 (13)	0.0046 (15)	0.0119 (13)
C12	0.0601 (16)	0.0368 (11)	0.0420 (13)	-0.0029 (11)	0.0080 (11)	0.0053 (10)
C13	0.0586 (16)	0.0459 (13)	0.0443 (13)	-0.0095 (11)	0.0030 (12)	0.0123 (10)
C14	0.0630 (17)	0.0678 (17)	0.0354 (13)	-0.0094 (13)	0.0095 (12)	0.0131 (12)
C15	0.0595 (16)	0.0587 (15)	0.0349 (12)	-0.0012 (12)	0.0147 (11)	-0.0012 (10)
C16	0.0359 (11)	0.0404 (11)	0.0324 (10)	-0.0029 (9)	0.0072 (9)	-0.0001 (9)
C17	0.0413 (12)	0.0374 (11)	0.0341 (11)	0.0010 (9)	0.0075 (9)	-0.0027 (8)
C18	0.0666 (18)	0.0526 (15)	0.0438 (14)	0.0124 (12)	0.0208 (13)	-0.0028 (11)
C19	0.086 (2)	0.0432 (14)	0.0623 (17)	0.0189 (14)	0.0222 (16)	-0.0097 (12)
C20	0.079 (2)	0.0317 (12)	0.0644 (17)	0.0067 (12)	0.0197 (15)	-0.0009 (12)
C21	0.0698 (17)	0.0313 (11)	0.0570 (15)	0.0013 (11)	0.0278 (13)	0.0019 (10)

Geometric parameters (\AA , $^\circ$)

Sn1—C9	2.097 (3)	C7—H7	0.9300
Sn1—C10	2.098 (3)	C8—H8	0.9300
Sn1—O1	2.1572 (14)	C9—H9A	0.90 (3)
Sn1—O2	2.2658 (15)	C9—H9B	0.92 (3)
Sn1—N1	2.2980 (18)	C9—H9C	0.87 (3)
Sn1—N3	2.5825 (18)	C10—H10A	0.84 (3)
Sn1—N4	2.6231 (19)	C10—H10B	0.96 (3)
O1—C1	1.316 (2)	C10—H10C	0.95 (3)
O2—C8	1.273 (3)	C11—H11A	0.9600
O3—C2	1.365 (3)	C11—H11B	0.9600
O3—C11	1.428 (3)	C11—H11C	0.9600
N1—C7	1.280 (3)	C12—C13	1.376 (3)
N1—N2	1.410 (3)	C12—H12	0.9300
N2—C8	1.298 (3)	C13—C14	1.369 (4)
N3—C12	1.338 (3)	C13—H13	0.9300
N3—C16	1.349 (3)	C14—C15	1.378 (3)
N4—C21	1.341 (3)	C14—H14	0.9300
N4—C17	1.343 (3)	C15—C16	1.385 (3)
C1—C6	1.410 (3)	C15—H15	0.9300
C1—C2	1.417 (3)	C16—C17	1.486 (3)
C2—C3	1.383 (3)	C17—C18	1.383 (3)

supplementary materials

C3—C4	1.391 (4)	C18—C19	1.379 (4)
C3—H3	0.9300	C18—H18	0.9300
C4—C5	1.352 (4)	C19—C20	1.360 (4)
C4—H4	0.9300	C19—H19	0.9300
C5—C6	1.412 (3)	C20—C21	1.389 (3)
C5—H5	0.9300	C20—H20	0.9300
C6—C7	1.441 (4)	C21—H21	0.9300
C9—Sn1—C10	169.76 (11)	N1—C7—H7	117.2
C9—Sn1—O1	93.97 (9)	C6—C7—H7	117.2
C10—Sn1—O1	90.20 (9)	O2—C8—N2	128.5 (2)
C9—Sn1—O2	94.23 (9)	O2—C8—H8	115.7
C10—Sn1—O2	87.60 (11)	N2—C8—H8	115.7
O1—Sn1—O2	145.23 (6)	Sn1—C9—H9A	111 (2)
C9—Sn1—N1	94.39 (9)	Sn1—C9—H9B	110 (2)
C10—Sn1—N1	95.65 (10)	H9A—C9—H9B	110 (3)
O1—Sn1—N1	77.12 (6)	Sn1—C9—H9C	110 (2)
O2—Sn1—N1	68.60 (6)	H9A—C9—H9C	109 (3)
C9—Sn1—N3	80.37 (9)	H9B—C9—H9C	106 (3)
C10—Sn1—N3	90.33 (9)	Sn1—C10—H10A	113 (2)
O1—Sn1—N3	138.83 (6)	Sn1—C10—H10B	109.5 (19)
O2—Sn1—N3	75.90 (6)	H10A—C10—H10B	109 (3)
N1—Sn1—N3	143.65 (6)	Sn1—C10—H10C	110 (2)
O1—Sn1—N4	76.91 (6)	H10A—C10—H10C	105 (3)
O2—Sn1—N4	136.43 (6)	H10B—C10—H10C	110 (3)
N1—Sn1—N4	153.65 (6)	O3—C11—H11A	109.5
N3—Sn1—N4	62.68 (6)	O3—C11—H11B	109.5
N4—Sn1—C9	91.77 (9)	H11A—C11—H11B	109.5
N4—Sn1—C10	80.06 (10)	O3—C11—H11C	109.5
Sn1—N4—C17	119.20 (14)	H11A—C11—H11C	109.5
Sn1—N4—C21	119.56 (14)	H11B—C11—H11C	109.5
C17—N4—C21	118.52 (20)	N3—C12—C13	124.0 (2)
C1—O1—Sn1	128.33 (14)	N3—C12—H12	118.0
C8—O2—Sn1	113.35 (15)	C13—C12—H12	118.0
C2—O3—C11	117.4 (2)	C14—C13—C12	118.1 (2)
C7—N1—N2	115.4 (2)	C14—C13—H13	120.9
C7—N1—Sn1	127.50 (16)	C12—C13—H13	120.9
N2—N1—Sn1	116.77 (14)	C13—C14—C15	119.3 (2)
C8—N2—N1	109.0 (2)	C13—C14—H14	120.3
C12—N3—C16	117.6 (2)	C15—C14—H14	120.3
C12—N3—Sn1	120.40 (16)	C14—C15—C16	119.5 (2)
C16—N3—Sn1	121.75 (14)	C14—C15—H15	120.2
C21—N4—C17	118.52 (19)	C16—C15—H15	120.2
O1—C1—C6	123.5 (2)	N3—C16—C15	121.5 (2)
O1—C1—C2	119.0 (2)	N3—C16—C17	116.54 (19)
C6—C1—C2	117.4 (2)	C15—C16—C17	121.9 (2)
O3—C2—C3	124.7 (2)	N4—C17—C18	121.4 (2)
O3—C2—C1	114.3 (2)	N4—C17—C16	116.58 (19)
C3—C2—C1	121.0 (2)	C18—C17—C16	122.0 (2)
C2—C3—C4	120.3 (3)	C19—C18—C17	119.4 (2)

C2—C3—H3	119.9	C19—C18—H18	120.3
C4—C3—H3	119.9	C17—C18—H18	120.3
C5—C4—C3	120.1 (2)	C20—C19—C18	119.6 (2)
C5—C4—H4	120.0	C20—C19—H19	120.2
C3—C4—H4	120.0	C18—C19—H19	120.2
C4—C5—C6	121.3 (3)	C19—C20—C21	118.6 (2)
C4—C5—H5	119.4	C19—C20—H20	120.7
C6—C5—H5	119.4	C21—C20—H20	120.7
C1—C6—C5	119.8 (2)	N4—C21—C20	122.5 (2)
C1—C6—C7	122.3 (2)	N4—C21—H21	118.7
C5—C6—C7	117.8 (2)	C20—C21—H21	118.7
N1—C7—C6	125.5 (2)		
C9—Sn1—O1—C1	-51.1 (2)	O3—C2—C3—C4	178.7 (3)
C10—Sn1—O1—C1	138.3 (2)	C1—C2—C3—C4	-2.8 (4)
O2—Sn1—O1—C1	52.2 (2)	C2—C3—C4—C5	-0.3 (5)
N1—Sn1—O1—C1	42.51 (18)	C3—C4—C5—C6	1.9 (5)
N3—Sn1—O1—C1	-131.00 (17)	O1—C1—C6—C5	-179.6 (2)
C9—Sn1—O2—C8	108.9 (2)	C2—C1—C6—C5	-2.6 (3)
C10—Sn1—O2—C8	-81.2 (2)	O1—C1—C6—C7	-1.9 (4)
O1—Sn1—O2—C8	5.7 (3)	C2—C1—C6—C7	175.1 (2)
N1—Sn1—O2—C8	15.83 (19)	C4—C5—C6—C1	-0.4 (4)
N3—Sn1—O2—C8	-172.1 (2)	C4—C5—C6—C7	-178.1 (3)
C9—Sn1—N1—C7	63.7 (2)	N2—N1—C7—C6	-177.0 (2)
C10—Sn1—N1—C7	-118.3 (2)	Sn1—N1—C7—C6	10.3 (4)
O1—Sn1—N1—C7	-29.4 (2)	C1—C6—C7—N1	14.3 (4)
O2—Sn1—N1—C7	156.5 (2)	C5—C6—C7—N1	-168.0 (2)
N3—Sn1—N1—C7	143.42 (19)	Sn1—O2—C8—N2	-16.6 (4)
C9—Sn1—N1—N2	-108.85 (18)	N1—N2—C8—O2	2.1 (4)
C10—Sn1—N1—N2	69.17 (19)	C16—N3—C12—C13	-0.5 (4)
O1—Sn1—N1—N2	158.07 (18)	Sn1—N3—C12—C13	-174.4 (2)
O2—Sn1—N1—N2	-16.02 (17)	N3—C12—C13—C14	-0.8 (4)
N3—Sn1—N1—N2	-29.1 (2)	C12—C13—C14—C15	0.9 (4)
C7—N1—N2—C8	-159.7 (2)	C13—C14—C15—C16	0.3 (4)
Sn1—N1—N2—C8	13.8 (3)	C12—N3—C16—C15	1.8 (3)
C9—Sn1—N3—C12	70.26 (19)	Sn1—N3—C16—C15	175.57 (18)
C10—Sn1—N3—C12	-114.1 (2)	C12—N3—C16—C17	-175.2 (2)
O1—Sn1—N3—C12	155.26 (17)	Sn1—N3—C16—C17	-1.5 (3)
O2—Sn1—N3—C12	-26.62 (18)	C14—C15—C16—N3	-1.7 (4)
N1—Sn1—N3—C12	-14.0 (2)	C14—C15—C16—C17	175.1 (2)
C9—Sn1—N3—C16	-103.34 (18)	C21—N4—C17—C18	-0.8 (4)
C10—Sn1—N3—C16	72.34 (18)	C21—N4—C17—C16	177.2 (2)
O1—Sn1—N3—C16	-18.3 (2)	N3—C16—C17—N4	15.3 (3)
O2—Sn1—N3—C16	159.78 (18)	C15—C16—C17—N4	-161.8 (2)
N1—Sn1—N3—C16	172.37 (15)	N3—C16—C17—C18	-166.7 (2)
Sn1—O1—C1—C6	-36.9 (3)	C15—C16—C17—C18	16.3 (4)
Sn1—O1—C1—C2	146.18 (17)	N4—C17—C18—C19	0.2 (4)
C11—O3—C2—C3	-5.9 (4)	C16—C17—C18—C19	-177.7 (3)
C11—O3—C2—C1	175.5 (2)	C17—C18—C19—C20	0.8 (5)
O1—C1—C2—O3	-0.1 (3)	C18—C19—C20—C21	-1.1 (5)


supplementary materials

C6—C1—C2—O3	−177.2 (2)	C17—N4—C21—C20	0.5 (4)
O1—C1—C2—C3	−178.7 (2)	C19—C20—C21—N4	0.5 (5)
C6—C1—C2—C3	4.2 (4)		

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C12—H12···O2	0.93	2.40	2.993 (3)	121
C21—H21···O1	0.93	2.36	2.968 (3)	123
C21—H21···O3	0.93	2.50	3.390 (3)	161

Fig. 1



supplementary materials

Fig. 2

