

2-Thienylcarbonylmethylene-triphenyl-phosphorane ylide

Seyyed Javad Sabounchei,^{a*} Vida Jodaian^a and Hamid Reza Khavasi^b

^aFaculty of Chemistry, Bu-Ali Sina University, Hamadan 65174, Iran, and

^bDepartment of Chemistry, Shahid Beheshti University, Evin, Tehran 1983963113,

Iran

Correspondence e-mail: jsabounchei@yahoo.co.uk


Received 21 January 2008; accepted 26 February 2008

Key indicators: single-crystal X-ray study; $T = 120\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.043; wR factor = 0.116; data-to-parameter ratio = 16.9.

In the molecule of the title compound, (2-thienylcarbonyl)-(triphenylphosphonio)methanide, $\text{C}_{24}\text{H}_{19}\text{OPS}$, the geometry around the P atom is nearly tetrahedral and the $\text{O}-\text{C}-\text{C}-\text{P}$ torsion angle is $2.80(3)^\circ$. The thiophene ring is twisted through an angle of $4.33(4)^\circ$ with respect to the plane of the carbonyl group. Inter- and intramolecular hydrogen bonds and $\text{C}-\text{H} \cdots \pi$ interactions are present in the crystal structure.

Related literature

For related literature, see: Allen *et al.* (1987); Bart (1969); Dunitz (1979).



Experimental

Crystal data

$\text{C}_{24}\text{H}_{19}\text{OPS}$	$V = 1971.6(5)\text{ \AA}^3$
$M_r = 386.43$	$Z = 4$
Monoclinic, Cc	$\text{Mo } K\alpha$ radiation
$a = 11.3076(17)\text{ \AA}$	$\mu = 0.26\text{ mm}^{-1}$
$b = 15.474(2)\text{ \AA}$	$T = 120(2)\text{ K}$
$c = 11.3540(16)\text{ \AA}$	$0.4 \times 0.25 \times 0.2\text{ mm}$
$\beta = 97.063(12)^\circ$	

Data collection

Stoe IPDSII diffractometer	7138 measured reflections
Absorption correction: numerical (<i>X-RED32</i> ; Stoe & Cie, 2005)	4130 independent reflections
$T_{\min} = 0.930$, $T_{\max} = 0.950$	4082 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.116$	$\Delta\rho_{\max} = 0.73\text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\min} = -0.48\text{ e \AA}^{-3}$
4130 reflections	Absolute structure: Flack (1983),
244 parameters	3361 Friedel pairs
2 restraints	Flack parameter: 0.03 (7)

Table 1
Selected geometric parameters (\AA , $^\circ$).

$\text{C}_6-\text{P}1$	$1.727(2)$	$\text{C}_{13}-\text{P}1$	$1.812(2)$
$\text{C}_7-\text{P}1$	$1.806(2)$	$\text{C}_{19}-\text{P}1$	$1.816(2)$
$\text{C}_6-\text{P}1-\text{C}_7$	$106.06(10)$	$\text{C}_6-\text{P}1-\text{C}_{19}$	$112.72(10)$
$\text{C}_6-\text{P}1-\text{C}_{13}$	$117.09(12)$	$\text{C}_7-\text{P}1-\text{C}_{19}$	$108.67(10)$
$\text{C}_7-\text{P}1-\text{C}_{13}$	$106.16(10)$	$\text{C}_{13}-\text{P}1-\text{C}_{19}$	$105.74(10)$

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{C}9-\text{H}9 \cdots \text{O}1^i$	0.93	2.34	3.147 (3)	145
$\text{C}14-\text{H}14 \cdots \text{O}1$	0.93	2.52	3.187 (3)	129
$\text{C}3-\text{H}3 \cdots \text{Cg}1^{ii}$	0.93	2.82	3.582 (3)	140
$\text{C}8-\text{H}8 \cdots \text{Cg}2^{ii}$	0.93	2.80	3.592 (2)	143
$\text{C}10-\text{H}10 \cdots \text{Cg}2^i$	0.93	2.95	3.734 (3)	143
$\text{C}23-\text{H}23 \cdots \text{Cg}3^{iii}$	0.93	2.82	3.465 (3)	127

Symmetry code: (i) $x - \frac{1}{2}, y - \frac{1}{2}, z$; (ii) $x, -y, z + \frac{1}{2}$; (iii) $x, -y, z - \frac{1}{2}$. $\text{Cg}1$ is the centroid of atoms $\text{C}7-\text{C}12$, $\text{Cg}2$ is the centroid of atoms $\text{C}19-\text{C}24$ and $\text{Cg}3$ is the centroid of atoms $\text{C}13-\text{C}18$.

Data collection: *X-AREA* (Stoe & Cie, 2005); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge Bu-Ali Sina University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2064).

References

- Allen, F. H., Kennard, O., Waston, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Bart, J. C. J. (1969). *J. Chem. Soc. B*, pp. 350–365.
- Dunitz, J. D. (1979). *X-ray Analysis and the Structure of Organic Molecules*. Ithaca: Cornell University Press.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2005). *X-RED32*, *X-AREA* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2008). E64, o647 [doi:10.1107/S1600536808005497]

2-Thienylcarbonylmethylene-triphenylphosphorane ylide

S. J. Sabounchei, V. Jodaian and H. R. Khavasi

Comment

Phosphoranes of the type $(C_6H_5)_3PCHCOCH_3S$ (TPPY) can coordinate to metals through either C or O atoms. The crystal and molecular structure of this ylide was determined successfully. The structural investigation with metrical parameters for the title compound, (I), show that how they vary with a change in delocalization in the metal derivatives as well as in other resonance stabilized ylides. In this molecule, the bond lengths and angles (Table 1) are generally within normal ranges (Allen *et al.*, 1987).

The P1—C6 bond length [1.727 (2) Å], is shorter than the other P—C bonds (Table 1) and longer than the equivalent bond lengths of 1.66 Å reported for methylenetriphenylphosphorane (Bart, 1969), which shows partial double-bond character for these two bonds.

The C4—S1 and C1—S1 bond lengths of 1.712 (3) Å and 1.719 (2) Å are longer than the other C—C bonds. These bond distances suggest resonance delocalization in the molecule (Fig. 2). The resonance formulation is supported by the near planarity of P1, C6, C5 and O1 in TPPY. The torsion angle O1—C2—C1—P1 of 2.80 (3)° also indicates resonance.

The thiophenyl group is twisted with respect to the plane containing the carbonyl group through angles of 4.33 (4)°. Bond angle of 118.30 (16)° for P1—C6—C5, indicate a distorted trigonal arrangement about C6. The non-bonded distances P1—O1 of 2.990 (3) Å of TPPY is significantly shorter than the sum of the van der Waals radii of P and O (3.3 Å) (Dunitz, 1979), indicating a strong intramolecular interaction between P^+ and O^- charge centers, which leads to the *cis* orientation. Packing diagram of TPPY is shown in Fig. 3. As it is clear from this diagram, there are some C—H···O inter- and intra-molecular interactions that seem to be effective in stability of packing (Table 2). There are four remarkable C—H···Cg (pi-ring) interactions; $[H3\cdots Cg1(C7/12)]^i = 2.82$ Å and $C3—H3\cdots Cg1 = 140^\circ$, $H8\cdots Cg2(C19/24)]^i = 2.80$ Å and $C8—H8\cdots Cg2 = 143^\circ$, $H10\cdots Cg2(C19/24)]^{ii} = 2.95$ Å and $C10—H10\cdots Cg2 = 143^\circ$, $H23\cdots Cg3(C13/18)]^{iii} = 2.82$ Å and $C23—H23\cdots Cg1 = 127^\circ$, with symmetry codes; (i) $X, -Y, 1/2+Z$, (ii) $-1/2+X, -1/2+Y, Z$ and (iii) $X, -Y, -1/2+Z$ which are effective in the stabilization of crystal packing.

Experimental

The title compound was prepared by addition of 2-bromo-acetothiophen (0.102 g, 0.5 mmol) in chloroform (25 ml) to a solution of triphenylphosphine (0.131 g, 0.5 mmol) in the same solvent (5 ml). The resulting pale pink solution was stirred for 12 h. The solution was concentrated under reduced pressure to 5 ml, and diethyl ether (20 ml) was added. The yellow solid formed was filtered off, washed with petroleum diethyl ether (10 ml), and dried under reduced pressure. In order to get the final product, all of the crude solid was transferred to an alkaline solution of 5% NaOH and stirred at 310 K for about 14 h, yielding the white precipitate. The product was washed several times with distilled water and air dried. The resulting solid was recrystallized from a chloroform-diethyl ether mixture (m.p. 496–498 K). Yield: 78%, 0.301 g.

supplementary materials

Refinement

H atoms were positioned geometrically, with C—H=0.93 Å for aromatic and methine H and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure with the atom-numbering scheme. Displacement ellipsoids are drawn at 30% probability level.



Fig. 2. Resonance in the TPPY.




Fig. 3. The packing of (I). Hydrogen bonds are shown as dashed lines. H atoms not participate in H bonding are omitted for clarity. Symmetry code: (i) $-1/2 + x, -1/2 + y, z$.

(2-Thienylcarbonyl)(triphenylphosphonio)methanide

Crystal data

C ₂₄ H ₁₉ O ₁ P ₁ S ₁	$F_{000} = 808$
$M_r = 386.43$	$D_x = 1.302 \text{ Mg m}^{-3}$
Monoclinic, <i>Cc</i>	Mo $K\alpha$ radiation
Hall symbol: C -2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 11.3076 (17) \text{ \AA}$	Cell parameters from 2000 reflections
$b = 15.474 (2) \text{ \AA}$	$\theta = 2.2\text{--}27.9^\circ$
$c = 11.3540 (16) \text{ \AA}$	$\mu = 0.26 \text{ mm}^{-1}$
$\beta = 97.063 (12)^\circ$	$T = 120 (2) \text{ K}$
$V = 1971.6 (5) \text{ \AA}^3$	Prism, colorless
$Z = 4$	$0.4 \times 0.25 \times 0.2 \text{ mm}$

Data collection

Stoe IPDSII diffractometer	$R_{\text{int}} = 0.059$
rotation method scans	$\theta_{\text{max}} = 25.9^\circ$

Absorption correction: numerical (X-RED32; Stoe & Cie, 2005)	$\theta_{\min} = 2.2^\circ$
$T_{\min} = 0.930$, $T_{\max} = 0.950$	$h = -14 \rightarrow 14$
7138 measured reflections	$k = -20 \rightarrow 20$
4130 independent reflections	$l = -14 \rightarrow 14$
4082 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.0801P)^2 + 1.5059P]$ where $P = (F_o^2 + 2F_c^2)/3$
$R[F^2 > 2\sigma(F^2)] = 0.042$	$(\Delta/\sigma)_{\max} = 0.007$
$wR(F^2) = 0.116$	$\Delta\rho_{\max} = 0.74 \text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\min} = -0.48 \text{ e \AA}^{-3}$
4130 reflections	Extinction correction: none
244 parameters	Absolute structure: Flack (1983), 3361 Friedel pairs
2 restraints	Flack parameter: 0.03 (7)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0560 (3)	0.37507 (16)	0.7865 (2)	0.0305 (5)
H1	0.0667	0.4255	0.8311	0.037*
C2	-0.0358 (2)	0.31826 (17)	0.7929 (2)	0.0297 (5)
H2	-0.0942	0.3254	0.8431	0.036*
C3	-0.0319 (2)	0.24653 (15)	0.7135 (2)	0.0262 (5)
H3	-0.0875	0.2019	0.7058	0.031*
C4	0.0640 (2)	0.25175 (14)	0.6502 (2)	0.0232 (4)
C5	0.1026 (2)	0.19364 (13)	0.5570 (2)	0.0224 (4)
C6	0.0300 (2)	0.12304 (14)	0.5188 (2)	0.0221 (4)
H6	-0.0331	0.1068	0.5593	0.026*
C7	-0.0633 (2)	-0.00552 (13)	0.3568 (2)	0.0208 (4)
C8	-0.0838 (2)	-0.06819 (14)	0.4413 (2)	0.0243 (5)
H8	-0.0315	-0.0733	0.5109	0.029*
C9	-0.1822 (2)	-0.12279 (15)	0.4217 (2)	0.0272 (5)
H9	-0.196	-0.1638	0.4783	0.033*
C10	-0.2600 (2)	-0.11563 (15)	0.3166 (2)	0.0274 (5)
H10	-0.3262	-0.1516	0.3034	0.033*
C11	-0.2385 (2)	-0.05476 (15)	0.2316 (2)	0.0295 (5)

supplementary materials

H11	-0.29	-0.0507	0.1613	0.035*
C12	-0.1407 (2)	0.00010 (14)	0.2511 (2)	0.0240 (4)
H12	-0.1267	0.0406	0.1939	0.029*
C13	0.1935 (2)	0.00050 (14)	0.4095 (2)	0.0212 (4)
C14	0.2881 (2)	0.01748 (15)	0.4970 (2)	0.0277 (5)
H14	0.2825	0.0619	0.5511	0.033*
C15	0.3913 (3)	-0.03241 (18)	0.5029 (3)	0.0340 (5)
H15	0.4547	-0.0212	0.5611	0.041*
C16	0.3994 (2)	-0.09902 (18)	0.4220 (3)	0.0345 (6)
H16	0.4685	-0.1321	0.4259	0.041*
C17	0.3045 (3)	-0.11623 (17)	0.3350 (2)	0.0327 (6)
H17	0.3102	-0.1607	0.281	0.039*
C18	0.2014 (3)	-0.06698 (14)	0.3290 (2)	0.0272 (5)
H18	0.1376	-0.0789	0.2714	0.033*
C19	0.0732 (2)	0.13744 (14)	0.26848 (19)	0.0210 (4)
C20	-0.0162 (2)	0.19924 (14)	0.2397 (2)	0.0233 (4)
H20	-0.0793	0.2037	0.2847	0.028*
C21	-0.0099 (2)	0.25406 (15)	0.1430 (2)	0.0280 (5)
H21	-0.0703	0.294	0.1219	0.034*
C22	0.0866 (3)	0.24901 (15)	0.0782 (2)	0.0284 (5)
H22	0.0913	0.2864	0.0149	0.034*
C23	0.1750 (2)	0.18886 (16)	0.1075 (2)	0.0269 (5)
H23	0.2394	0.1859	0.064	0.032*
C24	0.1684 (2)	0.13204 (15)	0.2025 (2)	0.0238 (4)
H24	0.2276	0.0908	0.2213	0.029*
O1	0.19781 (17)	0.21153 (10)	0.51549 (16)	0.0280 (4)
P1	0.06067 (5)	0.06711 (3)	0.39443 (5)	0.01906 (13)
S1	0.14871 (6)	0.34234 (4)	0.68674 (6)	0.02845 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0429 (15)	0.0251 (10)	0.0223 (11)	0.0022 (10)	-0.0003 (10)	-0.0063 (8)
C2	0.0360 (14)	0.0310 (12)	0.0225 (11)	0.0044 (10)	0.0055 (10)	-0.0033 (9)
C3	0.0328 (13)	0.0238 (10)	0.0224 (10)	0.0021 (9)	0.0053 (9)	-0.0005 (8)
C4	0.0281 (11)	0.0195 (9)	0.0218 (10)	-0.0002 (8)	0.0028 (9)	-0.0007 (7)
C5	0.0286 (11)	0.0182 (9)	0.0208 (10)	-0.0013 (8)	0.0042 (9)	0.0008 (7)
C6	0.0279 (12)	0.0186 (9)	0.0205 (10)	-0.0044 (8)	0.0063 (9)	-0.0017 (7)
C7	0.0244 (10)	0.0163 (9)	0.0218 (10)	-0.0007 (7)	0.0032 (8)	-0.0018 (7)
C8	0.0301 (13)	0.0212 (10)	0.0220 (11)	0.0002 (8)	0.0044 (9)	0.0001 (7)
C9	0.0341 (13)	0.0218 (10)	0.0274 (11)	-0.0056 (9)	0.0105 (10)	-0.0016 (8)
C10	0.0248 (11)	0.0218 (10)	0.0365 (13)	-0.0043 (8)	0.0074 (10)	-0.0063 (9)
C11	0.0287 (13)	0.0251 (10)	0.0331 (12)	0.0011 (9)	-0.0029 (10)	-0.0026 (9)
C12	0.0261 (12)	0.0206 (10)	0.0246 (10)	0.0003 (8)	0.0006 (9)	0.0015 (8)
C13	0.0229 (11)	0.0192 (9)	0.0218 (10)	0.0000 (8)	0.0037 (8)	0.0036 (8)
C14	0.0293 (13)	0.0253 (10)	0.0276 (11)	-0.0017 (9)	-0.0001 (10)	0.0018 (9)
C15	0.0291 (13)	0.0361 (13)	0.0357 (13)	0.0024 (10)	-0.0001 (11)	0.0084 (11)
C16	0.0282 (13)	0.0358 (13)	0.0407 (14)	0.0092 (10)	0.0097 (11)	0.0125 (11)

C17	0.0407 (15)	0.0290 (11)	0.0294 (12)	0.0107 (10)	0.0075 (11)	0.0021 (9)
C18	0.0329 (14)	0.0247 (12)	0.0243 (12)	0.0048 (9)	0.0047 (10)	0.0001 (8)
C19	0.0252 (11)	0.0192 (9)	0.0186 (10)	-0.0005 (8)	0.0024 (8)	-0.0009 (7)
C20	0.0282 (12)	0.0178 (9)	0.0243 (10)	0.0011 (8)	0.0051 (9)	0.0022 (7)
C21	0.0336 (13)	0.0234 (10)	0.0263 (11)	0.0018 (9)	0.0011 (10)	0.0045 (8)
C22	0.0384 (14)	0.0265 (11)	0.0201 (10)	-0.0044 (10)	0.0020 (10)	0.0042 (8)
C23	0.0323 (12)	0.0293 (11)	0.0200 (10)	-0.0031 (9)	0.0064 (9)	0.0007 (8)
C24	0.0270 (12)	0.0225 (9)	0.0226 (11)	-0.0011 (8)	0.0054 (9)	0.0004 (8)
O1	0.0299 (9)	0.0235 (8)	0.0325 (9)	-0.0059 (6)	0.0108 (8)	-0.0035 (6)
P1	0.0227 (3)	0.0161 (2)	0.0186 (2)	-0.00069 (19)	0.00337 (19)	0.00047 (18)
S1	0.0342 (3)	0.0220 (3)	0.0294 (3)	-0.0054 (2)	0.0047 (2)	-0.0052 (2)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.369 (4)	C13—C14	1.392 (4)
C1—S1	1.712 (3)	C13—C18	1.398 (3)
C1—H1	0.93	C13—P1	1.812 (2)
C2—C3	1.434 (3)	C14—C15	1.394 (4)
C2—H2	0.93	C14—H14	0.93
C3—C4	1.375 (3)	C15—C16	1.392 (4)
C3—H3	0.93	C15—H15	0.93
C4—C5	1.494 (3)	C16—C17	1.392 (4)
C4—S1	1.719 (2)	C16—H16	0.93
C5—O1	1.258 (3)	C17—C18	1.388 (4)
C5—C6	1.403 (3)	C17—H17	0.93
C6—P1	1.727 (2)	C18—H18	0.93
C6—H6	0.93	C19—C24	1.388 (3)
C7—C12	1.399 (3)	C19—C20	1.401 (3)
C7—C8	1.403 (3)	C19—P1	1.816 (2)
C7—P1	1.806 (2)	C20—C21	1.396 (3)
C8—C9	1.393 (4)	C20—H20	0.93
C8—H8	0.93	C21—C22	1.391 (4)
C9—C10	1.398 (4)	C21—H21	0.93
C9—H9	0.93	C22—C23	1.377 (4)
C10—C11	1.391 (4)	C22—H22	0.93
C10—H10	0.93	C23—C24	1.401 (3)
C11—C12	1.391 (3)	C23—H23	0.93
C11—H11	0.93	C24—H24	0.93
C12—H12	0.93		
C2—C1—S1	112.04 (18)	C13—C14—H14	120.1
C2—C1—H1	124	C15—C14—H14	120.1
S1—C1—H1	124	C16—C15—C14	120.1 (3)
C1—C2—C3	112.4 (2)	C16—C15—H15	120
C1—C2—H2	123.8	C14—C15—H15	120
C3—C2—H2	123.8	C17—C16—C15	120.1 (2)
C4—C3—C2	111.7 (2)	C17—C16—H16	119.9
C4—C3—H3	124.1	C15—C16—H16	119.9
C2—C3—H3	124.1	C18—C17—C16	120.0 (2)
C3—C4—C5	130.7 (2)	C18—C17—H17	120

supplementary materials

C3—C4—S1	111.94 (17)	C16—C17—H17	120
C5—C4—S1	117.31 (17)	C17—C18—C13	119.9 (3)
O1—C5—C6	123.4 (2)	C17—C18—H18	120.1
O1—C5—C4	118.1 (2)	C13—C18—H18	120.1
C6—C5—C4	118.5 (2)	C24—C19—C20	120.0 (2)
C5—C6—P1	118.30 (16)	C24—C19—P1	121.98 (18)
C5—C6—H6	120.9	C20—C19—P1	117.98 (16)
P1—C6—H6	120.9	C21—C20—C19	119.6 (2)
C12—C7—C8	119.4 (2)	C21—C20—H20	120.2
C12—C7—P1	123.35 (16)	C19—C20—H20	120.2
C8—C7—P1	117.22 (19)	C22—C21—C20	120.0 (2)
C9—C8—C7	120.4 (2)	C22—C21—H21	120
C9—C8—H8	119.8	C20—C21—H21	120
C7—C8—H8	119.8	C23—C22—C21	120.3 (2)
C8—C9—C10	119.7 (2)	C23—C22—H22	119.9
C8—C9—H9	120.2	C21—C22—H22	119.9
C10—C9—H9	120.2	C22—C23—C24	120.3 (2)
C11—C10—C9	120.1 (2)	C22—C23—H23	119.9
C11—C10—H10	120	C24—C23—H23	119.9
C9—C10—H10	120	C19—C24—C23	119.7 (2)
C12—C11—C10	120.4 (2)	C19—C24—H24	120.1
C12—C11—H11	119.8	C23—C24—H24	120.1
C10—C11—H11	119.8	C6—P1—C7	106.06 (10)
C11—C12—C7	120.0 (2)	C6—P1—C13	117.09 (12)
C11—C12—H12	120	C7—P1—C13	106.16 (10)
C7—C12—H12	120	C6—P1—C19	112.72 (10)
C14—C13—C18	120.2 (2)	C7—P1—C19	108.67 (10)
C14—C13—P1	121.02 (17)	C13—P1—C19	105.74 (10)
C18—C13—P1	118.8 (2)	C1—S1—C4	91.86 (13)
C13—C14—C15	119.7 (2)		
S1—C1—C2—C3	-0.8 (3)	C20—C21—C22—C23	1.3 (4)
C1—C2—C3—C4	0.4 (3)	C21—C22—C23—C24	0.2 (4)
C2—C3—C4—C5	-178.7 (2)	C20—C19—C24—C23	0.3 (3)
C2—C3—C4—S1	0.2 (3)	P1—C19—C24—C23	-178.83 (18)
C3—C4—C5—O1	-177.3 (3)	C22—C23—C24—C19	-1.0 (4)
S1—C4—C5—O1	3.9 (3)	C5—C6—P1—C7	-170.32 (19)
C3—C4—C5—C6	3.8 (4)	C5—C6—P1—C13	71.5 (2)
S1—C4—C5—C6	-175.02 (18)	C5—C6—P1—C19	-51.5 (2)
O1—C5—C6—P1	-9.7 (3)	C12—C7—P1—C6	115.89 (19)
C4—C5—C6—P1	169.12 (17)	C8—C7—P1—C6	-61.6 (2)
C12—C7—C8—C9	-1.7 (3)	C12—C7—P1—C13	-118.89 (19)
P1—C7—C8—C9	175.91 (17)	C8—C7—P1—C13	63.60 (19)
C7—C8—C9—C10	0.7 (3)	C12—C7—P1—C19	-5.6 (2)
C8—C9—C10—C11	0.6 (4)	C8—C7—P1—C19	176.93 (17)
C9—C10—C11—C12	-0.8 (4)	C14—C13—P1—C6	-23.8 (2)
C10—C11—C12—C7	-0.2 (3)	C18—C13—P1—C6	158.29 (17)
C8—C7—C12—C11	1.5 (3)	C14—C13—P1—C7	-141.96 (18)
P1—C7—C12—C11	-176.01 (18)	C18—C13—P1—C7	40.1 (2)
C18—C13—C14—C15	0.8 (3)	C14—C13—P1—C19	102.69 (19)

P1—C13—C14—C15	−177.07 (18)	C18—C13—P1—C19	−75.2 (2)
C13—C14—C15—C16	−0.1 (4)	C24—C19—P1—C6	131.09 (19)
C14—C15—C16—C17	−0.2 (4)	C20—C19—P1—C6	−48.1 (2)
C15—C16—C17—C18	0.0 (4)	C24—C19—P1—C7	−111.6 (2)
C16—C17—C18—C13	0.7 (4)	C20—C19—P1—C7	69.2 (2)
C14—C13—C18—C17	−1.1 (3)	C24—C19—P1—C13	2.0 (2)
P1—C13—C18—C17	176.84 (18)	C20—C19—P1—C13	−177.19 (18)
C24—C19—C20—C21	1.2 (4)	C2—C1—S1—C4	0.7 (2)
P1—C19—C20—C21	−179.61 (18)	C3—C4—S1—C1	−0.5 (2)
C19—C20—C21—C22	−2.0 (4)	C5—C4—S1—C1	178.5 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C9—H9···O1 ⁱ	0.93	2.34	3.147 (3)	145
C14—H14···O1	0.93	2.52	3.187 (3)	129
C3—H3···Cg1 ⁱⁱ	0.93	2.82	3.582 (3)	140
C8—H8···Cg2 ⁱⁱ	0.93	2.80	3.592 (2)	143
C10—H10···Cg2 ⁱ	0.93	2.95	3.734 (3)	143
C23—H23···Cg3 ⁱⁱⁱ	0.93	2.82	3.465 (3)	127

Symmetry codes: (i) $x-1/2, y-1/2, z$; (ii) $x, -y, z+1/2$; (iii) $x, -y, z-1/2$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3

