

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

2-Hydroxy-5-nitrobenzaldehyde  
2,4-dinitrophenylhydrazoneAbdassalam A. Tameem,<sup>a</sup> Bahruddin Saad,<sup>a</sup>  
Abdussalam M. Salhin,<sup>a</sup> Samuel Robinson Jebas<sup>b‡</sup> and  
Hoong-Kun Fun<sup>b\*</sup><sup>a</sup>School of Chemical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and <sup>b</sup>X-ray Crystallography unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia  
Correspondence e-mail: hkfun@usm.my


Received 1 March 2008; accepted 1 March 2008

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.051;  $wR$  factor = 0.146; data-to-parameter ratio = 22.2.

In the title compound,  $\text{C}_{13}\text{H}_9\text{N}_5\text{O}_7$ , one of the nitro groups is twisted away from the attached benzene ring by  $16.21$  ( $8^\circ$ ). The dihedral angle between the two benzene rings is  $4.63$  ( $1^\circ$ ). The molecular structure is stabilized by intramolecular  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bonds which generate an  $S(6)$  ring motif. The molecules pack as layers parallel to the  $ab$  plane; molecules of adjacent layers are linked into chains along the  $[101]$  direction through  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

For related literature, see: Cordis *et al.* (1998); Fun *et al.* (1996); Guillaumont & Nakamura (2000); Hanoune *et al.* (2006); Lamberton *et al.* (1974); Niknam *et al.* (2005); Raj & Kurup (2007); Salhin *et al.* (2007); Shan, Xu *et al.* (2003); Shan, Yu *et al.* (2003); Tameem *et al.* (2007); Uchiyama *et al.* (2003); Vogel *et al.* (2000); Zegota (1999); Zlotorzynska & Lai (1999). For ring motifs, see: Bernstein *et al.* (1995).



## Experimental

## Crystal data

$\text{C}_{13}\text{H}_9\text{N}_5\text{O}_7$   
 $M_r = 347.25$   
 Monoclinic,  $P2_1/n$   
 $a = 12.7543$  (5) Å

$b = 8.1898$  (3) Å  
 $c = 13.8618$  (5) Å  
 $\beta = 112.683$  (2)°  
 $V = 1335.94$  (9) Å<sup>3</sup>

‡ Permanent address: Department of Physics, Karunya University, Karunya Nagar, Coimbatore 641 114, India.

$Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.14$  mm<sup>-1</sup>

$T = 100.0$  (1) K  
 $0.29 \times 0.27 \times 0.11$  mm

## Data collection

Bruker SMART APEXII CCD  
 area-detector diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2005)  
 $T_{\min} = 0.960$ ,  $T_{\max} = 0.985$

24539 measured reflections  
 5195 independent reflections  
 3513 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.056$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$   
 $wR(F^2) = 0.145$   
 $S = 1.09$   
 5195 reflections  
 234 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.47$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.33$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N1}\cdots\text{O4}^i$	0.89 (2)	2.54 (2)	3.0666 (15)	118 (2)
$\text{N1}-\text{H1N1}\cdots\text{O1}$	0.89 (2)	2.07 (2)	2.6477 (15)	121 (2)
$\text{O5}-\text{H1O5}\cdots\text{N2}$	0.92 (3)	1.82 (3)	2.6656 (14)	150 (2)

Symmetry code: (i)  $x + \frac{1}{2}, -y + \frac{5}{2}, z + \frac{1}{2}$ .

Data collection: APEX2 (Bruker, 2005); cell refinement: APEX2; data reduction: SAINT (Bruker, 2005); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2003).

FKH and SRJ thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. SRJ thanks the Universiti Sains Malaysia for the award of a post-doctoral research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2567).

## References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cordis, G. A., Das, D. K. & Riedel, W. (1998). *J. Chromatogr. A*, **798**, 117–123.
- Fun, H.-K., Sivakumar, K., Lu, Z.-L., Duan, C.-Y., Tian, Y.-P. & You, X.-Z. (1996). *Acta Cryst.* **C52**, 986–988.
- Guillaumont, D. & Nakamura, S. (2000). *Dyes Pigm.* **46**, 85–92.
- Hanoune, B., Bris, T. L., Allou, L., Marchand, C. & Calve, S. L. (2006). *Atmos. Environ.* **40**, 5768–5775.
- Lamberton, J. A., Nelson, E. R. & Triffett, C. K. (1974). *Aust. J. Chem.* **27**, 1521–1529.
- Niknam, K., Kiasat, A. R. & Karimi, S. (2005). *Synth. Commun.* **35**, 2231–2236.
- Raj, B. N. B. & Kurup, M. R. P. (2007). *Spectrochim. Acta Part A*, **66**, 898–903.
- Salhin, A., Tameem, A. A., Saad, B., Ng, S.-L. & Fun, H.-K. (2007). *Acta Cryst.* **E63**, o2880.
- Shan, S., Xu, D.-J. & Hu, W.-X. (2003). *Acta Cryst.* **E59**, o1173–o1174.
- Shan, S., Yu, H.-G., Hu, W.-X. & Xu, D.-J. (2003). *Acta Cryst.* **E59**, o1886–o1887.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

Tameem, A. A., Salhin, A., Saad, B., Ng, S.-L. & Fun, H.-K. (2007). *Acta Cryst.* **E63**, o2502.  
Uchiyama, S., Ando, M. & Aoyagi, S. (2003). *J. Chromatogr.* **A996**, 95–102.

Vogel, M., Potter, W. & Karst, U. (2000). *J. Chromatogr.* **A886**, 303–307.  
Zegota, H. (1999). *J. Chromatogr.* **A863**, 227–233.  
Zlotorzynska, E. D. & Lai, E. P. C. (1999). *J. Chromatogr.* **A853**, 487–796.

**supplementary materials**

*Acta Cryst.* (2008). E64, o679-o680 [ doi:10.1107/S1600536808005825 ]

## 2-Hydroxy-5-nitrobenzaldehyde 2,4-dinitrophenylhydrazone

A. A. Tameem, B. Saad, A. M. Salhin, S. R. Jebas and H.-K. Fun

### Comment

Phenylhydrazone derivatives have been synthesized in order to investigate their structures and analytical applications. 2,4-Dinitrophenylhydrazones play an important role as stabilizers for the detection, characterization and protection of the carbonyl group of compounds than phenylhydrazones (Niknam *et al.*, 2005). 2,4-Dinitrophenylhydrazone derivatives are widely used in various forms of analytical chemistry (Lamberton *et al.*, 1974; Zegota, 1999; Cordis *et al.*, 1998; Zlotorzynska & Lai, 1999) and are also used as dyes (Guillaumont & Nakamura, 2000). They are also found to have versatile coordinating abilities towards different metal ions (Raj *et al.*, 2006). In addition, they are used to determine airborne aldehydes and ketones (Vogel *et al.*, 2000) and as detectors of formaldehyde (Hanoune *et al.*, 2006). These compounds can exist as E and Z stereoisomers (Uchiyama *et al.*, 2003). Their existence as a keto tautomer in the solid state with an E configuration across the C—N bond have been reported (Fun *et al.*, 1996). The title compound, whose structure is reported here, is one of the series of phenylhydrazone derivatives that we have prepared; the crystal structures of some of these compounds have been studied previously (Tameem *et al.*, 2007; Salhin *et al.*, 2007).

The bond lengths and angles in the title compound (Fig.1) have normal values. The molecule is nearly planar, with a maximum deviation from the mean plane of 0.487 (1) Å for atom O1. The dihedral angle between the two benzene rings is 4.63 (1)°. The C—N bond lengths in the hydrazone moiety agree well with those reported earlier (Shan, Xu *et al.*, 2003; Shan, Yu *et al.*, 2003). The asymmetry of the exocyclic angles at C7 [C7—C8—C13 = 118.2 (2)° and C7—C8—C9 = 123.3 (2)°] is more pronounced than that at C6 [N1—C6—C1 = 122.7 (1)° and N1—C6—C5 = 120.5 (1)°]. One of the hydrazone N atoms is involved in an O—H···N intramolecular hydrogen bond with the hydroxy group, while the other is involved in an N—H···O intramolecular hydrogen bond with the nitro group. Each of these hydrogen bonds generate an S(6) ring motif (Bernstein *et al.*, 1995).

The molecules are linked into a chain along the [1 0 1] direction through N—H—O hydrogen bonds. The molecules pack as layers parallel to the *ab* plane. Within the layer, weak  $\pi$ - $\pi$  interactions are observed between the C1—C6 and C8—C13 benzene rings, with a centroid-centroid distance of 3.7457 (8) Å.

### Experimental

2,4-Dinitrophenylhydrazine (400 mg, 2 mmol) in concentrated sulfuric acid (5 ml) was slowly added to a solution of 2-hydroxy-5-nitrobenzaldehyde (337 mg, 2 mmol) in ethanol (95%, 20 ml). The mixture was stirred for 15 min, and was left to stand at room temperature for 30 min. The resulting product was filtered and washed with 95% ethanol (20 ml) and the orange powder product was collected. Crystals suitable for X-ray diffraction analysis were grown by slow evaporation of a saturated solution of the resulted compound in ethanol.

## Refinement

O- and N-bound H atoms were located in a difference map and refined isotropically. The remaining H atoms were placed in calculated positions (C—H = 0.93 Å) and refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

## Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atomic numbering scheme. Hydrogen bonds are shown as dashed lines.




Fig. 2. The crystal packing of the title compound, viewed along the *c* axis. Hydrogen bonds are shown as dashed lines.

## 2-Hydroxy-5-nitrobenzaldehyde 2,4-dinitrophenylhydrazone

### Crystal data

$\text{C}_{13}\text{H}_9\text{N}_5\text{O}_7$

$M_r = 347.25$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2_1n$

$a = 12.7543\ (5)\ \text{\AA}$

$b = 8.1898\ (3)\ \text{\AA}$

$c = 13.8618\ (5)\ \text{\AA}$

$\beta = 112.683\ (2)^\circ$

$V = 1335.94\ (9)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 712$

$D_x = 1.726\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3255 reflections

$\theta = 2.8\text{--}33.1^\circ$

$\mu = 0.14\ \text{mm}^{-1}$

$T = 100.0\ (1)\ \text{K}$

Block, orange

$0.29 \times 0.27 \times 0.11\ \text{mm}$

### Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Detector resolution:  $8.33\ \text{pixels mm}^{-1}$

$\omega$  scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\text{min}} = 0.960$ ,  $T_{\text{max}} = 0.985$

24539 measured reflections

3513 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.056$

$\theta_{\text{max}} = 33.4^\circ$

$\theta_{\text{min}} = 2.8^\circ$

$h = -19 \rightarrow 19$

$k = -10 \rightarrow 12$

5195 independent reflections

$l = -21 \rightarrow 20$

*Refinement*

Refinement on  $F^2$

H atoms treated by a mixture of independent and constrained refinement

Least-squares matrix: full

$$w = 1/[\sigma^2(F_o^2) + (0.0708P)^2 + 0.0132P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$R[F^2 > 2\sigma(F^2)] = 0.050$

$(\Delta/\sigma)_{\max} = 0.001$

$wR(F^2) = 0.145$

$\Delta\rho_{\max} = 0.47 \text{ e } \text{\AA}^{-3}$

$S = 1.09$

$\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$

5195 reflections

Extinction correction: none

234 parameters

*Special details*

**Geometry.** Experimental. The low-temperature data was collected with the Oxford Cryosystem Cobra low-temperature attachment.

All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.71240 (8)	0.91583 (13)	0.22343 (8)	0.0228 (2)
O2	0.61439 (9)	1.12841 (14)	0.22968 (8)	0.0282 (2)
O3	0.39947 (10)	1.41385 (15)	-0.08171 (9)	0.0340 (3)
O4	0.45224 (10)	1.41148 (14)	-0.21219 (8)	0.0307 (3)
O5	0.94762 (8)	0.81549 (12)	-0.11896 (7)	0.0189 (2)
O6	1.34597 (9)	0.33913 (14)	0.11735 (8)	0.0300 (3)
O7	1.29887 (9)	0.41281 (13)	0.24496 (8)	0.0259 (2)
N1	0.80661 (9)	0.91651 (14)	0.08340 (9)	0.0164 (2)
N2	0.87737 (9)	0.86266 (13)	0.03680 (8)	0.0157 (2)
N3	0.66026 (9)	1.04241 (14)	0.18460 (8)	0.0187 (2)
N4	0.46053 (10)	1.36459 (15)	-0.12555 (9)	0.0208 (2)
N5	1.28791 (9)	0.41996 (14)	0.15301 (9)	0.0184 (2)
C1	0.64902 (11)	1.08800 (16)	0.07968 (9)	0.0160 (2)
C2	0.56488 (11)	1.20121 (16)	0.02874 (10)	0.0179 (2)
H2A	0.5209	1.2465	0.062	0.021*
C3	0.54779 (10)	1.24505 (16)	-0.07199 (10)	0.0172 (2)
C4	0.61284 (11)	1.17695 (16)	-0.12363 (10)	0.0171 (2)
H4A	0.5993	1.2067	-0.1921	0.02*
C5	0.69644 (11)	1.06624 (16)	-0.07233 (10)	0.0165 (2)
H5A	0.7392	1.0211	-0.1069	0.02*
C6	0.71925 (10)	1.01896 (15)	0.03210 (9)	0.0150 (2)
C7	0.96469 (10)	0.78013 (15)	0.09632 (9)	0.0151 (2)

## supplementary materials

---

H7A	0.9782	0.7678	0.1668	0.018*
C8	1.04195 (10)	0.70636 (15)	0.05474 (9)	0.0142 (2)
C9	1.03109 (10)	0.72528 (15)	-0.05001 (9)	0.0151 (2)
C10	1.10725 (11)	0.64726 (16)	-0.08556 (10)	0.0168 (2)
H10A	1.1009	0.6634	-0.154	0.02*
C11	1.19152 (10)	0.54678 (16)	-0.02006 (10)	0.0169 (2)
H11A	1.2412	0.4933	-0.044	0.02*
C12	1.20051 (10)	0.52732 (15)	0.08265 (10)	0.0159 (2)
C13	1.12875 (10)	0.60635 (15)	0.12052 (10)	0.0153 (2)
H13A	1.1382	0.593	0.19	0.018*
H1N1	0.8262 (16)	0.891 (2)	0.1508 (16)	0.038 (5)*
H1O5	0.905 (2)	0.854 (3)	-0.082 (2)	0.067 (8)*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0238 (5)	0.0253 (5)	0.0220 (5)	0.0056 (4)	0.0118 (4)	0.0049 (4)
O2	0.0377 (6)	0.0307 (6)	0.0253 (5)	0.0079 (5)	0.0222 (5)	-0.0004 (4)
O3	0.0317 (6)	0.0461 (7)	0.0298 (6)	0.0207 (5)	0.0181 (5)	0.0077 (5)
O4	0.0379 (6)	0.0342 (6)	0.0228 (5)	0.0153 (5)	0.0150 (5)	0.0087 (4)
O5	0.0201 (4)	0.0210 (5)	0.0162 (4)	0.0047 (4)	0.0076 (3)	0.0023 (3)
O6	0.0284 (5)	0.0335 (6)	0.0285 (5)	0.0145 (5)	0.0114 (4)	0.0014 (4)
O7	0.0269 (5)	0.0307 (6)	0.0197 (5)	0.0070 (4)	0.0086 (4)	0.0059 (4)
N1	0.0163 (5)	0.0196 (5)	0.0155 (5)	0.0030 (4)	0.0085 (4)	0.0005 (4)
N2	0.0158 (5)	0.0157 (5)	0.0177 (5)	0.0000 (4)	0.0089 (4)	-0.0017 (4)
N3	0.0193 (5)	0.0220 (6)	0.0183 (5)	-0.0015 (4)	0.0112 (4)	-0.0006 (4)
N4	0.0209 (5)	0.0224 (6)	0.0204 (5)	0.0049 (4)	0.0097 (4)	0.0004 (4)
N5	0.0170 (5)	0.0177 (5)	0.0203 (5)	0.0010 (4)	0.0070 (4)	0.0005 (4)
C1	0.0177 (5)	0.0183 (6)	0.0144 (5)	-0.0009 (4)	0.0088 (4)	-0.0011 (4)
C2	0.0176 (5)	0.0191 (6)	0.0195 (6)	0.0006 (5)	0.0099 (5)	-0.0033 (5)
C3	0.0161 (5)	0.0174 (6)	0.0194 (6)	0.0029 (5)	0.0081 (5)	0.0005 (4)
C4	0.0194 (6)	0.0174 (6)	0.0158 (5)	-0.0005 (5)	0.0085 (4)	-0.0005 (4)
C5	0.0169 (5)	0.0185 (6)	0.0160 (5)	0.0001 (5)	0.0085 (4)	-0.0023 (4)
C6	0.0147 (5)	0.0150 (6)	0.0163 (5)	-0.0019 (4)	0.0070 (4)	-0.0017 (4)
C7	0.0161 (5)	0.0151 (6)	0.0158 (5)	-0.0009 (4)	0.0081 (4)	-0.0004 (4)
C8	0.0151 (5)	0.0143 (5)	0.0142 (5)	-0.0009 (4)	0.0068 (4)	-0.0013 (4)
C9	0.0156 (5)	0.0142 (6)	0.0156 (5)	-0.0007 (4)	0.0062 (4)	-0.0002 (4)
C10	0.0190 (6)	0.0181 (6)	0.0156 (5)	-0.0011 (5)	0.0093 (4)	-0.0010 (4)
C11	0.0159 (5)	0.0174 (6)	0.0200 (6)	-0.0012 (4)	0.0096 (5)	-0.0024 (4)
C12	0.0134 (5)	0.0140 (6)	0.0190 (6)	0.0010 (4)	0.0048 (4)	0.0005 (4)
C13	0.0160 (5)	0.0147 (6)	0.0161 (5)	-0.0009 (4)	0.0072 (4)	0.0000 (4)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—N3	1.2372 (15)	C2—H2A	0.93
O2—N3	1.2291 (15)	C3—C4	1.4035 (18)
O3—N4	1.2262 (15)	C4—C5	1.3711 (18)
O4—N4	1.2259 (15)	C4—H4A	0.93
O5—C9	1.3446 (15)	C5—C6	1.4159 (17)

O5—H1O5	0.92 (3)	C5—H5A	0.93
O6—N5	1.2302 (15)	C7—C8	1.4514 (17)
O7—N5	1.2288 (14)	C7—H7A	0.93
N1—C6	1.3565 (16)	C8—C13	1.3962 (17)
N1—N2	1.3697 (15)	C8—C9	1.4128 (16)
N1—H1N1	0.89 (2)	C9—C10	1.4014 (18)
N2—C7	1.2920 (16)	C10—C11	1.3799 (18)
N3—C1	1.4541 (16)	C10—H10A	0.93
N4—C3	1.4530 (17)	C11—C12	1.3927 (17)
N5—C12	1.4589 (16)	C11—H11A	0.93
C1—C2	1.3876 (18)	C12—C13	1.3795 (17)
C1—C6	1.4187 (17)	C13—H13A	0.93
C2—C3	1.3753 (17)		
C9—O5—H1O5	105.4 (15)	C4—C5—H5A	119.2
C6—N1—N2	120.62 (11)	C6—C5—H5A	119.2
C6—N1—H1N1	122.4 (13)	N1—C6—C5	120.57 (11)
N2—N1—H1N1	116.6 (13)	N1—C6—C1	122.75 (11)
C7—N2—N1	115.45 (10)	C5—C6—C1	116.66 (11)
O2—N3—O1	122.71 (11)	N2—C7—C8	120.94 (11)
O2—N3—C1	118.60 (11)	N2—C7—H7A	119.5
O1—N3—C1	118.64 (11)	C8—C7—H7A	119.5
O4—N4—O3	123.45 (12)	C13—C8—C9	118.32 (11)
O4—N4—C3	118.12 (11)	C13—C8—C7	118.27 (11)
O3—N4—C3	118.43 (11)	C9—C8—C7	123.37 (11)
O7—N5—O6	123.08 (11)	O5—C9—C10	117.82 (11)
O7—N5—C12	118.31 (11)	O5—C9—C8	121.85 (11)
O6—N5—C12	118.60 (11)	C10—C9—C8	120.32 (11)
C2—C1—C6	122.19 (11)	C11—C10—C9	120.67 (11)
C2—C1—N3	116.05 (11)	C11—C10—H10A	119.7
C6—C1—N3	121.76 (11)	C9—C10—H10A	119.7
C3—C2—C1	118.67 (12)	C10—C11—C12	118.49 (12)
C3—C2—H2A	120.7	C10—C11—H11A	120.8
C1—C2—H2A	120.7	C12—C11—H11A	120.8
C2—C3—C4	121.38 (12)	C13—C12—C11	122.02 (11)
C2—C3—N4	119.01 (11)	C13—C12—N5	118.45 (11)
C4—C3—N4	119.61 (11)	C11—C12—N5	119.53 (11)
C5—C4—C3	119.51 (12)	C12—C13—C8	120.14 (11)
C5—C4—H4A	120.2	C12—C13—H13A	119.9
C3—C4—H4A	120.2	C8—C13—H13A	119.9
C4—C5—C6	121.55 (12)		
C6—N1—N2—C7	172.84 (11)	C2—C1—C6—C5	-2.48 (18)
O2—N3—C1—C2	-15.33 (17)	N3—C1—C6—C5	176.74 (11)
O1—N3—C1—C2	162.33 (11)	N1—N2—C7—C8	175.64 (11)
O2—N3—C1—C6	165.40 (12)	N2—C7—C8—C13	-174.24 (11)
O1—N3—C1—C6	-16.94 (18)	N2—C7—C8—C9	3.12 (19)
C6—C1—C2—C3	1.35 (19)	C13—C8—C9—O5	177.95 (11)
N3—C1—C2—C3	-177.92 (11)	C7—C8—C9—O5	0.58 (19)
C1—C2—C3—C4	0.5 (2)	C13—C8—C9—C10	-1.09 (18)



## supplementary materials

---

C1—C2—C3—N4	-179.46 (11)	C7—C8—C9—C10	-178.46 (11)
O4—N4—C3—C2	174.55 (13)	O5—C9—C10—C11	-176.95 (11)
O3—N4—C3—C2	-5.12 (19)	C8—C9—C10—C11	2.12 (19)
O4—N4—C3—C4	-5.37 (19)	C9—C10—C11—C12	-1.25 (19)
O3—N4—C3—C4	174.96 (13)	C10—C11—C12—C13	-0.63 (19)
C2—C3—C4—C5	-1.0 (2)	C10—C11—C12—N5	179.16 (11)
N4—C3—C4—C5	178.90 (11)	O7—N5—C12—C13	-4.77 (17)
C3—C4—C5—C6	-0.22 (19)	O6—N5—C12—C13	174.17 (12)
N2—N1—C6—C5	2.89 (18)	O7—N5—C12—C11	175.44 (12)
N2—N1—C6—C1	-175.04 (11)	O6—N5—C12—C11	-5.62 (18)
C4—C5—C6—N1	-176.16 (12)	C11—C12—C13—C8	1.65 (19)
C4—C5—C6—C1	1.89 (18)	N5—C12—C13—C8	-178.14 (11)
C2—C1—C6—N1	175.52 (12)	C9—C8—C13—C12	-0.75 (18)
N3—C1—C6—N1	-5.25 (19)	C7—C8—C13—C12	176.75 (11)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N1 $\cdots$ O4 <sup>i</sup>	0.89 (2)	2.54 (2)	3.0666 (15)	118 (2)
N1—H1N1 $\cdots$ O1	0.89 (2)	2.07 (2)	2.6477 (15)	121 (2)
O5—H1O5 $\cdots$ N2	0.92 (3)	1.82 (3)	2.6656 (14)	150 (2)

Symmetry codes: (i)  $x+1/2, -y+5/2, z+1/2$ .

Fig. 1




Fig. 2

