

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-[(2-Hydroxy-1-naphthyl)methylidene-amino]benzoic acid

Mehmet Akkurt,^{a*} Sema Öztürk Yıldırım,^a
Abdullah Mohamed Asiri^b and Vickie McKee^c

^aDepartment of Physics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey, ^bChemistry Department, Faculty of Science, King Abdul-Aziz University, P. O. Box 80203, Jeddah 21589, Saudi Arabia, and ^cDepartment of Chemistry, Loughborough University, Leicestershire LE11 3TU, England
Correspondence e-mail: akkurt@erciyes.edu.tr


Received 3 March 2008; accepted 5 March 2008

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.051; wR factor = 0.145; data-to-parameter ratio = 16.8.

The molecule of the title compound, $\text{C}_{18}\text{H}_{13}\text{NO}_3$, is almost planar, the dihedral angle between the naphthalene and benzene ring systems being $4.04(6)^\circ$. The molecular conformation and packing are stabilized by intramolecular $\text{O}-\text{H}\cdots\text{N}$ and intermolecular $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ interactions.

Related literature

For background, see: Asiri & Badahdah (2007).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{13}\text{NO}_3$
 $M_r = 291.29$

Monoclinic, $C2/c$
 $a = 14.7490(12)$ Å

$b = 4.9850(4)$ Å
 $c = 36.750(3)$ Å
 $\beta = 91.305(1)^\circ$
 $V = 2701.3(4)$ Å³
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 150(2)$ K
 $0.31 \times 0.19 \times 0.09$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.970$, $T_{\max} = 0.991$

13445 measured reflections
3520 independent reflections
2761 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.145$
 $S = 1.07$
3520 reflections

210 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.59$ e Å⁻³
 $\Delta\rho_{\min} = -0.33$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H}\text{O1}\cdots\text{N1}$	0.82	1.82	2.5572 (16)	148
$\text{O2}-\text{H}\text{O2}\cdots\text{O3}^{\text{i}}$	0.82	1.81	2.6281 (14)	171
$\text{C14}-\text{H14}\cdots\text{O2}^{\text{ii}}$	0.93	2.56	3.3611 (17)	145
$\text{C16}-\text{H16}\cdots\text{O1}^{\text{iii}}$	0.93	2.49	3.1542 (19)	128

Symmetry codes: (i) $-x + 1, -y - 1, -z$; (ii) $-x + \frac{1}{2}, -y - \frac{1}{2}, -z$; (iii) $x + \frac{1}{2}, y - \frac{1}{2}, z$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2705).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
Asiri, A. M. & Badahdah, K. O. (2007). *Molecules*, **12**, 1796–1804.
Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o682 [doi:10.1107/S1600536808006107]

4-[(2-Hydroxy-1-naphthyl)methylideneamino]benzoic acid

M. Akkurt, S. Ö. Yildirim, A. M. Asiri and V. McKee

Comment

2-Hydroxy Schiff base ligands and their complexes, derived from the reaction of salicylaldehyde and 2-hydroxy-1-naphthaldehyde with amines are of interest due to the existence of (O—H \cdots N and N—H \cdots O) type hydrogen bonds and tautomerism between the enol-imine and keto-enamine forms. Tautomerism in 2-hydroxy Schiff bases both in solution and in the solid state was investigated using different spectroscopic techniques (Asiri & Badahdah, 2007).

In the title compound, (I), the molecule is almost planar (Fig. 1). The maximum deviation of the non-H atoms from their mean plane is 0.087 (1) Å for O3. The dihedral angle between the naphthalene ring and the benzene ring is 4.04 (6)°.

The molecular conformation is stabilized by an intramolecular O—H \cdots N hydrogen bond (Table 1). Then, classical inversion dimers are formed by head-to-head O—H \cdots O linkages of the carboxylic acid groups. Finally, C—H \cdots O interactions link the dimers into sheets (Fig. 2).

Experimental

A solution of 4-aminobenzoic acid (5.0 g, 36.5 mmol) in hot ethanol was mixed with an ethanolic solution of 2-hydroxynaphthaldehyde (7.23 g, 36.5 mmol) and the resulting mixture was refluxed for 3 h. The mixture was cooled to recover the crude product. Orange laths of (I) were recrystallized from ethanol. IR ν (cm⁻¹); 1683.3 (C=O), 1588.1 (C=N), 1427.1 (C=C), 1301.4 (C—O) and 1152 (C—N). [*M.p.*: > 573 K, yield: 54.7%].

Refinement

The H atoms were positioned geometrically (C—H = 0.93 Å, O—H = 0.82 Å) and refined as riding with with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of (I), with displacement ellipsoids for the non-H atoms are drawn at the 50% probability level.




Fig. 2. The packing for (I) showing hydrogen bonds as dashed lines.

4-[(2-Hydroxy-1-naphthyl)methylideneamino]benzoic acid

Crystal data

$C_{18}H_{13}NO_3$

$M_r = 291.29$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 14.7490\ (12)\ \text{\AA}$

$b = 4.9850\ (4)\ \text{\AA}$

$c = 36.750\ (3)\ \text{\AA}$

$\beta = 91.305\ (1)^\circ$

$V = 2701.3\ (4)\ \text{\AA}^3$

$Z = 8$

$F_{000} = 1216$

$D_x = 1.433\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71069\ \text{\AA}$

Cell parameters from 3973 reflections

$\theta = 2.2\text{--}28.8^\circ$

$\mu = 0.10\ \text{mm}^{-1}$

$T = 150\ (2)\ \text{K}$

Lath, orange

$0.31 \times 0.19 \times 0.09\ \text{mm}$

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: sealed tube

Monochromator: graphite

$T = 150\ (2)\ \text{K}$

ϕ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2005)

$T_{\min} = 0.970$, $T_{\max} = 0.991$

13445 measured reflections

3520 independent reflections

2761 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 28.9^\circ$

$\theta_{\min} = 2.2^\circ$

$h = -20 \rightarrow 19$

$k = -6 \rightarrow 6$

$l = -49 \rightarrow 49$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.050$	H-atom parameters constrained
$wR(F^2) = 0.145$	$w = 1/[\sigma^2(F_o^2) + (0.0792P)^2 + 1.0642P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
3520 reflections	$(\Delta/\sigma)_{\max} < 0.001$
210 parameters	$\Delta\rho_{\max} = 0.59 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > 2\sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.11588 (7)	0.8430 (3)	0.11623 (4)	0.0398 (4)
O2	0.39188 (7)	-0.3435 (2)	0.00066 (3)	0.0295 (3)
O3	0.52323 (6)	-0.2551 (2)	0.03002 (3)	0.0277 (3)
N1	0.26607 (8)	0.6069 (2)	0.10400 (3)	0.0232 (3)
C1	0.15748 (10)	0.9891 (3)	0.13926 (4)	0.0293 (4)
C2	0.10809 (11)	1.1871 (4)	0.15946 (5)	0.0387 (5)
C3	0.14890 (11)	1.3471 (3)	0.18415 (5)	0.0368 (5)
C4	0.24458 (10)	1.3336 (3)	0.19194 (4)	0.0273 (4)
C5	0.28528 (12)	1.5100 (3)	0.21737 (4)	0.0333 (4)
C6	0.37653 (13)	1.5035 (3)	0.22449 (4)	0.0367 (5)
C7	0.43000 (12)	1.3207 (3)	0.20591 (5)	0.0382 (5)
C8	0.39165 (11)	1.1460 (3)	0.18077 (4)	0.0328 (4)
C9	0.29754 (10)	1.1446 (3)	0.17322 (4)	0.0235 (4)
C10	0.25387 (9)	0.9658 (3)	0.14694 (4)	0.0227 (3)
C11	0.30277 (9)	0.7702 (3)	0.12869 (4)	0.0222 (3)
C12	0.31172 (9)	0.4082 (3)	0.08444 (4)	0.0216 (3)
C13	0.26106 (9)	0.2595 (3)	0.05899 (4)	0.0237 (4)
C14	0.30229 (9)	0.0589 (3)	0.03908 (4)	0.0236 (4)

supplementary materials

C15	0.39461 (9)	0.0041 (3)	0.04452 (4)	0.0212 (3)
C16	0.44479 (9)	0.1537 (3)	0.07000 (4)	0.0237 (4)
C17	0.40437 (9)	0.3544 (3)	0.08974 (4)	0.0237 (4)
C18	0.44065 (9)	-0.2106 (3)	0.02406 (4)	0.0224 (3)
HO1	0.15250	0.74990	0.10550	0.0600*
H2	0.04600	1.20430	0.15520	0.053 (6)*
HO2	0.42310	-0.45920	-0.00880	0.0440*
H3	0.11410	1.47080	0.19660	0.046 (5)*
H5	0.24950	1.63300	0.22950	0.041 (5)*
H6	0.40270	1.61950	0.24150	0.045 (5)*
H7	0.49230	1.31630	0.21050	0.058 (6)*
H8	0.42870	1.02670	0.16860	0.037 (5)*
H11	0.36440	0.75340	0.13410	0.0270*
H13	0.19960	0.29500	0.05540	0.0280*
H14	0.26850	-0.03930	0.02210	0.031 (4)*
H16	0.50620	0.11760	0.07370	0.022 (4)*
H17	0.43850	0.45380	0.10650	0.031 (4)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0220 (5)	0.0457 (7)	0.0513 (7)	0.0052 (5)	-0.0055 (5)	-0.0177 (6)
O2	0.0250 (5)	0.0288 (5)	0.0348 (6)	-0.0002 (4)	0.0003 (4)	-0.0125 (4)
O3	0.0212 (5)	0.0280 (5)	0.0340 (5)	0.0019 (4)	0.0017 (4)	-0.0052 (4)
N1	0.0220 (5)	0.0239 (6)	0.0238 (6)	0.0009 (4)	0.0003 (4)	-0.0034 (4)
C1	0.0236 (7)	0.0307 (7)	0.0336 (8)	0.0035 (6)	0.0012 (6)	-0.0026 (6)
C2	0.0244 (7)	0.0432 (9)	0.0485 (10)	0.0085 (7)	0.0031 (7)	-0.0090 (8)
C3	0.0340 (8)	0.0371 (8)	0.0396 (9)	0.0111 (7)	0.0078 (7)	-0.0078 (7)
C4	0.0359 (8)	0.0237 (7)	0.0224 (7)	0.0030 (6)	0.0047 (5)	0.0017 (5)
C5	0.0487 (9)	0.0257 (7)	0.0257 (7)	0.0055 (7)	0.0049 (6)	-0.0032 (6)
C6	0.0533 (10)	0.0281 (7)	0.0285 (8)	-0.0041 (7)	-0.0049 (7)	-0.0053 (6)
C7	0.0379 (9)	0.0361 (8)	0.0402 (9)	-0.0015 (7)	-0.0091 (7)	-0.0079 (7)
C8	0.0319 (8)	0.0295 (7)	0.0368 (8)	0.0039 (6)	-0.0038 (6)	-0.0084 (6)
C9	0.0287 (7)	0.0204 (6)	0.0213 (6)	0.0014 (5)	0.0020 (5)	0.0011 (5)
C10	0.0233 (6)	0.0226 (6)	0.0222 (6)	0.0009 (5)	0.0012 (5)	0.0000 (5)
C11	0.0209 (6)	0.0230 (6)	0.0228 (6)	0.0005 (5)	0.0000 (5)	0.0001 (5)
C12	0.0219 (6)	0.0215 (6)	0.0214 (6)	0.0000 (5)	0.0016 (5)	-0.0002 (5)
C13	0.0182 (6)	0.0264 (7)	0.0264 (7)	0.0009 (5)	-0.0023 (5)	-0.0018 (5)
C14	0.0223 (6)	0.0246 (6)	0.0237 (7)	-0.0021 (5)	-0.0022 (5)	-0.0031 (5)
C15	0.0206 (6)	0.0210 (6)	0.0221 (6)	-0.0017 (5)	0.0018 (5)	-0.0010 (5)
C16	0.0188 (6)	0.0261 (7)	0.0263 (7)	-0.0006 (5)	-0.0001 (5)	-0.0033 (5)
C17	0.0210 (6)	0.0258 (7)	0.0243 (6)	-0.0020 (5)	-0.0016 (5)	-0.0048 (5)
C18	0.0223 (6)	0.0214 (6)	0.0235 (6)	-0.0025 (5)	0.0027 (5)	-0.0013 (5)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.265 (2)	C12—C17	1.4018 (19)
O2—C18	1.2914 (18)	C12—C13	1.397 (2)
O3—C18	1.2524 (16)	C13—C14	1.388 (2)

O1—HO1	0.8200	C14—C15	1.3987 (19)
O2—HO2	0.8200	C15—C16	1.396 (2)
N1—C12	1.4047 (18)	C15—C18	1.482 (2)
N1—C11	1.3254 (18)	C16—C17	1.379 (2)
C1—C2	1.442 (2)	C2—H2	0.9300
C1—C10	1.448 (2)	C3—H3	0.9300
C2—C3	1.341 (3)	C5—H5	0.9300
C3—C4	1.435 (2)	C6—H6	0.9300
C4—C9	1.413 (2)	C7—H7	0.9300
C4—C5	1.408 (2)	C8—H8	0.9300
C5—C6	1.366 (3)	C11—H11	0.9300
C6—C7	1.394 (2)	C13—H13	0.9300
C7—C8	1.381 (2)	C14—H14	0.9300
C8—C9	1.409 (2)	C16—H16	0.9300
C9—C10	1.454 (2)	C17—H17	0.9300
C10—C11	1.394 (2)		
C1—O1—HO1	109.00	C14—C15—C18	121.63 (13)
C18—O2—HO2	109.00	C15—C16—C17	120.76 (13)
C11—N1—C12	126.19 (12)	C12—C17—C16	119.83 (13)
O1—C1—C2	119.70 (14)	O3—C18—C15	119.63 (13)
C2—C1—C10	117.37 (13)	O2—C18—O3	123.39 (13)
O1—C1—C10	122.92 (14)	O2—C18—C15	116.99 (12)
C1—C2—C3	122.09 (15)	C1—C2—H2	119.00
C2—C3—C4	122.11 (15)	C3—C2—H2	119.00
C3—C4—C9	119.09 (14)	C2—C3—H3	119.00
C3—C4—C5	120.38 (14)	C4—C3—H3	119.00
C5—C4—C9	120.52 (14)	C4—C5—H5	119.00
C4—C5—C6	121.06 (14)	C6—C5—H5	119.00
C5—C6—C7	119.19 (15)	C5—C6—H6	120.00
C6—C7—C8	120.81 (16)	C7—C6—H6	120.00
C7—C8—C9	121.42 (14)	C6—C7—H7	120.00
C4—C9—C10	119.40 (13)	C8—C7—H7	120.00
C4—C9—C8	116.99 (13)	C7—C8—H8	119.00
C8—C9—C10	123.59 (13)	C9—C8—H8	119.00
C1—C10—C11	118.61 (13)	N1—C11—H11	118.00
C1—C10—C9	119.90 (13)	C10—C11—H11	118.00
C9—C10—C11	121.48 (12)	C12—C13—H13	120.00
N1—C11—C10	123.46 (12)	C14—C13—H13	120.00
N1—C12—C17	122.74 (13)	C13—C14—H14	120.00
C13—C12—C17	119.77 (13)	C15—C14—H14	120.00
N1—C12—C13	117.49 (12)	C15—C16—H16	120.00
C12—C13—C14	120.09 (12)	C17—C16—H16	120.00
C13—C14—C15	120.15 (13)	C12—C17—H17	120.00
C14—C15—C16	119.41 (13)	C16—C17—H17	120.00
C16—C15—C18	118.96 (12)		
C12—N1—C11—C10	179.29 (14)	C7—C8—C9—C10	179.74 (15)
C11—N1—C12—C13	-178.82 (14)	C4—C9—C10—C1	2.2 (2)
C11—N1—C12—C17	1.6 (2)	C4—C9—C10—C11	-177.85 (14)

supplementary materials

O1—C1—C2—C3	-179.95 (18)	C8—C9—C10—C1	-176.23 (14)
C10—C1—C2—C3	0.8 (3)	C8—C9—C10—C11	3.8 (2)
O1—C1—C10—C9	178.67 (15)	C1—C10—C11—N1	1.4 (2)
O1—C1—C10—C11	-1.3 (2)	C9—C10—C11—N1	-178.55 (14)
C2—C1—C10—C9	-2.1 (2)	N1—C12—C13—C14	-179.43 (13)
C2—C1—C10—C11	177.93 (15)	C17—C12—C13—C14	0.1 (2)
C1—C2—C3—C4	0.5 (3)	N1—C12—C17—C16	179.04 (13)
C2—C3—C4—C5	178.31 (16)	C13—C12—C17—C16	-0.5 (2)
C2—C3—C4—C9	-0.5 (2)	C12—C13—C14—C15	0.3 (2)
C3—C4—C5—C6	-178.57 (15)	C13—C14—C15—C16	-0.3 (2)
C9—C4—C5—C6	0.2 (2)	C13—C14—C15—C18	179.30 (14)
C3—C4—C9—C8	177.61 (14)	C14—C15—C16—C17	-0.1 (2)
C3—C4—C9—C10	-0.9 (2)	C18—C15—C16—C17	-179.67 (14)
C5—C4—C9—C8	-1.1 (2)	C14—C15—C18—O2	0.7 (2)
C5—C4—C9—C10	-179.65 (14)	C14—C15—C18—O3	-179.25 (14)
C4—C5—C6—C7	0.7 (2)	C16—C15—C18—O2	-179.75 (13)
C5—C6—C7—C8	-0.5 (2)	C16—C15—C18—O3	0.3 (2)
C6—C7—C8—C9	-0.5 (2)	C15—C16—C17—C12	0.5 (2)
C7—C8—C9—C4	1.3 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—HO1 \cdots N1	0.82	1.82	2.5572 (16)	148
O2—HO2 \cdots O3 ⁱ	0.82	1.81	2.6281 (14)	171
C14—H14 \cdots O2 ⁱⁱ	0.93	2.56	3.3611 (17)	145
C16—H16 \cdots O1 ⁱⁱⁱ	0.93	2.49	3.1542 (19)	128

Symmetry codes: (i) $-x+1, -y-1, -z$; (ii) $-x+1/2, -y-1/2, -z$; (iii) $x+1/2, y-1/2, z$.

Fig. 1




Fig. 2

