

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

6-Amino-3-methyl-4-(4-nitrophenyl)-1-phenylpyrazolo[3,4-*b*]pyridine-5-carbonitrile

Xin-Ying Zhang,* Xiao-Yan Li, Xia Wang, Xue-Sen Fan* and Gui-Rong Qu

School of Chemical and Environmental Sciences, Henan Key Laboratory for Environmental Pollution Control, Henan Normal University, Xinxiang, Henan 453007, People's Republic of China

Correspondence e-mail: xyzh518@sohu.com, xuesen.fan@yahoo.com


Received 2 March 2008; accepted 5 March 2008

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.042; wR factor = 0.122; data-to-parameter ratio = 13.3.

The title compound, $\text{C}_{20}\text{H}_{14}\text{N}_6\text{O}_2$, contains four rings. The dihedral angle between the pyridine ring and the pyrazole ring is $1.9(1)^\circ$, *i.e.* almost coplanar, which gives rise to a conjugated structure. The dihedral angle between the nitro-substituted phenyl ring and the pyridine ring is $76.3(1)^\circ$ and that between the pyrazole ring and the non-substituted phenyl ring is $40.5(1)^\circ$. In the crystal structure, symmetry-related molecules are linked by $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds.

Related literature

For related structures, see: Quiroga *et al.* (1999); Zhu *et al.* (2005). For the biological and pharmacological activities, see: Kamal *et al.* (1991); Straub *et al.* (2001); Sekikawa *et al.* (1973).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{14}\text{N}_6\text{O}_2$
 $M_r = 370.37$
 Monoclinic, $C2/c$

$a = 16.470(11)$ Å
 $b = 9.742(7)$ Å
 $c = 23.46(2)$ Å

$\beta = 105.857(8)^\circ$
 $V = 3621(5)$ Å³
 $Z = 8$
 Mo $K\alpha$ radiation

$\mu = 0.09$ mm⁻¹
 $T = 294(2)$ K
 $0.39 \times 0.25 \times 0.15$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.945$, $T_{\max} = 0.986$
 13443 measured reflections
 3374 independent reflections
 2287 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.121$
 $S = 1.02$
 3374 reflections
 254 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.18$ e Å⁻³
 $\Delta\rho_{\min} = -0.16$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N5}-\text{H5A}\cdots\text{O2}^i$	0.86	2.13	2.981 (3)	168
$\text{C14}-\text{H14}\cdots\text{N2}^i$	0.93	2.61	3.529 (3)	168

Symmetry codes: (i) $x, -y, z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by the National Natural Science Foundation of China (No. 20772025) and the Program for Science & Technology Innovation Talents in Universities of Henan Province (No. 2008 HASTIT006).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2046).

References

- Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Kamal, A. M., Atalla, A. A., Mohamed, T. A. A. & Geies, A. (1991). *Z. Naturforsch. Teil B*, **46**, 541–544.
 Quiroga, J., Alvarado, M., Insuasty, B., Moreno, R., Ravina, E., Estevez, I. & de Almedia, R. H. (1999). *J. Heterocycl. Chem.* **36**, 1311–1316.
 Sekikawa, I., Nishie, J., Tono-oka, S., Tanaka, Y. & Kakimoto, S. (1973). *J. Heterocycl. Chem.* **10**, 931–932.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Straub, A., Stasch, J.-P., Alonso-Alija, C., Benet-Buchholz, J., Ducke, B., Feurer, A. & Furstner, C. (2001). *Bioorg. Med. Chem. Lett.* **11**, 781–784.
 Zhu, S.-L., Tu, S.-J., Li, T.-J., Zhang, X.-J., Ji, S.-J. & Zhang, Y. (2005). *Chin. J. Org. Chem.* **25**, 987–990.

supplementary materials

Acta Cryst. (2008). E64, o689 [doi:10.1107/S1600536808006156]

6-Amino-3-methyl-4-(4-nitrophenyl)-1-phenylpyrazolo[3,4-*b*]pyridine-5-carbonitrile

X.-Y. Zhang, X.-Y. Li, X. Wang, X.-S. Fan and G.-R. Qu

Comment

The structure of the title compound, (I), is shown below. Dimensions are available in the archived CIF.

For related literature, see Quiroga *et al.* (1999); Zhu *et al.* (2005). For the biological and pharmacological activities, see Kamal *et al.* (1991); Straub *et al.* (2001); Sekikawa *et al.* (1973).

Experimental

The title compound was prepared by the following procedure: To 1 ml of 1-butyl-3-methylimidazolium tetrafluoroborate ([bmim][BF₄]) were added 4-nitrobenzaldehyde (1 mmol), malononitrile (1 mmol) and 5-amino-3-methyl-1-phenylpyrazole (1 mmol). The reaction mixture was stirred at 80°C for 10 hrs. The yellow solid product that was obtained was collected by suction and rinsed with water and ethanol (yield 93%). Single crystals of the title compound were obtained by slow evaporation from ethanol.

Refinement

H-atoms were included in calculated positions and treated as riding atoms: N—H = 0.86 Å and C—H = 0.93 – 0.96 Å with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{CH}_3)$ and $1.2U_{\text{eq}}(\text{NH}_2, \text{CH})$.

Figures




Fig. 1. Molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level.

6-Amino-3-methyl-4-(4-nitrophenyl)-1-phenylpyrazolo[3,4-*b*]pyridine- 5-carbonitrile

Crystal data

C₂₀H₁₄N₆O₂

$M_r = 370.37$

Monoclinic, *C*2/*c*

Hall symbol: -*C* 2yc

$a = 16.470$ (11) Å

$b = 9.742$ (7) Å

$F_{000} = 1536$

$D_x = 1.359$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2825 reflections

$\theta = 2.5$ – 21.6°

$\mu = 0.09$ mm⁻¹

supplementary materials

$c = 23.46 (2) \text{ \AA}$
 $\beta = 105.857 (8)^\circ$
 $V = 3621 (5) \text{ \AA}^3$
 $Z = 8$

$T = 294 (2) \text{ K}$
Block, yellow
 $0.39 \times 0.25 \times 0.15 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	3374 independent reflections
Radiation source: fine-focus sealed tube	2287 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 294(2) \text{ K}$	$\theta_{\text{max}} = 25.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -19 \rightarrow 19$
$T_{\text{min}} = 0.945$, $T_{\text{max}} = 0.986$	$k = -11 \rightarrow 11$
13443 measured reflections	$l = -28 \rightarrow 28$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.122$	$w = 1/[\sigma^2(F_o^2) + (0.0546P)^2 + 1.3917P]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
3374 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
254 parameters	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.16 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes)

are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and

goodness of fit S are based on F^2 , conventional R -factors R are based

on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F, and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.14291 (11)	0.3249 (2)	0.22385 (8)	0.0539 (5)
C2	0.13161 (11)	0.17956 (19)	0.22657 (8)	0.0508 (4)
C3	0.12907 (11)	0.06720 (19)	0.18900 (8)	0.0514 (5)
C4	0.11731 (12)	-0.0608 (2)	0.21136 (8)	0.0579 (5)
C5	0.10829 (12)	-0.0751 (2)	0.27046 (8)	0.0586 (5)
C6	0.11882 (11)	0.1536 (2)	0.28254 (8)	0.0515 (5)
C7	0.16473 (14)	0.4076 (2)	0.17675 (9)	0.0675 (6)
H7A	0.1171	0.4109	0.1424	0.101*
H7B	0.2117	0.3663	0.1665	0.101*
H7C	0.1795	0.4992	0.1910	0.101*
C8	0.11094 (16)	-0.1792 (2)	0.17454 (10)	0.0771 (7)
C9	0.13676 (11)	0.08314 (19)	0.12762 (8)	0.0516 (5)
C10	0.06928 (13)	0.1307 (3)	0.08327 (9)	0.0750 (7)
H10	0.0187	0.1510	0.0919	0.090*
C11	0.07559 (14)	0.1485 (2)	0.02644 (9)	0.0749 (7)
H11	0.0300	0.1808	-0.0034	0.090*
C12	0.15017 (13)	0.11769 (19)	0.01483 (8)	0.0573 (5)
C13	0.21832 (13)	0.0685 (2)	0.05755 (9)	0.0656 (6)
H13	0.2684	0.0471	0.0484	0.079*
C14	0.21111 (12)	0.0516 (2)	0.11416 (8)	0.0617 (5)
H14	0.2568	0.0184	0.1437	0.074*
C15	0.11107 (11)	0.3128 (2)	0.36614 (8)	0.0539 (5)
C16	0.14851 (12)	0.2328 (2)	0.41484 (9)	0.0610 (5)
H16	0.1792	0.1550	0.4107	0.073*
C17	0.13988 (13)	0.2697 (2)	0.47001 (9)	0.0674 (6)
H17	0.1640	0.2155	0.5029	0.081*
C18	0.09585 (14)	0.3862 (3)	0.47639 (10)	0.0732 (7)
H18	0.0910	0.4111	0.5136	0.088*
C19	0.05918 (14)	0.4654 (2)	0.42781 (11)	0.0722 (6)
H19	0.0294	0.5440	0.4322	0.087*
C20	0.06610 (12)	0.4293 (2)	0.37226 (10)	0.0636 (5)
H20	0.0408	0.4828	0.3394	0.076*
N1	0.12015 (10)	0.27857 (17)	0.30914 (7)	0.0569 (4)
N2	0.13518 (10)	0.38354 (17)	0.27275 (7)	0.0592 (4)
N3	0.10738 (10)	0.03211 (17)	0.30563 (6)	0.0575 (4)

supplementary materials

N4	0.1045 (2)	-0.2721 (2)	0.14393 (10)	0.1197 (9)
N5	0.10113 (12)	-0.20117 (18)	0.29226 (8)	0.0784 (6)
H5A	0.0962	-0.2102	0.3276	0.094*
H5B	0.1015	-0.2726	0.2708	0.094*
N6	0.15858 (14)	0.14160 (19)	-0.04520 (8)	0.0750 (5)
O1	0.22406 (12)	0.1080 (2)	-0.05634 (7)	0.0962 (6)
O2	0.09960 (14)	0.1956 (2)	-0.08097 (7)	0.1125 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0489 (10)	0.0631 (12)	0.0496 (11)	0.0009 (9)	0.0132 (8)	0.0005 (9)
C2	0.0476 (10)	0.0611 (12)	0.0439 (10)	0.0024 (8)	0.0128 (8)	-0.0009 (9)
C3	0.0479 (10)	0.0626 (12)	0.0425 (10)	0.0093 (9)	0.0102 (8)	0.0001 (9)
C4	0.0672 (13)	0.0611 (12)	0.0430 (10)	0.0062 (9)	0.0111 (9)	-0.0021 (9)
C5	0.0655 (12)	0.0633 (13)	0.0455 (11)	-0.0049 (10)	0.0128 (9)	-0.0017 (9)
C6	0.0499 (11)	0.0610 (12)	0.0439 (10)	-0.0028 (9)	0.0135 (8)	-0.0063 (9)
C7	0.0721 (13)	0.0687 (13)	0.0632 (13)	-0.0008 (11)	0.0210 (10)	0.0070 (11)
C8	0.1150 (19)	0.0602 (14)	0.0533 (13)	0.0113 (13)	0.0184 (12)	0.0015 (11)
C9	0.0545 (11)	0.0572 (11)	0.0428 (10)	0.0114 (9)	0.0128 (8)	-0.0004 (8)
C10	0.0606 (13)	0.1155 (19)	0.0501 (11)	0.0348 (12)	0.0173 (10)	0.0073 (12)
C11	0.0726 (14)	0.1049 (18)	0.0449 (11)	0.0368 (13)	0.0120 (10)	0.0089 (11)
C12	0.0760 (13)	0.0552 (11)	0.0441 (10)	0.0137 (10)	0.0221 (9)	-0.0006 (9)
C13	0.0623 (12)	0.0805 (14)	0.0601 (13)	0.0194 (11)	0.0267 (10)	0.0038 (10)
C14	0.0560 (12)	0.0791 (14)	0.0490 (11)	0.0208 (10)	0.0130 (9)	0.0070 (10)
C15	0.0470 (10)	0.0664 (12)	0.0498 (11)	-0.0113 (9)	0.0157 (8)	-0.0135 (9)
C16	0.0535 (11)	0.0732 (13)	0.0557 (12)	-0.0040 (10)	0.0141 (9)	-0.0105 (10)
C17	0.0603 (12)	0.0911 (16)	0.0513 (11)	-0.0140 (11)	0.0161 (9)	-0.0092 (11)
C18	0.0684 (14)	0.0973 (18)	0.0617 (14)	-0.0247 (13)	0.0310 (11)	-0.0278 (13)
C19	0.0646 (13)	0.0787 (15)	0.0832 (17)	-0.0096 (11)	0.0370 (12)	-0.0239 (13)
C20	0.0565 (12)	0.0698 (13)	0.0672 (13)	-0.0063 (10)	0.0216 (10)	-0.0086 (11)
N1	0.0641 (10)	0.0628 (10)	0.0464 (9)	-0.0078 (8)	0.0193 (7)	-0.0086 (8)
N2	0.0627 (10)	0.0615 (10)	0.0540 (10)	-0.0048 (8)	0.0171 (8)	-0.0040 (8)
N3	0.0638 (10)	0.0638 (11)	0.0449 (9)	-0.0083 (8)	0.0151 (7)	-0.0043 (8)
N4	0.206 (3)	0.0672 (14)	0.0819 (15)	0.0173 (16)	0.0322 (16)	-0.0113 (13)
N5	0.1225 (16)	0.0619 (11)	0.0510 (10)	-0.0128 (10)	0.0238 (10)	-0.0003 (8)
N6	0.1076 (16)	0.0713 (12)	0.0516 (10)	0.0168 (11)	0.0311 (11)	0.0007 (9)
O1	0.1160 (14)	0.1135 (14)	0.0769 (11)	0.0135 (11)	0.0567 (11)	0.0013 (10)
O2	0.1504 (18)	0.1383 (16)	0.0511 (9)	0.0640 (14)	0.0312 (10)	0.0233 (10)

Geometric parameters (\AA , $^\circ$)

C1—N2	1.318 (3)	C12—C13	1.371 (3)
C1—C2	1.431 (3)	C12—N6	1.471 (3)
C1—C7	1.489 (3)	C13—C14	1.375 (3)
C2—C3	1.399 (3)	C13—H13	0.9300
C2—C6	1.408 (3)	C14—H14	0.9300
C3—C4	1.386 (3)	C15—C16	1.381 (3)
C3—C9	1.488 (3)	C15—C20	1.384 (3)

C4—C8	1.428 (3)	C15—N1	1.425 (2)
C4—C5	1.441 (3)	C16—C17	1.387 (3)
C5—N3	1.333 (2)	C16—H16	0.9300
C5—N5	1.348 (3)	C17—C18	1.376 (3)
C6—N3	1.336 (2)	C17—H17	0.9300
C6—N1	1.365 (2)	C18—C19	1.372 (3)
C7—H7A	0.9600	C18—H18	0.9300
C7—H7B	0.9600	C19—C20	1.384 (3)
C7—H7C	0.9600	C19—H19	0.9300
C8—N4	1.142 (3)	C20—H20	0.9300
C9—C10	1.378 (3)	N1—N2	1.397 (2)
C9—C14	1.380 (3)	N5—H5A	0.8600
C10—C11	1.376 (3)	N5—H5B	0.8600
C10—H10	0.9300	N6—O2	1.216 (2)
C11—C12	1.362 (3)	N6—O1	1.222 (2)
C11—H11	0.9300		
N2—C1—C2	110.21 (17)	C13—C12—N6	118.86 (19)
N2—C1—C7	120.72 (18)	C12—C13—C14	118.58 (18)
C2—C1—C7	129.00 (18)	C12—C13—H13	120.7
C3—C2—C6	117.42 (18)	C14—C13—H13	120.7
C3—C2—C1	136.70 (18)	C13—C14—C9	120.68 (17)
C6—C2—C1	105.87 (16)	C13—C14—H14	119.7
C4—C3—C2	116.66 (17)	C9—C14—H14	119.7
C4—C3—C9	121.18 (17)	C16—C15—C20	120.50 (19)
C2—C3—C9	122.16 (17)	C16—C15—N1	120.44 (18)
C3—C4—C8	119.51 (18)	C20—C15—N1	119.04 (18)
C3—C4—C5	120.85 (17)	C15—C16—C17	119.3 (2)
C8—C4—C5	119.60 (19)	C15—C16—H16	120.4
N3—C5—N5	117.52 (18)	C17—C16—H16	120.4
N3—C5—C4	122.75 (19)	C18—C17—C16	120.4 (2)
N5—C5—C4	119.73 (18)	C18—C17—H17	119.8
N3—C6—N1	126.23 (17)	C16—C17—H17	119.8
N3—C6—C2	127.62 (17)	C19—C18—C17	119.9 (2)
N1—C6—C2	106.15 (17)	C19—C18—H18	120.0
C1—C7—H7A	109.5	C17—C18—H18	120.0
C1—C7—H7B	109.5	C18—C19—C20	120.5 (2)
H7A—C7—H7B	109.5	C18—C19—H19	119.7
C1—C7—H7C	109.5	C20—C19—H19	119.7
H7A—C7—H7C	109.5	C19—C20—C15	119.4 (2)
H7B—C7—H7C	109.5	C19—C20—H20	120.3
N4—C8—C4	178.2 (3)	C15—C20—H20	120.3
C10—C9—C14	119.01 (18)	C6—N1—N2	110.91 (15)
C10—C9—C3	120.08 (17)	C6—N1—C15	130.13 (16)
C14—C9—C3	120.91 (16)	N2—N1—C15	118.94 (16)
C11—C10—C9	120.98 (19)	C1—N2—N1	106.83 (16)
C11—C10—H10	119.5	C5—N3—C6	114.61 (17)
C9—C10—H10	119.5	C5—N5—H5A	120.0
C12—C11—C10	118.46 (18)	C5—N5—H5B	120.0
C12—C11—H11	120.8	H5A—N5—H5B	120.0

supplementary materials

C10—C11—H11	120.8	O2—N6—O1	123.6 (2)
C11—C12—C13	122.28 (18)	O2—N6—C12	117.6 (2)
C11—C12—N6	118.84 (18)	O1—N6—C12	118.81 (19)
N2—C1—C2—C3	176.8 (2)	N6—C12—C13—C14	177.45 (19)
C7—C1—C2—C3	-6.2 (4)	C12—C13—C14—C9	0.1 (3)
N2—C1—C2—C6	-1.8 (2)	C10—C9—C14—C13	0.7 (3)
C7—C1—C2—C6	175.12 (18)	C3—C9—C14—C13	-178.96 (19)
C6—C2—C3—C4	-2.4 (2)	C20—C15—C16—C17	0.6 (3)
C1—C2—C3—C4	179.1 (2)	N1—C15—C16—C17	178.90 (17)
C6—C2—C3—C9	176.39 (16)	C15—C16—C17—C18	-1.2 (3)
C1—C2—C3—C9	-2.1 (3)	C16—C17—C18—C19	0.9 (3)
C2—C3—C4—C8	177.67 (19)	C17—C18—C19—C20	0.0 (3)
C9—C3—C4—C8	-1.1 (3)	C18—C19—C20—C15	-0.5 (3)
C2—C3—C4—C5	0.0 (3)	C16—C15—C20—C19	0.2 (3)
C9—C3—C4—C5	-178.80 (17)	N1—C15—C20—C19	-178.06 (17)
C3—C4—C5—N3	2.7 (3)	N3—C6—N1—N2	178.85 (17)
C8—C4—C5—N3	-174.99 (19)	C2—C6—N1—N2	-1.3 (2)
C3—C4—C5—N5	-176.71 (18)	N3—C6—N1—C15	0.7 (3)
C8—C4—C5—N5	5.6 (3)	C2—C6—N1—C15	-179.46 (17)
C3—C2—C6—N3	2.7 (3)	C16—C15—N1—C6	40.1 (3)
C1—C2—C6—N3	-178.33 (18)	C20—C15—N1—C6	-141.6 (2)
C3—C2—C6—N1	-177.13 (15)	C16—C15—N1—N2	-137.89 (18)
C1—C2—C6—N1	1.82 (19)	C20—C15—N1—N2	40.4 (2)
C3—C4—C8—N4	-45 (11)	C2—C1—N2—N1	1.0 (2)
C5—C4—C8—N4	132 (10)	C7—C1—N2—N1	-176.20 (16)
C4—C3—C9—C10	102.6 (2)	C6—N1—N2—C1	0.2 (2)
C2—C3—C9—C10	-76.2 (3)	C15—N1—N2—C1	178.58 (15)
C4—C3—C9—C14	-77.8 (3)	N5—C5—N3—C6	176.90 (17)
C2—C3—C9—C14	103.5 (2)	C4—C5—N3—C6	-2.5 (3)
C14—C9—C10—C11	-0.9 (3)	N1—C6—N3—C5	179.64 (18)
C3—C9—C10—C11	178.8 (2)	C2—C6—N3—C5	-0.2 (3)
C9—C10—C11—C12	0.2 (4)	C11—C12—N6—O2	4.2 (3)
C10—C11—C12—C13	0.7 (4)	C13—C12—N6—O2	-174.2 (2)
C10—C11—C12—N6	-177.6 (2)	C11—C12—N6—O1	-176.6 (2)
C11—C12—C13—C14	-0.8 (3)	C13—C12—N6—O1	5.1 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N5—H5A \cdots O2 ⁱ	0.86	2.13	2.981 (3)	168
C14—H14 \cdots N2 ⁱⁱ	0.93	2.61	3.529 (3)	168

Symmetry codes: (i) $x, -y, z+1/2$; (ii) $-x+1/2, y-1/2, -z+1/2$.

Fig. 1

