

2-Amino-N-(4-methylphenylsulfonyl)-N-phenylbenzenesulfonamide

Zu-Wei Song

College of Science, Qingdao Agricultural University, Qingdao 266109, People's Republic of China

Correspondence e-mail: songzuwei2008@yahoo.cn


Received 20 February 2008; accepted 18 March 2008

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.040; wR factor = 0.121; data-to-parameter ratio = 17.6.

In the title molecule, $\text{C}_{19}\text{H}_{18}\text{N}_2\text{O}_4\text{S}_2$, the phenyl ring makes dihedral angles of 33.99 (2) and 43.70 (3)° with the two methyl-substituted benzene rings. Intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into centrosymmetric dimers. The crystal packing exhibits weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the crystal structures of related compounds, see: Henschel *et al.* (1996). For details of the biological activities of sulfonamide-containing compounds, see: Kamoshita *et al.* (1987). For related literature, see: Allen *et al.* (1987); Zhang *et al.* (2007).



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{18}\text{N}_2\text{O}_4\text{S}_2$
 $M_r = 402.47$

 Monoclinic, $P2_1/c$
 $a = 14.6245$ (3) Å

 $b = 10.0454$ (2) Å
 $c = 13.4735$ (4) Å
 $\beta = 107.478$ (2)°
 $V = 1887.99$ (7) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.31$ mm⁻¹
 $T = 293$ (2) K
 $0.52 \times 0.32 \times 0.25$ mm

Data collection

 Rigaku R-Axis RAPID IP area-detector diffractometer
 Absorption correction: multi-scan (ABSCOR; Higashi 1995)
 $T_{\min} = 0.855$, $T_{\max} = 0.926$

 17437 measured reflections
 4312 independent reflections
 3531 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.121$
 $S = 1.07$
 4312 reflections

 245 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.40$ e Å⁻³
 $\Delta\rho_{\min} = -0.26$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2B}\cdots\text{O3}^{\text{i}}$	0.86	2.20	3.062 (2)	176
$\text{C19}-\text{H19A}\cdots\text{O1}^{\text{ii}}$	0.96	2.57	3.517 (3)	169
$\text{C19}-\text{H19C}\cdots\text{O4}^{\text{iii}}$	0.96	2.58	3.538 (3)	174
$\text{N2}-\text{H2C}\cdots\text{O2}$	0.86	2.23	2.893 (2)	133

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2388).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Henschel, D., Hiemisch, O., Blaschette, A. & Jones, P. G. (1996). *Z. Naturforsch. Teil B*, **51**, 1313–1315.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Kamoshita, K., Matsumoto, H. & Nagano, E. (1987). US Patent 4 670 046.
 Rigaku (2004). *RAPID-AUTO*. Version 3.0. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zhang, Z. B., Zhou, S. Y. & Nie, J. (2007). *J. Mol. Catal. A Chem.* **265**, 9–14.

supplementary materials

Acta Cryst. (2008). E64, o744 [doi:10.1107/S1600536808007447]

2-Amino-*N*-(4-methylphenylsulfonyl)-*N*-phenylbenzenesulfonamide

Z.-W. Song

Comment

Many compounds containing sulfonimide groups possess a broad spectrum of biological activities and can be used as herbicides (Kamoshita *et al.*, 1987). In addition, some compounds containing sulfonimide groups can be used as catalysts (Zhang *et al.*, 2007). Herein we report the crystal structure of the title compound, (I).

In (I) (Fig. 1), all bond lengths and angles are normal (Allen *et al.*, 1987) and in a good agreement with those reported previously (Henschel *et al.*, 1996). The phenyl ring (C7—C12) makes the dihedral angles of 33.99 (2) and 43.70 (3)°, respectively, with two benzene rings (C1—C6/N1 and C13—C19). The intermolecular N—H···O hydrogen bonds (Table 1) link the molecules into centrosymmetric dimers. The crystal packing exhibits also weak intermolecular C—H···O hydrogen bonds (Table 1).

Experimental

A solution of methylsulfonyl chloride (1 mmol) was dissolved in anhydrous CH₂Cl₂ (10 ml), and dropwise added over a period of 10 min to a solution of 2-amino-*N*-methyl-benzenesulfonamide (1 mmol) and DMAP₂ (3 mmol) in CH₂Cl₂ (10 ml) at 273 K. The mixture was stirred for 4 h at room temperature. The organic phase was washed with 2 N HCl twice, and dried over anhydrous Na₂SO₄. The solvent was removed and the residue was purified by flash chromatography (1:1 cyclohexane:dichloromethane) to give (I) as a white solid (294 mg, 73%). Single crystals suitable for X-ray measurements were obtained by recrystallization from ethanol and dichloromethane at room temperature.

Refinement

H atoms were positioned geometrically with C—H = 0.93 or 0.96 Å, N—H = 0.86 Å, and refined using a riding model, with $U_{iso}(H) = 1.2$ (1.5 for methyl groups) times $U_{eq}(C, N)$.

Figures




Fig. 1. The molecular structure of (I) showing the atomic numbering and 40% probability displacement ellipsoids.

2-Amino-*N*-(4-methylphenylsulfonyl)-*N*-phenylbenzenesulfonamide

Crystal data

$C_{19}H_{18}N_2O_4S_2$	$F(000) = 840$
$M_r = 402.47$	$D_x = 1.416 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 2652 reflections
$a = 14.6245 (3) \text{ \AA}$	$\theta = 2.6\text{--}25.6^\circ$
$b = 10.0454 (2) \text{ \AA}$	$\mu = 0.31 \text{ mm}^{-1}$
$c = 13.4735 (4) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 107.478 (2)^\circ$	Platelet, yellow
$V = 1887.99 (7) \text{ \AA}^3$	$0.52 \times 0.32 \times 0.25 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-Axis RAPID IP area-detector diffractometer	4312 independent reflections
Radiation source: Rotating Anode graphite	3531 reflections with $I > 2\sigma(I)$
ω oscillation scans	$R_{\text{int}} = 0.038$
Absorption correction: multi-scan (ABSCOR; Higashi 1995)	$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.5^\circ$
$T_{\text{min}} = 0.855$, $T_{\text{max}} = 0.927$	$h = -18 \rightarrow 18$
17437 measured reflections	$k = -13 \rightarrow 13$
	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.121$	H-atom parameters constrained
$S = 1.07$	$w = 1/[\sigma^2(F_o^2) + (0.0647P)^2 + 0.3731P]$
4312 reflections	where $P = (F_o^2 + 2F_c^2)/3$
245 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.40 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.18008 (3)	0.28081 (5)	0.24068 (4)	0.04085 (14)
S2	0.24948 (3)	0.51762 (5)	0.14469 (3)	0.04136 (14)
O1	0.24248 (10)	0.20876 (14)	0.32523 (12)	0.0550 (4)
O2	0.16312 (11)	0.23184 (16)	0.13781 (11)	0.0573 (4)
O3	0.18520 (10)	0.46901 (19)	0.05005 (10)	0.0595 (4)
O4	0.24789 (12)	0.65454 (14)	0.17142 (12)	0.0571 (4)
N1	0.22958 (11)	0.43445 (16)	0.24350 (11)	0.0386 (3)
N2	-0.00565 (13)	0.3985 (2)	0.09409 (13)	0.0651 (6)
H2B	-0.0542	0.4371	0.0515	0.078*
H2C	0.0415	0.3727	0.0731	0.078*
C1	-0.00348 (13)	0.3782 (2)	0.19378 (15)	0.0431 (4)
C2	-0.08053 (14)	0.4204 (2)	0.22702 (18)	0.0541 (5)
H2D	-0.1332	0.4599	0.1794	0.065*
C3	-0.08033 (17)	0.4052 (3)	0.3272 (2)	0.0667 (6)
H3B	-0.1326	0.4346	0.3469	0.080*
C4	-0.00295 (18)	0.3463 (3)	0.4010 (2)	0.0691 (7)
H4B	-0.0034	0.3370	0.4696	0.083*
C5	0.07352 (16)	0.3022 (2)	0.37162 (16)	0.0540 (5)
H5A	0.1251	0.2618	0.4202	0.065*
C6	0.07447 (12)	0.31772 (19)	0.26886 (14)	0.0399 (4)
C7	0.25885 (13)	0.49745 (18)	0.34475 (13)	0.0375 (4)
C8	0.19747 (14)	0.5849 (2)	0.37156 (16)	0.0459 (4)
H8A	0.1393	0.6076	0.3234	0.055*
C9	0.22318 (18)	0.6384 (2)	0.47042 (19)	0.0586 (6)
H9A	0.1821	0.6970	0.4894	0.070*
C10	0.3098 (2)	0.6048 (3)	0.54093 (18)	0.0650 (6)
H10A	0.3268	0.6406	0.6077	0.078*
C11	0.37175 (19)	0.5185 (3)	0.51344 (18)	0.0640 (6)
H11A	0.4302	0.4969	0.5615	0.077*
C12	0.34685 (14)	0.4643 (2)	0.41479 (15)	0.0497 (5)
H12A	0.3883	0.4066	0.3956	0.060*
C13	0.36661 (12)	0.47731 (19)	0.14622 (13)	0.0384 (4)
C14	0.44113 (15)	0.5618 (2)	0.19508 (16)	0.0514 (5)
H14A	0.4301	0.6370	0.2301	0.062*
C15	0.53243 (15)	0.5326 (2)	0.19103 (18)	0.0562 (5)
H15A	0.5829	0.5890	0.2239	0.067*
C16	0.55032 (14)	0.4216 (2)	0.13933 (16)	0.0494 (5)
C17	0.47460 (17)	0.3389 (2)	0.09216 (19)	0.0590 (6)

supplementary materials

H17A	0.4860	0.2635	0.0576	0.071*
C18	0.38235 (15)	0.3642 (2)	0.09434 (18)	0.0523 (5)
H18A	0.3322	0.3071	0.0620	0.063*
C19	0.64941 (17)	0.3926 (3)	0.1334 (2)	0.0718 (7)
H19A	0.6850	0.4742	0.1402	0.108*
H19B	0.6451	0.3521	0.0676	0.108*
H19C	0.6815	0.3331	0.1887	0.108*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0370 (2)	0.0369 (2)	0.0459 (3)	0.00456 (18)	0.00835 (18)	-0.00020 (18)
S2	0.0365 (2)	0.0506 (3)	0.0373 (2)	0.00913 (19)	0.01168 (18)	0.00992 (19)
O1	0.0466 (8)	0.0449 (8)	0.0664 (9)	0.0112 (6)	0.0064 (7)	0.0118 (7)
O2	0.0573 (9)	0.0564 (9)	0.0567 (8)	0.0020 (7)	0.0148 (7)	-0.0172 (7)
O3	0.0401 (7)	0.0969 (12)	0.0360 (7)	0.0060 (8)	0.0031 (6)	0.0088 (7)
O4	0.0668 (10)	0.0444 (8)	0.0697 (9)	0.0160 (7)	0.0354 (8)	0.0182 (7)
N1	0.0390 (8)	0.0422 (8)	0.0361 (7)	-0.0013 (6)	0.0133 (6)	0.0008 (6)
N2	0.0426 (9)	0.1027 (17)	0.0438 (9)	0.0165 (10)	0.0037 (7)	0.0140 (10)
C1	0.0340 (8)	0.0449 (10)	0.0468 (10)	-0.0028 (8)	0.0067 (7)	0.0042 (8)
C2	0.0347 (9)	0.0589 (13)	0.0679 (13)	0.0040 (9)	0.0145 (9)	0.0136 (11)
C3	0.0513 (12)	0.0765 (16)	0.0834 (16)	0.0059 (12)	0.0371 (12)	0.0146 (14)
C4	0.0654 (15)	0.0889 (19)	0.0633 (14)	0.0073 (14)	0.0350 (12)	0.0226 (13)
C5	0.0492 (11)	0.0615 (13)	0.0510 (11)	0.0031 (10)	0.0147 (9)	0.0201 (10)
C6	0.0326 (8)	0.0389 (9)	0.0460 (9)	-0.0015 (7)	0.0085 (7)	0.0053 (8)
C7	0.0395 (9)	0.0388 (9)	0.0352 (8)	-0.0038 (7)	0.0128 (7)	-0.0001 (7)
C8	0.0444 (10)	0.0430 (10)	0.0532 (11)	0.0009 (8)	0.0189 (8)	-0.0003 (8)
C9	0.0690 (14)	0.0522 (12)	0.0646 (13)	-0.0004 (11)	0.0353 (11)	-0.0124 (10)
C10	0.0861 (18)	0.0632 (14)	0.0464 (12)	-0.0049 (13)	0.0208 (11)	-0.0143 (10)
C11	0.0643 (14)	0.0713 (15)	0.0459 (11)	0.0031 (12)	0.0005 (10)	-0.0055 (11)
C12	0.0429 (10)	0.0574 (12)	0.0459 (10)	0.0052 (9)	0.0089 (8)	-0.0034 (9)
C13	0.0341 (8)	0.0462 (10)	0.0355 (8)	0.0042 (7)	0.0116 (7)	0.0048 (7)
C14	0.0454 (11)	0.0551 (12)	0.0541 (11)	-0.0023 (9)	0.0157 (9)	-0.0107 (10)
C15	0.0390 (10)	0.0646 (14)	0.0620 (13)	-0.0083 (10)	0.0104 (9)	-0.0056 (11)
C16	0.0384 (10)	0.0553 (12)	0.0573 (12)	0.0063 (9)	0.0189 (9)	0.0121 (9)
C17	0.0546 (12)	0.0540 (12)	0.0748 (15)	0.0035 (10)	0.0291 (11)	-0.0118 (11)
C18	0.0416 (10)	0.0533 (12)	0.0627 (12)	-0.0037 (9)	0.0168 (9)	-0.0106 (10)
C19	0.0484 (12)	0.0763 (17)	0.100 (2)	0.0100 (12)	0.0359 (13)	0.0127 (15)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.4206 (15)	C8—C9	1.379 (3)
S1—O1	1.4248 (14)	C8—H8A	0.9300
S1—N1	1.7002 (16)	C9—C10	1.378 (3)
S1—C6	1.7378 (19)	C9—H9A	0.9300
S2—O4	1.4237 (16)	C10—C11	1.382 (4)
S2—O3	1.4241 (15)	C10—H10A	0.9300
S2—N1	1.6698 (15)	C11—C12	1.380 (3)
S2—C13	1.7543 (18)	C11—H11A	0.9300

N1—C7	1.447 (2)	C12—H12A	0.9300
N2—C1	1.349 (3)	C13—C14	1.382 (3)
N2—H2B	0.8600	C13—C18	1.389 (3)
N2—H2C	0.8600	C14—C15	1.384 (3)
C1—C2	1.398 (3)	C14—H14A	0.9300
C1—C6	1.414 (2)	C15—C16	1.380 (3)
C2—C3	1.357 (3)	C15—H15A	0.9300
C2—H2D	0.9300	C16—C17	1.377 (3)
C3—C4	1.394 (3)	C16—C19	1.504 (3)
C3—H3B	0.9300	C17—C18	1.382 (3)
C4—C5	1.367 (3)	C17—H17A	0.9300
C4—H4B	0.9300	C18—H18A	0.9300
C5—C6	1.397 (3)	C19—H19A	0.9600
C5—H5A	0.9300	C19—H19B	0.9600
C7—C8	1.380 (3)	C19—H19C	0.9600
C7—C12	1.389 (3)		
O2—S1—O1	119.11 (10)	C9—C8—C7	119.4 (2)
O2—S1—N1	106.54 (9)	C9—C8—H8A	120.3
O1—S1—N1	106.09 (8)	C7—C8—H8A	120.3
O2—S1—C6	112.39 (9)	C10—C9—C8	119.9 (2)
O1—S1—C6	109.36 (9)	C10—C9—H9A	120.1
N1—S1—C6	101.64 (8)	C8—C9—H9A	120.1
O4—S2—O3	120.08 (10)	C9—C10—C11	120.6 (2)
O4—S2—N1	105.17 (8)	C9—C10—H10A	119.7
O3—S2—N1	108.43 (9)	C11—C10—H10A	119.7
O4—S2—C13	108.07 (9)	C12—C11—C10	120.1 (2)
O3—S2—C13	108.12 (9)	C12—C11—H11A	120.0
N1—S2—C13	106.17 (8)	C10—C11—H11A	120.0
C7—N1—S2	117.37 (12)	C11—C12—C7	118.8 (2)
C7—N1—S1	114.93 (11)	C11—C12—H12A	120.6
S2—N1—S1	127.69 (9)	C7—C12—H12A	120.6
C1—N2—H2B	120.0	C14—C13—C18	121.05 (18)
C1—N2—H2C	120.0	C14—C13—S2	119.43 (15)
H2B—N2—H2C	120.0	C18—C13—S2	119.45 (15)
N2—C1—C2	119.54 (18)	C13—C14—C15	118.9 (2)
N2—C1—C6	123.36 (18)	C13—C14—H14A	120.6
C2—C1—C6	117.10 (18)	C15—C14—H14A	120.6
C3—C2—C1	121.5 (2)	C16—C15—C14	121.5 (2)
C3—C2—H2D	119.2	C16—C15—H15A	119.3
C1—C2—H2D	119.2	C14—C15—H15A	119.3
C2—C3—C4	121.1 (2)	C17—C16—C15	118.16 (19)
C2—C3—H3B	119.5	C17—C16—C19	120.9 (2)
C4—C3—H3B	119.5	C15—C16—C19	121.0 (2)
C5—C4—C3	119.3 (2)	C16—C17—C18	122.3 (2)
C5—C4—H4B	120.3	C16—C17—H17A	118.9
C3—C4—H4B	120.3	C18—C17—H17A	118.9
C4—C5—C6	120.2 (2)	C17—C18—C13	118.12 (19)
C4—C5—H5A	119.9	C17—C18—H18A	120.9
C6—C5—H5A	119.9	C13—C18—H18A	120.9

supplementary materials

C5—C6—C1	120.73 (18)	C16—C19—H19A	109.5
C5—C6—S1	117.82 (14)	C16—C19—H19B	109.5
C1—C6—S1	120.92 (14)	H19A—C19—H19B	109.5
C8—C7—C12	121.20 (18)	C16—C19—H19C	109.5
C8—C7—N1	119.59 (16)	H19A—C19—H19C	109.5
C12—C7—N1	119.15 (17)	H19B—C19—H19C	109.5
O4—S2—N1—C7	-27.73 (15)	S2—N1—C7—C8	85.99 (19)
O3—S2—N1—C7	-157.36 (13)	S1—N1—C7—C8	-95.07 (18)
C13—S2—N1—C7	86.67 (14)	S2—N1—C7—C12	-96.67 (19)
O4—S2—N1—S1	153.48 (12)	S1—N1—C7—C12	82.28 (19)
O3—S2—N1—S1	23.85 (15)	C12—C7—C8—C9	-1.3 (3)
C13—S2—N1—S1	-92.12 (13)	N1—C7—C8—C9	176.02 (18)
O2—S1—N1—C7	-175.64 (13)	C7—C8—C9—C10	0.4 (3)
O1—S1—N1—C7	-47.78 (15)	C8—C9—C10—C11	0.4 (4)
C6—S1—N1—C7	66.53 (14)	C9—C10—C11—C12	-0.4 (4)
O2—S1—N1—S2	3.18 (14)	C10—C11—C12—C7	-0.4 (4)
O1—S1—N1—S2	131.04 (12)	C8—C7—C12—C11	1.2 (3)
C6—S1—N1—S2	-114.65 (12)	N1—C7—C12—C11	-176.1 (2)
N2—C1—C2—C3	-178.3 (2)	O4—S2—C13—C14	16.10 (18)
C6—C1—C2—C3	0.6 (3)	O3—S2—C13—C14	147.52 (16)
C1—C2—C3—C4	-0.2 (4)	N1—S2—C13—C14	-96.30 (17)
C2—C3—C4—C5	-0.6 (4)	O4—S2—C13—C18	-161.03 (16)
C3—C4—C5—C6	0.8 (4)	O3—S2—C13—C18	-29.62 (19)
C4—C5—C6—C1	-0.4 (3)	N1—S2—C13—C18	86.56 (17)
C4—C5—C6—S1	171.3 (2)	C18—C13—C14—C15	0.5 (3)
N2—C1—C6—C5	178.5 (2)	S2—C13—C14—C15	-176.61 (17)
C2—C1—C6—C5	-0.3 (3)	C13—C14—C15—C16	0.1 (3)
N2—C1—C6—S1	7.1 (3)	C14—C15—C16—C17	-0.6 (3)
C2—C1—C6—S1	-171.74 (16)	C14—C15—C16—C19	178.6 (2)
O2—S1—C6—C5	148.79 (17)	C15—C16—C17—C18	0.6 (4)
O1—S1—C6—C5	14.18 (19)	C19—C16—C17—C18	-178.7 (2)
N1—S1—C6—C5	-97.68 (17)	C16—C17—C18—C13	0.0 (4)
O2—S1—C6—C1	-39.51 (19)	C14—C13—C18—C17	-0.5 (3)
O1—S1—C6—C1	-174.13 (15)	S2—C13—C18—C17	176.55 (17)
N1—S1—C6—C1	74.01 (17)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2B \cdots O3 ⁱ	0.86	2.20	3.062 (2)	176.
C19—H19A \cdots O1 ⁱⁱ	0.96	2.57	3.517 (3)	169.
C19—H19C \cdots O4 ⁱⁱⁱ	0.96	2.58	3.538 (3)	174.
N2—H2C \cdots O2	0.86	2.23	2.893 (2)	133.

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x+1, y+1/2, -z+1/2$; (iii) $-x+1, y-1/2, -z+1/2$.

Fig. 1

