

3-(2-Chloroanilino)isobenzofuran-1(3H)-one¹

Mustafa Odabaşoğlu^a and Orhan Büyükgüngör^{b*}

^aDepartment of Chemistry, Faculty of Arts and Sciences, Ondokuz Mayıs University, TR-55139 Kurupelit Samsun, Turkey, and ^bDepartment of Physics, Faculty of Arts and Sciences, Ondokuz Mayıs University, TR-55139 Kurupelit Samsun, Turkey
Correspondence e-mail: orhanb@omu.edu.tr


Received 19 March 2008; accepted 20 March 2008

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.030; wR factor = 0.068; data-to-parameter ratio = 14.5.

In the molecule of the title compound, $\text{C}_{14}\text{H}_{10}\text{ClNO}_2$, the essentially planar phthalide group is oriented at a dihedral angle of $59.43(4)^\circ$ with respect to the substituted aromatic ring. In the crystal structure, intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, generating $R_4^4(21)$ ring motifs to form a three-dimensional network.

Related literature

For general background, see: Aoki *et al.* (1973, 1974); Tsui & Tan (1997); Roy & Sarkar (2005); Bellasio (1974, 1975). For related structures, see: Büyükgüngör & Odabaşoğlu (2006); Odabaşoğlu & Büyükgüngör (2006). For ring motif details, see: Bernstein *et al.* (1995); Etter (1990).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{10}\text{ClNO}_2$	$V = 1245.25(19)\text{ \AA}^3$
$M_r = 259.68$	$Z = 4$
Monoclinic, Cc	$\text{Mo K}\alpha$ radiation
$a = 9.2485(8)\text{ \AA}$	$\mu = 0.30\text{ mm}^{-1}$
$b = 22.7915(13)\text{ \AA}$	$T = 296\text{ K}$
$c = 7.1111(6)\text{ \AA}$	$0.51 \times 0.34 \times 0.11\text{ mm}$
$\beta = 123.823(6)^\circ$	

¹ 3-Substituted phthalides. Part XXXIV.

Data collection

Stoe IPDSII diffractometer	7423 measured reflections
Absorption correction: integration (<i>X-RED32</i> ; Stoe & Cie, 2002)	2433 independent reflections
$T_{\min} = 0.880$, $T_{\max} = 0.969$	2200 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.067$	$\Delta\rho_{\text{max}} = 0.12\text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\text{min}} = -0.15\text{ e \AA}^{-3}$
2433 reflections	Absolute structure: Flack (1983), 1205 Friedel pairs
168 parameters	Flack parameter: 0.01 (5)
2 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 \cdots O1 ⁱ	0.84 (3)	2.29 (3)	3.091 (2)	159 (2)
C4—H4 \cdots O2 ⁱⁱ	0.93	2.54	3.397 (2)	153

Symmetry codes: (i) $x, -y + 1, z - \frac{1}{2}$; (ii) $x + 1, -y + 1, z + \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDSII diffractometer (purchased under grant F.279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2437).

References

- Aoki, K., Furusho, T., Kimura, T., Satake, K. & Funayama, S. (1973). Jpn Patent No. 7 324 724.
- Aoki, K., Furusho, T., Kimura, T., Satake, K. & Funayama, S. (1974). *Chem. Abstr.* **80**, 129246.
- Bellasio, E. (1974). German Patent No. 2 422 193.
- Bellasio, E. (1975). *Chem. Abstr.* **83**, 9788.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Büyükgüngör, O. & Odabaşoğlu, M. (2006). *Acta Cryst. E* **62**, o2003–o2004.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Odabaşoğlu, M. & Büyükgüngör, O. (2006). *Acta Cryst. E* **62**, o1879–o1881.
- Roy, H. N. & Sarkar, M. S. (2005). *Synth. Commun.* **35**, 2177–2181.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
- Tsi, D. & Tan, B. K. H. (1997). *Phytother. Res.* **11**, 576–582.

supplementary materials

Acta Cryst. (2008). E64, o754 [doi:10.1107/S160053680800771X]

3-(2-Chloroanilino)isobenzofuran-1(3H)-one

M. Odabasoglu and O. Büyükgüngör

Comment

Phthalides (isobenzofuranones) are five-membered lactones found in plants and they are known to show diverse biological activities, such as fungicidal, bactericidal, herbicidal, analgesic, pesticidal, hypotensive and vasorelaxant activities (Aoki *et al.*, 1973; Tsi & Tan, 1997; Roy & Sarkar, 2005). In addition, phthalide derivatives are useful in the treatment of circulatory and heart-related diseases (Bellasio, 1974). As part of our ongoing research on 3-substituted phthalides, the title compound, (I), has been synthesized and its crystal structure is reported here.

In the molecule of (I), (Fig. 1), rings A (C2-C7), B (C1/C2/C7/C8/O2) and C (C9-C14) are, of course, planar. The dihedral angles between them are A/B = 2.45 (4) $^{\circ}$, A/C = 59.93 (4) $^{\circ}$ and B/C = 58.90 (4) $^{\circ}$. So, rings A and B are also nearly coplanar. Ring C is oriented with respect to the coplanar ring system at a dihedral angle of 59.43 (4) $^{\circ}$. The geometry of (I) does not show any significant difference from the average geometry found for 3-(4-chloroanilino)isobenzofuran-1(3H)-one (Büyükgüngör & Odabaşoğlu, 2006).

In the crystal structure, intermolecular C-H \cdots O and N-H \cdots O hydrogen bonds (Table 1) link the molecules, generating R₄⁴(21) (Fig. 2) ring motifs (Bernstein *et al.*, 1995; Etter, 1990), to form a three-dimensional network, in which they may be effective in the stabilization of the structure.

Experimental

The title compound was prepared according to the method described by Odabaşoğlu & Büyükgüngör (2006), using phthalaldehydic acid and 2-chloroaniline as starting materials (yield: 84%). Crystals of (I) suitable for X-ray analysis were obtained by slow evaporation of an ethanol-DMF (1:1) solution at room temperature.

Refinement

H atom (for NH) was located in difference synthesis and refined freely [N-H = 0.84 (3) Å and U_{iso}(H) = 0.061 (6) Å²]. The remaining H atoms were positioned geometrically, with C-H = 0.93 and 0.98 Å for aromatic and methine H, respectively, and constrained to ride on their parent atoms with U_{iso}(H) = 1.2U_{eq}(C).

supplementary materials

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.



Fig. 2. A partial packing diagram of (I), showing the formation of $R_4^4(21)$ ring motif. Hydrogen bonds are shown as dashed lines [symmetry codes: (i) $x - 1/2, 1 - y, z - 1/2$; (ii) $x - 1, y, z$; (iii) $x, 1 - y, 1/2 + z$]. H atoms not involved in hydrogen bondings have been omitted for clarity.

3-(2-chloroanilino)isobenzofuran-1(3H)-one

Crystal data

$C_{14}H_{10}ClNO_2$	$F_{000} = 536$
$M_r = 259.68$	$D_x = 1.385 \text{ Mg m}^{-3}$
Monoclinic, Cc	Mo $K\alpha$ radiation
Hall symbol: C -2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 9.2485 (8) \text{ \AA}$	Cell parameters from 7423 reflections
$b = 22.7915 (13) \text{ \AA}$	$\theta = 1.8\text{--}28.0^\circ$
$c = 7.1111 (6) \text{ \AA}$	$\mu = 0.30 \text{ mm}^{-1}$
$\beta = 123.823 (6)^\circ$	$T = 296 \text{ K}$
$V = 1245.25 (19) \text{ \AA}^3$	Prism, colorless
$Z = 4$	$0.51 \times 0.34 \times 0.11 \text{ mm}$

Data collection

Stoe IPDS II diffractometer	2433 independent reflections
Monochromator: plane graphite	2200 reflections with $I > 2\sigma(I)$
Detector resolution: 6.67 pixels mm^{-1}	$R_{\text{int}} = 0.038$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 26.0^\circ$
w-scan rotation method	$\theta_{\text{min}} = 1.8^\circ$
Absorption correction: integration (X-RED32; Stoe & Cie, 2002)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.880, T_{\text{max}} = 0.969$	$k = -28 \rightarrow 28$
7423 measured reflections	$l = -8 \rightarrow 8$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.029$	$w = 1/[\sigma^2(F_o^2) + (0.0412P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.067$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.05$	$\Delta\rho_{\max} = 0.12 \text{ e \AA}^{-3}$
2433 reflections	$\Delta\rho_{\min} = -0.15 \text{ e \AA}^{-3}$
168 parameters	Extinction correction: none
2 restraints	Absolute structure: Flack (1983), with 1205 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.01 (5)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.32584 (7)	0.29431 (2)	-0.03793 (8)	0.06502 (16)
O1	0.48884 (19)	0.56863 (6)	0.4723 (2)	0.0568 (4)
O2	0.43774 (16)	0.47290 (5)	0.4752 (2)	0.0462 (3)
N1	0.4488 (2)	0.37749 (6)	0.3398 (3)	0.0446 (3)
H1	0.462 (3)	0.3829 (10)	0.233 (5)	0.061 (6)*
C1	0.5400 (2)	0.51868 (7)	0.5030 (3)	0.0418 (4)
C2	0.7123 (2)	0.49617 (8)	0.5728 (3)	0.0401 (4)
C3	0.8602 (3)	0.52646 (9)	0.6291 (3)	0.0507 (4)
H3	0.8604	0.5672	0.6209	0.061*
C4	1.0076 (3)	0.49429 (11)	0.6977 (3)	0.0616 (5)
H4	1.1095	0.5134	0.7368	0.074*
C5	1.0055 (3)	0.43377 (11)	0.7092 (4)	0.0622 (6)
H5	1.1068	0.4129	0.7568	0.075*
C6	0.8566 (3)	0.40333 (9)	0.6516 (3)	0.0545 (5)

supplementary materials

H6	0.8558	0.3626	0.6584	0.065*
C7	0.7097 (2)	0.43588 (7)	0.5838 (3)	0.0404 (4)
C8	0.5336 (2)	0.41624 (7)	0.5234 (3)	0.0400 (3)
H8	0.5456	0.3977	0.6558	0.048*
C9	0.2979 (2)	0.34797 (7)	0.2808 (3)	0.0385 (4)
C10	0.2263 (3)	0.30630 (8)	0.1070 (3)	0.0473 (4)
C11	0.0825 (3)	0.27393 (9)	0.0501 (3)	0.0610 (6)
H11	0.0388	0.2462	-0.0651	0.073*
C12	0.0027 (3)	0.28242 (11)	0.1632 (4)	0.0678 (6)
H12	-0.0959	0.2610	0.1235	0.081*
C13	0.0701 (3)	0.32296 (9)	0.3359 (4)	0.0588 (5)
H13	0.0174	0.3285	0.4141	0.071*
C14	0.2156 (2)	0.35546 (8)	0.3937 (3)	0.0488 (4)
H14	0.2592	0.3828	0.5101	0.059*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0858 (4)	0.0647 (3)	0.0552 (2)	-0.0121 (3)	0.0459 (3)	-0.0150 (2)
O1	0.0704 (10)	0.0411 (7)	0.0633 (8)	0.0097 (6)	0.0399 (8)	0.0004 (6)
O2	0.0393 (6)	0.0438 (7)	0.0576 (7)	0.0011 (5)	0.0282 (6)	-0.0031 (5)
N1	0.0508 (9)	0.0446 (8)	0.0439 (8)	-0.0085 (7)	0.0297 (7)	-0.0057 (6)
C1	0.0468 (10)	0.0416 (9)	0.0371 (8)	0.0008 (8)	0.0234 (7)	-0.0028 (7)
C2	0.0406 (9)	0.0446 (9)	0.0332 (8)	-0.0032 (7)	0.0192 (7)	-0.0032 (6)
C3	0.0494 (11)	0.0577 (11)	0.0431 (8)	-0.0141 (9)	0.0245 (8)	-0.0071 (8)
C4	0.0403 (11)	0.0914 (16)	0.0487 (11)	-0.0170 (10)	0.0221 (9)	-0.0092 (10)
C5	0.0379 (10)	0.0905 (17)	0.0527 (11)	0.0122 (10)	0.0218 (9)	0.0000 (10)
C6	0.0496 (11)	0.0555 (11)	0.0552 (10)	0.0111 (9)	0.0271 (9)	0.0032 (9)
C7	0.0379 (9)	0.0445 (9)	0.0363 (8)	0.0015 (7)	0.0192 (7)	-0.0013 (7)
C8	0.0408 (9)	0.0381 (8)	0.0404 (8)	0.0015 (7)	0.0220 (7)	0.0001 (7)
C9	0.0399 (9)	0.0313 (8)	0.0405 (8)	0.0008 (7)	0.0201 (7)	0.0033 (6)
C10	0.0552 (11)	0.0431 (9)	0.0378 (8)	-0.0032 (8)	0.0223 (8)	-0.0001 (7)
C11	0.0629 (14)	0.0537 (11)	0.0523 (11)	-0.0203 (10)	0.0233 (10)	-0.0118 (9)
C12	0.0591 (13)	0.0671 (14)	0.0748 (14)	-0.0228 (10)	0.0358 (12)	-0.0053 (10)
C13	0.0551 (12)	0.0593 (13)	0.0726 (13)	-0.0092 (10)	0.0422 (11)	-0.0001 (10)
C14	0.0535 (11)	0.0456 (10)	0.0522 (10)	-0.0025 (8)	0.0324 (9)	-0.0022 (7)

Geometric parameters (\AA , $^\circ$)

N1—H1	0.84 (3)	C8—N1	1.400 (2)
C1—O1	1.205 (2)	C8—O2	1.494 (2)
C1—O2	1.347 (2)	C8—H8	0.9800
C1—C2	1.472 (3)	C9—N1	1.387 (2)
C2—C7	1.377 (2)	C9—C14	1.391 (2)
C2—C3	1.378 (3)	C9—C10	1.399 (2)
C3—C4	1.376 (3)	C10—C11	1.370 (3)
C3—H3	0.9300	C10—Cl1	1.744 (2)
C4—C5	1.383 (4)	C11—C12	1.375 (3)
C4—H4	0.9300	C11—H11	0.9300

C5—C6	1.385 (3)	C12—C13	1.377 (3)
C5—H5	0.9300	C12—H12	0.9300
C6—C7	1.378 (3)	C13—C14	1.383 (3)
C6—H6	0.9300	C13—H13	0.9300
C7—C8	1.503 (3)	C14—H14	0.9300
C1—O2—C8	110.92 (13)	N1—C8—O2	112.25 (14)
C8—N1—H1	117.4 (16)	N1—C8—C7	114.17 (15)
C9—N1—C8	122.34 (16)	O2—C8—C7	102.62 (13)
C9—N1—H1	115.2 (17)	N1—C8—H8	109.2
O1—C1—O2	122.15 (17)	O2—C8—H8	109.2
O1—C1—C2	129.22 (17)	C7—C8—H8	109.2
O2—C1—C2	108.63 (14)	N1—C9—C14	123.22 (15)
C7—C2—C3	121.94 (17)	N1—C9—C10	119.92 (15)
C7—C2—C1	108.50 (15)	C14—C9—C10	116.80 (16)
C3—C2—C1	129.53 (17)	C11—C10—C9	122.08 (18)
C4—C3—C2	117.61 (19)	C11—C10—Cl1	119.09 (14)
C4—C3—H3	121.2	C9—C10—Cl1	118.82 (14)
C2—C3—H3	121.2	C10—C11—C12	120.02 (19)
C3—C4—C5	120.61 (19)	C10—C11—H11	120.0
C3—C4—H4	119.7	C12—C11—H11	120.0
C5—C4—H4	119.7	C11—C12—C13	119.48 (19)
C4—C5—C6	121.76 (19)	C11—C12—H12	120.3
C4—C5—H5	119.1	C13—C12—H12	120.3
C6—C5—H5	119.1	C12—C13—C14	120.4 (2)
C7—C6—C5	117.26 (19)	C12—C13—H13	119.8
C7—C6—H6	121.4	C14—C13—H13	119.8
C5—C6—H6	121.4	C13—C14—C9	121.18 (17)
C2—C7—C6	120.82 (17)	C13—C14—H14	119.4
C2—C7—C8	109.32 (14)	C9—C14—H14	119.4
C6—C7—C8	129.80 (16)		
O1—C1—O2—C8	179.77 (16)	C2—C7—C8—O2	0.52 (17)
C2—C1—O2—C8	-0.24 (16)	C6—C7—C8—O2	177.69 (17)
O1—C1—C2—C7	-179.43 (17)	N1—C8—O2—C1	-123.18 (15)
O2—C1—C2—C7	0.59 (17)	C7—C8—O2—C1	-0.16 (16)
O1—C1—C2—C3	2.8 (3)	O2—C8—N1—C9	-73.5 (2)
O2—C1—C2—C3	-177.14 (15)	C7—C8—N1—C9	170.20 (15)
C7—C2—C3—C4	0.0 (3)	C14—C9—N1—C8	1.9 (3)
C1—C2—C3—C4	177.43 (17)	C10—C9—N1—C8	-174.97 (15)
C2—C3—C4—C5	0.0 (3)	N1—C9—C10—C11	176.68 (18)
C3—C4—C5—C6	0.3 (3)	C14—C9—C10—C11	-0.3 (2)
C4—C5—C6—C7	-0.5 (3)	N1—C9—C10—Cl1	-2.1 (2)
C3—C2—C7—C6	-0.2 (3)	C14—C9—C10—Cl1	-179.13 (13)
C1—C2—C7—C6	-178.15 (15)	C9—C10—C11—C12	0.8 (3)
C3—C2—C7—C8	177.25 (14)	Cl1—C10—C11—C12	179.56 (18)
C1—C2—C7—C8	-0.68 (18)	C10—C11—C12—C13	-1.0 (4)
C5—C6—C7—C2	0.5 (3)	C11—C12—C13—C14	0.8 (4)
C5—C6—C7—C8	-176.40 (17)	C12—C13—C14—C9	-0.3 (3)
C2—C7—C8—N1	122.25 (16)	N1—C9—C14—C13	-176.81 (18)

supplementary materials

C6—C7—C8—N1

−60.6 (2)

C10—C9—C14—C13

0.1 (3)

Hydrogen-bond geometry (Å, °)

D—H···A

D—H

H···A

D···A

D—H···A

N1—H1···O1ⁱ

0.84 (3)

2.29 (3)

3.091 (2)

159 (2)

C4—H4···O2ⁱⁱ

0.93


2.54

3.397 (2)

153

Symmetry codes: (i) $x, -y+1, z-1/2$; (ii) $x+1, -y+1, z+1/2$.

Fig. 1



supplementary materials

Fig. 2

