

N'-(2-Hydroxynaphthylidene)-4-methoxybenzohydrazide

Chun-Bao Tang

Department of Chemistry, Jiaying University, Meizhou 514015, People's Republic of China

Correspondence e-mail: chunbao_tang@163.com


Received 21 March 2008; accepted 25 March 2008

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.056; wR factor = 0.137; data-to-parameter ratio = 16.0.

The title Schiff base compound, $\text{C}_{19}\text{H}_{16}\text{N}_2\text{O}_3$, was derived from the condensation reaction of 2-hydroxy-1-naphthaldehyde with 4-methoxybenzohydrazide. The dihedral angle between the benzene ring and the naphthalene ring system is $6.8(2)^\circ$. In the crystal structure, molecules are linked through intermolecular $\text{N}-\text{H}\cdots\text{O}$ intermolecular hydrogen bonds, forming chains running along the c axis.

Related literature

For related structures, see: Tang (2006, 2007a,b,c,d). For reference structural data, see: Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{16}\text{N}_2\text{O}_3$ $M_r = 320.34$ Monoclinic, P_{2_1}/c $a = 11.159(2)\text{ \AA}$ $b = 15.790(3)\text{ \AA}$ $c = 8.8300(18)\text{ \AA}$ $\beta = 91.70(3)^\circ$ $V = 1555.2(5)\text{ \AA}^3$ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.09\text{ mm}^{-1}$
 $T = 298(2)\text{ K}$ $0.32 \times 0.32 \times 0.30\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.971$, $T_{\max} = 0.972$ 13232 measured reflections
3550 independent reflections
2161 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.136$
 $S = 1.04$
3550 reflections
222 parameters
1 restraintH atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.20\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots N1	0.82	1.86	2.582 (2)	146
N2—H2 \cdots O2 ⁱ	0.904 (9)	1.957 (12)	2.834 (2)	163 (2)

Symmetry code: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Financial support from the Jiaying University Research Fund is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2478).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bruker (2002). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Tang, C.-B. (2006). *Acta Cryst. E* **62**, m2629–m2630.
Tang, C.-B. (2007a). *Acta Cryst. E* **63**, m2654.
Tang, C.-B. (2007b). *Acta Cryst. E* **63**, m2785–m2786.
Tang, C.-B. (2007c). *Acta Cryst. E* **63**, o4545.
Tang, C.-B. (2007d). *Acta Cryst. E* **63**, o4841.

supplementary materials

Acta Cryst. (2008). E64, o768 [doi:10.1107/S1600536808008076]

N'-(2-Hydroxynaphthylidene)-4-methoxybenzohydrazide

C.-B. Tang

Comment

Recently, the author has reported the structures of several Schiff base compounds (Tang, 2006, 2007*a,b,c,d*) and, in continuation of work in this area, reports herein the structure of the title compound, (I), Fig. 1, a new Schiff base compound.

In the title compound (Fig. 1), the dihedral angle between the benzene ring and the naphtyl ring is 6.8 (2)°. The torsion angles C1—C11—N1—N2, C11—N1—N2—C12, and N1—N2—C12—C13 are 1.3 (2), 17.0 (2), and 1.5 (2)°, respectively. All the bond lengths are within normal values (Allen *et al.*, 1987).

In the crystal structure of the compound, molecules are linked through N—H···O intermolecular hydrogen bonds (Table 1), forming chains running along the *c* axis (Fig. 2).

Experimental

2-Hydroxy-1-naphthylaldehyde (0.1 mmol, 17.2 mg) and 4-methoxybenzohydrazide (0.1 mmol, 16.6 mg) were dissolved in an ethanol solution (20 ml). The mixture was stirred at reflux for 10 min to give a clear colorless solution. Colorless needle-like crystals of the compound were formed by slow evaporation of the solvent over several days.

Refinement

H2 atom was located from a difference Fourier map and refined isotropically, with N—H distance restrained to 0.90 (1) Å. Other H atoms were constrained to ideal geometries, with C—H = 0.93–0.96 Å, O—H = 0.82 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$, $1.5U_{\text{eq}}(\text{C}19 \text{ and } \text{O}1)$.

Figures




Fig. 1. The molecular structure of the compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.



Fig. 2. Molecular packing of (I) with hydrogen bonds drawn as dashed lines.

supplementary materials

N¹-(2-Hydroxynaphthylidene)-4-methoxybenzohydrazide

Crystal data

C ₁₉ H ₁₆ N ₂ O ₃	$F_{000} = 672$
$M_r = 320.34$	$D_x = 1.368 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 11.159 (2) \text{ \AA}$	Cell parameters from 1378 reflections
$b = 15.790 (3) \text{ \AA}$	$\theta = 2.2\text{--}24.5^\circ$
$c = 8.8300 (18) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 91.70 (3)^\circ$	$T = 298 (2) \text{ K}$
$V = 1555.2 (5) \text{ \AA}^3$	Cut from a needle, colorless
$Z = 4$	$0.32 \times 0.32 \times 0.30 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	3550 independent reflections
Radiation source: fine-focus sealed tube	2161 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.052$
$T = 298(2) \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 1.8^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -14 \rightarrow 14$
$T_{\text{min}} = 0.971, T_{\text{max}} = 0.972$	$k = -20 \rightarrow 20$
13232 measured reflections	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.056$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.136$	$w = 1/[\sigma^2(F_o^2) + (0.0391P)^2 + 0.168P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\text{max}} = 0.001$
3550 reflections	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
222 parameters	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.44682 (14)	0.12307 (9)	0.77419 (16)	0.0569 (4)
H1	0.4089	0.1584	0.7244	0.085*
O2	0.24817 (13)	0.32717 (8)	0.67361 (14)	0.0515 (4)
O3	0.02184 (13)	0.57256 (9)	0.17402 (17)	0.0602 (5)
N1	0.30518 (14)	0.17750 (9)	0.55897 (17)	0.0415 (4)
N2	0.25139 (15)	0.24056 (10)	0.47098 (17)	0.0423 (4)
C1	0.35505 (16)	0.03145 (12)	0.5862 (2)	0.0362 (4)
C2	0.42732 (17)	0.04559 (13)	0.7145 (2)	0.0431 (5)
C3	0.48691 (18)	-0.02246 (15)	0.7874 (2)	0.0520 (6)
H3	0.5370	-0.0121	0.8715	0.062*
C4	0.47221 (19)	-0.10234 (14)	0.7367 (2)	0.0526 (6)
H4	0.5124	-0.1462	0.7868	0.063*
C5	0.39740 (17)	-0.12109 (13)	0.6096 (2)	0.0434 (5)
C6	0.3824 (2)	-0.20495 (13)	0.5558 (3)	0.0561 (6)
H6	0.4230	-0.2489	0.6051	0.067*
C7	0.3101 (2)	-0.22238 (14)	0.4342 (3)	0.0608 (6)
H7	0.3007	-0.2779	0.4007	0.073*
C8	0.2498 (2)	-0.15638 (14)	0.3591 (3)	0.0586 (6)
H8	0.2005	-0.1683	0.2750	0.070*
C9	0.26197 (18)	-0.07449 (12)	0.4075 (2)	0.0458 (5)
H9	0.2198	-0.0318	0.3566	0.055*
C10	0.33734 (16)	-0.05348 (11)	0.5331 (2)	0.0371 (5)
C11	0.29956 (17)	0.10221 (12)	0.5053 (2)	0.0388 (5)
H11	0.2594	0.0926	0.4131	0.047*
C12	0.22727 (17)	0.31534 (11)	0.5375 (2)	0.0376 (5)
C13	0.17287 (17)	0.38188 (11)	0.4404 (2)	0.0365 (4)
C14	0.18459 (18)	0.46552 (12)	0.4857 (2)	0.0454 (5)
H14	0.2259	0.4778	0.5761	0.054*
C15	0.13651 (19)	0.53130 (12)	0.4000 (2)	0.0493 (5)
H15	0.1466	0.5871	0.4317	0.059*
C16	0.07353 (17)	0.51333 (12)	0.2674 (2)	0.0427 (5)
C17	0.05869 (18)	0.43022 (12)	0.2214 (2)	0.0453 (5)
H17	0.0147	0.4180	0.1329	0.054*

supplementary materials

C18	0.10864 (17)	0.36569 (12)	0.3058 (2)	0.0406 (5)
H18	0.0995	0.3101	0.2725	0.049*
C19	0.0515 (2)	0.65852 (13)	0.2009 (3)	0.0685 (7)
H19A	0.0224	0.6755	0.2974	0.103*
H19B	0.0153	0.6931	0.1227	0.103*
H19C	0.1370	0.6652	0.2007	0.103*
H2	0.251 (2)	0.2301 (14)	0.3703 (12)	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0614 (11)	0.0580 (10)	0.0505 (10)	-0.0051 (8)	-0.0100 (8)	-0.0063 (8)
O2	0.0791 (11)	0.0422 (8)	0.0326 (8)	0.0005 (7)	-0.0063 (7)	-0.0019 (6)
O3	0.0683 (11)	0.0440 (9)	0.0674 (11)	0.0126 (7)	-0.0113 (8)	0.0084 (8)
N1	0.0508 (11)	0.0356 (9)	0.0380 (9)	0.0008 (8)	0.0009 (8)	0.0047 (7)
N2	0.0608 (11)	0.0320 (9)	0.0339 (9)	0.0046 (8)	-0.0029 (9)	0.0013 (7)
C1	0.0350 (10)	0.0388 (11)	0.0350 (11)	0.0019 (8)	0.0038 (8)	0.0062 (8)
C2	0.0391 (11)	0.0500 (13)	0.0405 (12)	-0.0026 (10)	0.0036 (9)	0.0034 (10)
C3	0.0414 (12)	0.0728 (16)	0.0413 (13)	0.0038 (11)	-0.0065 (10)	0.0114 (11)
C4	0.0465 (13)	0.0549 (14)	0.0564 (14)	0.0122 (10)	0.0035 (11)	0.0204 (11)
C5	0.0371 (11)	0.0461 (12)	0.0474 (12)	0.0058 (9)	0.0077 (9)	0.0119 (10)
C6	0.0589 (15)	0.0399 (13)	0.0700 (16)	0.0138 (11)	0.0121 (12)	0.0144 (11)
C7	0.0684 (16)	0.0394 (13)	0.0750 (17)	0.0035 (11)	0.0085 (14)	-0.0022 (12)
C8	0.0658 (16)	0.0512 (14)	0.0586 (15)	-0.0033 (11)	-0.0020 (12)	-0.0065 (11)
C9	0.0500 (13)	0.0380 (11)	0.0492 (13)	0.0013 (9)	-0.0004 (10)	0.0014 (9)
C10	0.0349 (11)	0.0396 (11)	0.0370 (11)	0.0022 (9)	0.0062 (9)	0.0069 (9)
C11	0.0421 (12)	0.0395 (11)	0.0348 (11)	-0.0008 (9)	0.0006 (9)	0.0035 (9)
C12	0.0423 (11)	0.0359 (11)	0.0346 (11)	-0.0056 (8)	0.0005 (9)	-0.0003 (8)
C13	0.0397 (11)	0.0349 (10)	0.0350 (11)	-0.0017 (8)	0.0033 (8)	-0.0002 (8)
C14	0.0545 (13)	0.0409 (12)	0.0403 (12)	-0.0017 (10)	-0.0045 (10)	-0.0061 (9)
C15	0.0609 (14)	0.0322 (11)	0.0547 (14)	0.0017 (10)	-0.0011 (11)	-0.0036 (10)
C16	0.0408 (11)	0.0412 (12)	0.0458 (12)	0.0050 (9)	-0.0008 (9)	0.0032 (9)
C17	0.0454 (13)	0.0479 (13)	0.0421 (12)	-0.0002 (10)	-0.0077 (10)	-0.0004 (10)
C18	0.0467 (12)	0.0351 (11)	0.0397 (11)	-0.0029 (9)	-0.0020 (9)	-0.0046 (9)
C19	0.0857 (19)	0.0414 (13)	0.0784 (18)	0.0172 (12)	0.0019 (14)	0.0076 (12)

Geometric parameters (\AA , $^\circ$)

O1—C2	1.347 (2)	C7—C8	1.397 (3)
O1—H1	0.8200	C7—H7	0.9300
O2—C12	1.231 (2)	C8—C9	1.367 (3)
O3—C16	1.363 (2)	C8—H8	0.9300
O3—C19	1.415 (2)	C9—C10	1.411 (3)
N1—C11	1.281 (2)	C9—H9	0.9300
N1—N2	1.388 (2)	C11—H11	0.9300
N2—C12	1.350 (2)	C12—C13	1.475 (3)
N2—H2	0.904 (9)	C13—C14	1.385 (2)
C1—C2	1.389 (3)	C13—C18	1.394 (3)
C1—C10	1.432 (2)	C14—C15	1.384 (3)

C1—C11	1.455 (2)	C14—H14	0.9300
C2—C3	1.409 (3)	C15—C16	1.377 (3)
C3—C4	1.347 (3)	C15—H15	0.9300
C3—H3	0.9300	C16—C17	1.382 (3)
C4—C5	1.410 (3)	C17—C18	1.371 (3)
C4—H4	0.9300	C17—H17	0.9300
C5—C6	1.415 (3)	C18—H18	0.9300
C5—C10	1.421 (3)	C19—H19A	0.9600
C6—C7	1.352 (3)	C19—H19B	0.9600
C6—H6	0.9300	C19—H19C	0.9600
C2—O1—H1	109.5	C9—C10—C5	117.31 (18)
C16—O3—C19	117.63 (17)	C9—C10—C1	123.40 (17)
C11—N1—N2	116.28 (16)	C5—C10—C1	119.29 (18)
C12—N2—N1	118.18 (15)	N1—C11—C1	121.04 (18)
C12—N2—H2	126.4 (15)	N1—C11—H11	119.5
N1—N2—H2	114.0 (15)	C1—C11—H11	119.5
C2—C1—C10	119.24 (17)	O2—C12—N2	121.54 (17)
C2—C1—C11	120.36 (18)	O2—C12—C13	121.46 (17)
C10—C1—C11	120.39 (17)	N2—C12—C13	116.99 (16)
O1—C2—C1	123.23 (18)	C14—C13—C18	117.57 (18)
O1—C2—C3	116.45 (18)	C14—C13—C12	118.53 (17)
C1—C2—C3	120.31 (19)	C18—C13—C12	123.89 (17)
C4—C3—C2	120.7 (2)	C15—C14—C13	121.75 (19)
C4—C3—H3	119.6	C15—C14—H14	119.1
C2—C3—H3	119.6	C13—C14—H14	119.1
C3—C4—C5	121.63 (19)	C16—C15—C14	119.34 (19)
C3—C4—H4	119.2	C16—C15—H15	120.3
C5—C4—H4	119.2	C14—C15—H15	120.3
C4—C5—C6	121.66 (19)	O3—C16—C15	124.67 (18)
C4—C5—C10	118.73 (19)	O3—C16—C17	115.39 (18)
C6—C5—C10	119.6 (2)	C15—C16—C17	119.94 (18)
C7—C6—C5	121.3 (2)	C18—C17—C16	120.17 (18)
C7—C6—H6	119.4	C18—C17—H17	119.9
C5—C6—H6	119.4	C16—C17—H17	119.9
C6—C7—C8	119.5 (2)	C17—C18—C13	121.20 (18)
C6—C7—H7	120.2	C17—C18—H18	119.4
C8—C7—H7	120.2	C13—C18—H18	119.4
C9—C8—C7	121.0 (2)	O3—C19—H19A	109.5
C9—C8—H8	119.5	O3—C19—H19B	109.5
C7—C8—H8	119.5	H19A—C19—H19B	109.5
C8—C9—C10	121.29 (19)	O3—C19—H19C	109.5
C8—C9—H9	119.4	H19A—C19—H19C	109.5
C10—C9—H9	119.4	H19B—C19—H19C	109.5

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···N1	0.82	1.86	2.582 (2)	146
N2—H2···O2 ⁱ	0.904 (9)	1.957 (12)	2.834 (2)	163 (2)

supplementary materials

Symmetry codes: (i) $x, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

