

(*meso*-5,5,7,12,12,14-Hexamethyl-1,4,8,11-tetraazacyclotetradecane)-nickel(II) bis(*O,O'*-dibenzyl dithiophosphate)

Bin Xie,* Li-Ke Zou, Yi-Guo He, Jian-Shen Feng and Xiu-Lan Zhang

Department of Chemistry, Sichuan University of Science & Engineering, Zigong, Sichuan 643000, People's Republic of China
Correspondence e-mail: zoulake@yahoo.com.cn


Received 15 March 2008; accepted 28 March 2008

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.050; wR factor = 0.128; data-to-parameter ratio = 16.0.

In the title salt-type 1:2 adduct, $[\text{Ni}(\text{C}_{16}\text{H}_{36}\text{N}_4)](\text{C}_{14}\text{H}_{14}\text{O}_2\text{PS}_2)_2$ or $[\text{Ni}(\text{tet-a})][\text{S}_2\text{P}(\text{OCH}_2\text{Ph})_2]_2$, where tet-a is *meso*-5,5,7,12,12,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane, the $[\text{Ni}(\text{tet-a})]^{2+}$ complex cation exhibits a relatively undistorted square-planar geometry about the Ni atom, which lies on an inversion centre and is coordinated by four macrocyclic N atoms. The two *O,O'*-bis(2-phenylmethyl) dithiophosphate anions act as counter-ions to balance the charge and they interact with the complex through $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds. Important geometric data include Ni—N distances of 1.958 (3) and 1.963 (3) Å.

Related literature

For related literature, see: Burchell *et al.* (2000); Ali *et al.* (2004); Allen (2002); Li *et al.* (2006); Liu *et al.* (1997).



Experimental

Crystal data

$[\text{Ni}(\text{C}_{16}\text{H}_{36}\text{N}_4)](\text{C}_{14}\text{H}_{14}\text{O}_2\text{PS}_2)_2$
 $M_r = 961.88$
Monoclinic, $P2_1/c$
 $a = 16.371$ (5) Å
 $b = 14.917$ (5) Å
 $c = 9.964$ (4) Å
 $\beta = 103.11$ (3)°

$V = 2369.9$ (15) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.70$ mm⁻¹
 $T = 290$ (2) K
0.40 × 0.38 × 0.36 mm

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\text{min}} = 0.768$, $T_{\text{max}} = 0.787$
5571 measured reflections

4393 independent reflections
2880 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$
3 standard reflections every 300 reflections
intensity decay: 1.1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.127$
 $S = 1.01$
4393 reflections

275 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.52$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{S1}$	0.91	2.61	3.390 (3)	144
$\text{N2}-\text{H2}\cdots\text{S2}^i$	0.91	2.50	3.394 (3)	169

Symmetry code: (i) $-x + 1, -y + 1, -z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Financial assistance from the Science and Technology Office of Zigong City, China (Project No. 06 G2045 and 07GX008) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2324).

References

- Ali, M., Ray, A., Sheldrick, W. S., Mayer-Figge, H., Gao, S. & Sahmesd, A. I. (2004). *New J. Chem.* **28**, 412–417.
Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
Burchell, C. J., Ferguson, G., Lough, A. J. & Glidewell, C. (2000). *Acta Cryst.* **B56**, 1054–1062.
Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
Li, Z., Li, J. & Du, S. (2006). *J. Mol. Struct.* **783**, 116–121.
Liu, C. W., Pitts, J. T. & Fackler, J. P. (1997). *Polyhedron*, **16**, 3899–3909.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m622 [doi:10.1107/S1600536808008398]

(*meso*-5,5,7,12,12,14-Hexamethyl-1,4,8,11-tetraazacyclotetradecane)nickel(II) bis(*O,O'*-dibenzyl dithiophosphate)

B. Xie, L.-K. Zou, Y.-G. He, J.-S. Feng and X.-L. Zhang

Comment

Considerable interest in the chemistry of the *O,O'*-dialkyl dithiophosphate complexes of transition metal results from their value to industry, namely, as anti-oxidants, additives to lubricating oils, flotation reagents, insecticides (Liu *et al.*, 1997; Li *et al.*, 2006). We report here the synthesis and crystal structure of salt-type adduct [Ni(tet-a)][S₂P(OCH₂Ph)₂]₂ (tet-a = *meso*-5,5,7,12,12,14-hexamethyl-1,4,8,11-tetraazacyclotetradecane, a tetradentate macrocyclic nitrogen base (Burchell *et al.*, 2000; Ali *et al.*, 2004))

The Ni^{II} in the complex cation [Ni(tet-a)]²⁺ lies on an inversion center and is coordinated by four nitrogen atoms of tetaaza macrocycle in undistorted square-planar geometry (Fig. 1). The bond lengths and angles within the complex may be considered normal in comparison with the Cambridge Structural Database results (Allen, 2002).

The two *O,O'*-di(2-phenylmethyl) dithiophosphates act as counter-ions to balance the charge and they interact with the complex through N—H...S hydrogen bonds (Table 1).

Experimental

A hot solution of tet-a.2H₂O (0.77 g, 2 mmol) and Ni(OAc)₂.2H₂O (0.25 g, 1 mmol) in 20 ml methanol was quickly added to a hot solution of (PhCH₂O)₂PSSNH₂(C₂H₅)₂ (0.77 g, 2 mmol) in 20 ml methanol. The mixture was refluxed for 6 h, then cooled to room temperature, the orange-red precipitate was collected by filtration, washed with small amounts of methanol. A solution of adduct (1) in DMSO was kept at room temperature, and red block crystals suitable for X-ray diffraction studies were obtained in eight months.

Refinement

All H atoms attached to C and N atom were fixed geometrically and treated as riding with C—H = 0.98 Å (methine), 0.97 Å (methylene), 0.96 Å (methyl) or 0.93 Å (aromatic) and N—H = 0.91 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C, N})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl})$.

Figures




Fig. 1. A View of the title compound showing the atom-labelling scheme. Only one of the two anions is represented for clarity. Displacement ellipsoids are drawn at the 30% probability level. Hydrogen-bonds are shown as dashed lines. For the sake of clarity, H atoms bonded to C atoms have been omitted. H atoms are represented as small spheres of arbitrary radii. [Symmetry code: (i) $-x + 1, -y + 1, -z$].

supplementary materials

(*meso*-5,5,7,12,12,14-Hexamethyl-1,4,8,11-tetraazacyclotetradecane)nickel(II) bis(*O,O'*-dibenzyl dithiophosphate)

Crystal data

[Ni(C₁₆H₃₆N₄)](C₁₄H₁₄O₂PS₂)₂

$M_r = 961.88$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 16.371$ (5) Å

$b = 14.917$ (5) Å

$c = 9.964$ (4) Å

$\beta = 103.11$ (3)°

$V = 2369.9$ (15) Å³

$Z = 2$

$F_{000} = 1020$

$D_x = 1.348$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 19 reflections

$\theta = 4.4$ – 5.5 °

$\mu = 0.70$ mm⁻¹

$T = 290$ (2) K

Block, red

$0.40 \times 0.38 \times 0.36$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 290$ (2) K

$\omega/2\theta$ scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.768$, $T_{\max} = 0.787$

5571 measured reflections

4393 independent reflections

2880 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\max} = 25.5$ °

$\theta_{\min} = 1.3$ °

$h = -7 \rightarrow 19$

$k = -18 \rightarrow 0$

$l = -12 \rightarrow 11$

3 standard reflections

every 300 reflections

intensity decay: 1.1%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.049$

$wR(F^2) = 0.127$

$S = 1.01$

4393 reflections

275 parameters

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0689P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.52$ e Å⁻³

$\Delta\rho_{\min} = -0.34$ e Å⁻³

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.5000	0.5000	0.0000	0.02975 (18)
S1	0.39590 (6)	0.45749 (7)	0.26928 (12)	0.0569 (3)
S2	0.28652 (6)	0.64383 (7)	0.13356 (11)	0.0498 (3)
P1	0.29136 (6)	0.52753 (7)	0.22790 (10)	0.0406 (3)
O1	0.26427 (15)	0.53665 (18)	0.3726 (3)	0.0509 (7)
O2	0.21601 (15)	0.47114 (17)	0.1325 (3)	0.0521 (7)
N1	0.57052 (16)	0.53521 (17)	0.1790 (3)	0.0341 (7)
H1	0.5357	0.5294	0.2379	0.041*
N2	0.55476 (16)	0.38269 (17)	0.0308 (3)	0.0332 (6)
H2	0.5975	0.3845	-0.0138	0.040*
C1	0.6346 (2)	0.4648 (2)	0.2251 (4)	0.0543 (12)
H1A	0.6539	0.4663	0.3245	0.065*
H1B	0.6824	0.4747	0.1844	0.065*
C2	0.5955 (2)	0.3765 (2)	0.1806 (4)	0.0478 (10)
H2A	0.5540	0.3619	0.2330	0.057*
H2B	0.6378	0.3298	0.1960	0.057*
C3	0.5074 (2)	0.2981 (2)	-0.0165 (3)	0.0353 (8)
H3	0.4636	0.2913	0.0354	0.042*
C4	0.4652 (2)	0.3050 (2)	-0.1675 (4)	0.0397 (9)
H4A	0.4439	0.2462	-0.1989	0.048*
H4B	0.5077	0.3204	-0.2174	0.048*
C5	0.5636 (2)	0.2146 (2)	0.0096 (4)	0.0452 (9)
H5A	0.6067	0.2194	-0.0413	0.068*
H5B	0.5304	0.1621	-0.0197	0.068*
H5C	0.5889	0.2101	0.1062	0.068*
C6	0.6063 (2)	0.6282 (2)	0.2081 (4)	0.0359 (8)
C7	0.6734 (2)	0.6441 (3)	0.1263 (4)	0.0542 (11)
H7A	0.6523	0.6262	0.0323	0.081*
H7B	0.7225	0.6096	0.1658	0.081*
H7C	0.6877	0.7066	0.1293	0.081*
C8	0.6443 (2)	0.6385 (3)	0.3636 (4)	0.0543 (11)
H8A	0.6602	0.6998	0.3838	0.081*
H8B	0.6928	0.6008	0.3896	0.081*

supplementary materials

H8C	0.6035	0.6212	0.4144	0.081*
C9	0.1889 (3)	0.5872 (3)	0.3782 (4)	0.0624 (12)
H9A	0.1446	0.5736	0.2982	0.075*
H9B	0.2003	0.6510	0.3783	0.075*
C10	0.1620 (2)	0.5621 (3)	0.5064 (4)	0.0511 (10)
C11	0.1766 (3)	0.6179 (3)	0.6203 (5)	0.0639 (12)
H11	0.2020	0.6734	0.6175	0.077*
C12	0.1527 (3)	0.5897 (5)	0.7399 (5)	0.0884 (18)
H12	0.1623	0.6266	0.8171	0.106*
C13	0.1153 (4)	0.5082 (6)	0.7433 (7)	0.103 (2)
H13	0.0995	0.4899	0.8230	0.123*
C14	0.1010 (4)	0.4537 (5)	0.6318 (8)	0.108 (2)
H14	0.0759	0.3980	0.6349	0.130*
C15	0.1237 (3)	0.4810 (4)	0.5149 (6)	0.0822 (15)
H15	0.1130	0.4435	0.4383	0.099*
C16	0.2008 (2)	0.3800 (3)	0.1649 (5)	0.0604 (12)
H16A	0.1953	0.3756	0.2596	0.072*
H16B	0.2473	0.3425	0.1542	0.072*
C17	0.1214 (2)	0.3495 (2)	0.0688 (4)	0.0472 (10)
C18	0.0461 (3)	0.3895 (3)	0.0682 (5)	0.0681 (13)
H18	0.0439	0.4365	0.1287	0.082*
C19	-0.0272 (3)	0.3613 (4)	-0.0213 (5)	0.0795 (16)
H19	-0.0781	0.3884	-0.0193	0.095*
C20	-0.0245 (3)	0.2952 (4)	-0.1100 (6)	0.0853 (18)
H20	-0.0738	0.2751	-0.1682	0.102*
C21	0.0503 (4)	0.2567 (4)	-0.1159 (6)	0.108 (2)
H21	0.0522	0.2126	-0.1811	0.130*
C22	0.1229 (3)	0.2831 (3)	-0.0256 (6)	0.0808 (16)
H22	0.1734	0.2557	-0.0286	0.097*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0297 (3)	0.0263 (3)	0.0308 (3)	0.0014 (2)	0.0017 (2)	0.0013 (3)
S1	0.0404 (5)	0.0557 (6)	0.0758 (8)	0.0085 (5)	0.0157 (5)	0.0121 (6)
S2	0.0506 (6)	0.0459 (6)	0.0535 (6)	-0.0007 (4)	0.0129 (5)	0.0074 (5)
P1	0.0349 (5)	0.0421 (5)	0.0440 (6)	-0.0003 (4)	0.0070 (4)	0.0019 (4)
O1	0.0459 (14)	0.0648 (17)	0.0443 (16)	0.0157 (13)	0.0152 (12)	0.0131 (13)
O2	0.0455 (14)	0.0413 (14)	0.0633 (18)	-0.0079 (11)	-0.0011 (13)	0.0018 (13)
N1	0.0367 (14)	0.0292 (14)	0.0336 (16)	0.0000 (12)	0.0023 (12)	0.0011 (12)
N2	0.0332 (14)	0.0297 (15)	0.0358 (16)	0.0036 (11)	0.0059 (12)	0.0022 (12)
C1	0.052 (2)	0.037 (2)	0.059 (3)	0.0101 (17)	-0.019 (2)	-0.0037 (19)
C2	0.058 (2)	0.035 (2)	0.039 (2)	0.0084 (17)	-0.0113 (18)	0.0016 (17)
C3	0.0396 (18)	0.0274 (16)	0.038 (2)	-0.0049 (14)	0.0082 (15)	-0.0018 (15)
C4	0.0458 (19)	0.0307 (18)	0.041 (2)	-0.0024 (15)	0.0071 (16)	-0.0022 (16)
C5	0.060 (2)	0.0301 (19)	0.044 (2)	0.0053 (16)	0.0098 (18)	0.0000 (17)
C6	0.0390 (18)	0.0301 (18)	0.037 (2)	-0.0041 (14)	0.0046 (15)	-0.0051 (15)
C7	0.041 (2)	0.056 (2)	0.065 (3)	-0.0041 (18)	0.0109 (19)	0.006 (2)

C8	0.058 (2)	0.050 (2)	0.045 (2)	0.0027 (19)	-0.0075 (19)	-0.0090 (19)
C9	0.060 (2)	0.075 (3)	0.056 (3)	0.025 (2)	0.020 (2)	0.014 (2)
C10	0.044 (2)	0.061 (3)	0.050 (3)	0.019 (2)	0.0147 (18)	0.011 (2)
C11	0.058 (3)	0.073 (3)	0.060 (3)	0.019 (2)	0.012 (2)	-0.001 (2)
C12	0.083 (4)	0.128 (5)	0.055 (3)	0.050 (4)	0.018 (3)	-0.003 (3)
C13	0.097 (4)	0.138 (6)	0.091 (5)	0.063 (4)	0.061 (4)	0.053 (5)
C14	0.114 (5)	0.089 (5)	0.146 (6)	0.021 (4)	0.080 (5)	0.043 (5)
C15	0.095 (4)	0.077 (4)	0.086 (4)	-0.002 (3)	0.043 (3)	0.001 (3)
C16	0.055 (2)	0.039 (2)	0.081 (3)	-0.0046 (18)	0.004 (2)	0.005 (2)
C17	0.043 (2)	0.038 (2)	0.059 (3)	-0.0055 (17)	0.0095 (18)	0.0008 (19)
C18	0.060 (3)	0.079 (3)	0.064 (3)	0.020 (2)	0.010 (2)	-0.016 (3)
C19	0.043 (2)	0.118 (5)	0.076 (4)	0.011 (3)	0.009 (2)	0.010 (3)
C20	0.066 (3)	0.060 (3)	0.114 (5)	-0.018 (3)	-0.014 (3)	0.002 (3)
C21	0.113 (5)	0.062 (3)	0.126 (5)	0.007 (3)	-0.023 (4)	-0.044 (4)
C22	0.064 (3)	0.062 (3)	0.109 (4)	0.012 (2)	0.006 (3)	-0.021 (3)

Geometric parameters (Å, °)

Ni1—N2	1.958 (3)	C7—H7C	0.9600
Ni1—N1	1.963 (3)	C8—H8A	0.9600
S1—P1	1.9674 (14)	C8—H8B	0.9600
S2—P1	1.9661 (15)	C8—H8C	0.9600
P1—O1	1.608 (3)	C9—C10	1.490 (6)
P1—O2	1.613 (3)	C9—H9A	0.9700
O1—C9	1.459 (4)	C9—H9B	0.9700
O2—C16	1.432 (4)	C10—C15	1.374 (7)
N1—C1	1.482 (4)	C10—C11	1.385 (6)
N1—C6	1.509 (4)	C11—C12	1.400 (7)
N1—H1	0.9100	C11—H11	0.9300
N2—C2	1.493 (4)	C12—C13	1.366 (9)
N2—C3	1.501 (4)	C12—H12	0.9300
N2—H2	0.9100	C13—C14	1.354 (9)
C1—C2	1.488 (5)	C13—H13	0.9300
C1—H1A	0.9700	C14—C15	1.363 (8)
C1—H1B	0.9700	C14—H14	0.9300
C2—H2A	0.9700	C15—H15	0.9300
C2—H2B	0.9700	C16—C17	1.500 (5)
C3—C4	1.510 (4)	C16—H16A	0.9700
C3—C5	1.535 (4)	C16—H16B	0.9700
C3—H3	0.9800	C17—C18	1.369 (5)
C4—C6 ⁱ	1.520 (4)	C17—C22	1.370 (6)
C4—H4A	0.9700	C18—C19	1.388 (6)
C4—H4B	0.9700	C18—H18	0.9300
C5—H5A	0.9600	C19—C20	1.332 (7)
C5—H5B	0.9600	C19—H19	0.9300
C5—H5C	0.9600	C20—C21	1.366 (7)
C6—C4 ⁱ	1.520 (4)	C20—H20	0.9300
C6—C7	1.528 (5)	C21—C22	1.376 (6)
C6—C8	1.541 (5)	C21—H21	0.9300

supplementary materials

C7—H7A	0.9600	C22—H22	0.9300
C7—H7B	0.9600		
N2—Ni1—N1	86.71 (11)	C6—C7—H7B	109.5
N2 ⁱ —Ni1—N1	93.29 (11)	H7A—C7—H7B	109.5
O1—P1—O2	104.07 (15)	C6—C7—H7C	109.5
O1—P1—S2	111.40 (12)	H7A—C7—H7C	109.5
O2—P1—S2	103.65 (11)	H7B—C7—H7C	109.5
O1—P1—S1	105.11 (11)	C6—C8—H8A	109.5
O2—P1—S1	111.05 (11)	C6—C8—H8B	109.5
S2—P1—S1	120.50 (7)	H8A—C8—H8B	109.5
C9—O1—P1	119.1 (2)	C6—C8—H8C	109.5
C16—O2—P1	120.7 (2)	H8A—C8—H8C	109.5
C1—N1—C6	112.0 (2)	H8B—C8—H8C	109.5
C1—N1—Ni1	108.7 (2)	O1—C9—C10	108.6 (3)
C6—N1—Ni1	122.9 (2)	O1—C9—H9A	110.0
C1—N1—H1	103.6	C10—C9—H9A	110.0
C6—N1—H1	103.6	O1—C9—H9B	110.0
Ni1—N1—H1	103.6	C10—C9—H9B	110.0
C2—N2—C3	110.1 (2)	H9A—C9—H9B	108.3
C2—N2—Ni1	107.3 (2)	C15—C10—C11	118.4 (4)
C3—N2—Ni1	121.11 (19)	C15—C10—C9	120.1 (4)
C2—N2—H2	105.8	C11—C10—C9	121.5 (4)
C3—N2—H2	105.8	C10—C11—C12	119.2 (5)
Ni1—N2—H2	105.8	C10—C11—H11	120.4
N1—C1—C2	108.0 (3)	C12—C11—H11	120.4
N1—C1—H1A	110.1	C13—C12—C11	120.1 (6)
C2—C1—H1A	110.1	C13—C12—H12	119.9
N1—C1—H1B	110.1	C11—C12—H12	119.9
C2—C1—H1B	110.1	C14—C13—C12	120.6 (6)
H1A—C1—H1B	108.4	C14—C13—H13	119.7
C1—C2—N2	107.9 (3)	C12—C13—H13	119.7
C1—C2—H2A	110.1	C13—C14—C15	119.5 (6)
N2—C2—H2A	110.1	C13—C14—H14	120.2
C1—C2—H2B	110.1	C15—C14—H14	120.2
N2—C2—H2B	110.1	C14—C15—C10	122.2 (6)
H2A—C2—H2B	108.4	C14—C15—H15	118.9
N2—C3—C4	110.0 (3)	C10—C15—H15	118.9
N2—C3—C5	112.4 (2)	O2—C16—C17	108.3 (3)
C4—C3—C5	110.3 (3)	O2—C16—H16A	110.0
N2—C3—H3	108.0	C17—C16—H16A	110.0
C4—C3—H3	108.0	O2—C16—H16B	110.0
C5—C3—H3	108.0	C17—C16—H16B	110.0
C3—C4—C6 ⁱ	117.4 (3)	H16A—C16—H16B	108.4
C3—C4—H4A	107.9	C18—C17—C22	118.0 (4)
C6 ⁱ —C4—H4A	107.9	C18—C17—C16	121.3 (4)
C3—C4—H4B	107.9	C22—C17—C16	120.6 (4)
C6 ⁱ —C4—H4B	107.9	C17—C18—C19	121.1 (4)
H4A—C4—H4B	107.2	C17—C18—H18	119.4

C3—C5—H5A	109.5	C19—C18—H18	119.4
C3—C5—H5B	109.5	C20—C19—C18	119.8 (5)
H5A—C5—H5B	109.5	C20—C19—H19	120.1
C3—C5—H5C	109.5	C18—C19—H19	120.1
H5A—C5—H5C	109.5	C19—C20—C21	120.3 (5)
H5B—C5—H5C	109.5	C19—C20—H20	119.8
N1—C6—C4 ⁱ	108.0 (2)	C21—C20—H20	119.8
N1—C6—C7	109.6 (3)	C20—C21—C22	120.1 (5)
C4 ⁱ —C6—C7	111.1 (3)	C20—C21—H21	120.0
N1—C6—C8	109.6 (3)	C22—C21—H21	120.0
C4 ⁱ —C6—C8	108.3 (3)	C17—C22—C21	120.6 (5)
C7—C6—C8	110.3 (3)	C17—C22—H22	119.7
C6—C7—H7A	109.5	C21—C22—H22	119.7

Symmetry codes: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots S1	0.91	2.61	3.390 (3)	144
N2—H2 \cdots S2 ⁱ	0.91	2.50	3.394 (3)	169

Symmetry codes: (i) $-x+1, -y+1, -z$.

Fig. 1

