

Tris[tris(1,10-phenanthroline- κ^2N,N')-iron(II)] dodecatungstoferrate dihydrate

 Feng-Xia Ma,^{a,b} Ya-Guang Chen^{a*} and Dong-Mei Shi^a

^aDepartment of Chemistry, Northeast Normal University, Changchun 130024, People's Republic of China, and ^bBasic Science Department, Jilin Agricultural Science and Technology College, Jilin 132101, People's Republic of China

Correspondence e-mail: chenyg146@nenu.edu.cn


Received 24 March 2008; accepted 10 April 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.027$ Å; H-atom completeness 95%; disorder in solvent or counterion; R factor = 0.036; wR factor = 0.094; data-to-parameter ratio = 11.0.

The title compound, $[\text{Fe}(\text{C}_{12}\text{H}_8\text{N}_2)_3]_3[\text{FeW}_{12}\text{O}_{40}] \cdot 2\text{H}_2\text{O}$, was prepared under hydrothermal conditions. The discrete Keggin-type $[\text{FeW}_{12}\text{O}_{40}]^{6-}$ heteropolyoxoanion has threefold symmetry, with the Fe^{II} atom located on the threefold rotation axis. The central FeO_4 tetrahedron in the anion shares its O atoms with four W_3O_{13} trinuclear units, each of which is made up of three edge-shared WO_6 octahedral units. The Fe^{II} atom in the complex cation, *viz* $[\text{Fe}(\text{phen})_3]^{2+}$ (phen is 1,10-phenanthroline), shows a slightly distorted octahedral geometry defined by six N atoms from three phen ligands. The polyoxoanions pack together with the cations, with the disordered water molecules located in voids; the site occupancy factor for each water O atom is 0.33.

Related literature

For related literature, see: Brown (2002); Misono (1987); Pope (1983).



Experimental

Crystal data

$[\text{Fe}(\text{C}_{12}\text{H}_8\text{N}_2)_3]_3[\text{FeW}_{12}\text{O}_{40}] \cdot 2\text{H}_2\text{O}$
 $M_r = 4727.47$

Trigonal, $R\bar{3}$
 $a = 25.088$ (5) Å

$c = 17.231$ (5) Å
 $V = 9392$ (4) Å³
 $Z = 3$
 Mo $K\alpha$ radiation

$\mu = 11.51$ mm⁻¹
 $T = 293$ (2) K
 $0.24 \times 0.21 \times 0.20$ mm

Data collection

Bruker SMART APEX CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (*SADABS*; Sheldrick, 1996)
 $T_{\text{min}} = 0.075$, $T_{\text{max}} = 0.104$

16084 measured reflections
 6154 independent reflections
 5557 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.093$
 $S = 1.03$
 6154 reflections
 560 parameters
 1 restraint

H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.62$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.55$ e Å⁻³
 Absolute structure: Flack (1983),
 3713 Friedel pairs
 Flack parameter: -0.006 (6)

Table 1
 Selected bond lengths (Å).

Fe1—O13	1.826 (8)	W2—O14 ⁱ	1.954 (9)
Fe1—O1	1.833 (18)	W2—O11 ⁱ	1.963 (10)
Fe2—N6	1.969 (11)	W2—O13 ⁱ	2.211 (8)
Fe2—N3	1.971 (12)	W3—O6	1.742 (11)
Fe2—N1	1.972 (10)	W3—O2 ⁱ	1.903 (9)
Fe2—N5	1.979 (12)	W3—O2	1.911 (9)
Fe2—N2	1.986 (11)	W3—O11	1.951 (10)
Fe2—N4	2.001 (11)	W3—O10	1.961 (10)
W1—O3	1.692 (9)	W3—O13	2.229 (10)
W1—O4	1.898 (9)	W4—O8	1.678 (9)
W1—O7	1.943 (8)	W4—O9	1.896 (8)
W1—O12	1.950 (8)	W4—O4	1.904 (9)
W1—O1	2.230 (9)	W4—O10	1.944 (11)
W2—O5	1.702 (10)	W4—O14	2.001 (9)
W2—O7	1.858 (9)	W4—O13	2.221 (8)
W2—O9	1.940 (9)		

Symmetry code: (i) $-x + y + 1, -x + 1, z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Analysis and Testing Foundation of Northeast Normal University, Changchun, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2126).

References

- Brown, I. D. (2002). *The Chemical Bond in Inorganic Chemistry: the Bond Valence Model*. Oxford University Press.
 Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Misono, M. (1987). *Cat. Rev. Sci. Eng.* **29**, 269–321.
 Pope, M. T. (1983). *Heteropoly and Isopoly Oxometalates*. Berlin: Springer.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m672 [doi:10.1107/S1600536808009896]

Tris[tris(1,10-phenanthroline- κ^2N,N')iron(II)] dodecatungstoferrate dihydrate

F.-X. Ma, Y.-G. Chen and D.-M. Shi

Comment

Polyoxometalates (POMs) have attracted attention in recent years, not only because of their structural diversity but also because of their potential applications in medicine, material science, especially in catalysis (Misono, 1987; Pope, 1983).

The structure of the title compound is built up from three complex cations, $[\text{Fe}(\text{phen})_3]^{2+}$ (phen = 1,10-phenanthroline), one Keggin-type anion, $[\text{FeW}_{12}\text{O}_{40}]^{6-}$, and two disordered water molecules. The heteropolyoxoanion $[\text{FeW}_{12}\text{O}_{40}]^{6-}$ has threefold symmetry with the Fe1 atom located on the threefold rotation axis. The Fe2 atom in the cation is coordinated by six N atoms from three phen molecules, forming a distorted FeN_6 octahedron with Fe—N bond distances ranging from 1.969 (11) to 2.001 (11) Å (Fig. 1; Table 1).

Results of bond valence sum (BVS) calculations (Brown, 2002) are in accordance with expected values for hexavalent tungsten (average 6.15 valence units for the 12 W atoms) and divalent iron (1.92 valence units for the Fe atom in the complex cation).

Experimental

A mixture of ammonium tungstate monohydrate (1.005 g, 0.6 mmol), $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (0.157 g, 0.6 mmol), phen. H_2O (0.256 g, 1.3 mmol), NH_4VO_3 (0.053 g, 0.5 mmol), oxalic acid dihydrate (0.205 g, 1.6 mmol) and H_2O (10 ml) was adjusted to pH = 5.8 by addition of 2 mol L^{-1} NaOH under stirring for 30 min. The final solution was transferred into a 25 ml Teflon-lined autoclave and was heated at 453 K for 96 h. Then the autoclave was cooled in a rate of 10 K h^{-1} to room temperature. Red block-like crystals were filtered off, washed with distilled water, and dried at ambient temperature (40% yield on W).

Refinement

H atoms on C atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. H atoms of the disordered water molecules were not located. In the final difference Fourier map, the highest peak is 2.83 Å away from O2W and the deepest hole is 0.23 Å from Fe1.

Figures




Fig. 1. The structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms and disordered water molecules are omitted for clarity. [Symmetry codes: (i) $-y + 1, x - y, z$; (ii) $-x + y + 1, -x + 1, z$.]

Tris[tris(1,10-phenanthroline- κ^2N,N')iron(II)] dodecatungstoferrate dihydrate

Crystal data

$[\text{Fe}(\text{C}_{12}\text{H}_8\text{N}_2)_3]_3[\text{FeW}_{12}\text{O}_{40}] \cdot 2\text{H}_2\text{O}$

$M_r = 4727.47$

Trigonal, $R\bar{3}$

Hall symbol: $R\bar{3}$

$a = 25.088 (5) \text{ \AA}$

$b = 25.088 (5) \text{ \AA}$

$c = 17.231 (5) \text{ \AA}$

$\alpha = 90^\circ$

$\beta = 90^\circ$

$\gamma = 120^\circ$

$V = 9392 (4) \text{ \AA}^3$

$Z = 3$

$F_{000} = 6593.7$

$D_x = 2.527 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5571 reflections

$\theta = 1.5\text{--}25.1^\circ$

$\mu = 11.51 \text{ mm}^{-1}$

$T = 293 (2) \text{ K}$

Block, red

$0.24 \times 0.21 \times 0.20 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293(2) \text{ K}$

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.075, T_{\max} = 0.104$

16084 measured reflections

6154 independent reflections

5557 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

$\theta_{\max} = 25.1^\circ$

$\theta_{\min} = 1.5^\circ$

$h = -29 \rightarrow 29$

$k = -29 \rightarrow 29$

$l = -20 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0566P)^2 + 7.5378P]$

$wR(F^2) = 0.093$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\max} = 0.001$
6154 reflections	$\Delta\rho_{\max} = 1.62 \text{ e } \text{\AA}^{-3}$
560 parameters	$\Delta\rho_{\min} = -1.55 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 3713 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: $-0.006 (6)$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Fe1	0.6667	0.3333	0.3946 (2)	0.0268 (6)	
Fe2	0.32586 (8)	0.15648 (8)	0.72949 (11)	0.0297 (4)	
W1	0.58806 (3)	0.33016 (3)	0.22372 (4)	0.03811 (16)	
W2	0.58842 (3)	0.41729 (3)	0.39586 (3)	0.04278 (17)	
W3	0.58253 (3)	0.24702 (3)	0.55437 (4)	0.04819 (18)	
W4	0.50462 (2)	0.24669 (3)	0.39934 (3)	0.04051 (16)	
C1	0.2245 (6)	0.0581 (6)	0.8268 (8)	0.035 (3)	
H1A	0.2201	0.0887	0.8513	0.043*	
C2	0.1869 (6)	-0.0018 (6)	0.8499 (9)	0.040 (3)	
H2A	0.1596	-0.0099	0.8907	0.048*	
C3	0.1885 (6)	-0.0479 (6)	0.8161 (8)	0.042 (3)	
H3A	0.1614	-0.0885	0.8298	0.050*	
C4	0.2355 (6)	-0.0320 (5)	0.7551 (8)	0.035 (3)	
C5	0.2712 (5)	0.0293 (5)	0.7356 (8)	0.035 (3)	
C6	0.3150 (6)	0.0474 (6)	0.6755 (8)	0.044 (3)	
C7	0.3259 (7)	0.0014 (6)	0.6414 (9)	0.052 (4)	
C8	0.2083 (6)	0.1176 (7)	0.6442 (9)	0.048 (4)	
H8A	0.1871	0.0856	0.6792	0.057*	
C9	0.1760 (9)	0.1237 (9)	0.5801 (13)	0.079 (7)	
H9A	0.1338	0.0965	0.5779	0.095*	
C10	0.1990 (11)	0.1623 (11)	0.5263 (12)	0.080 (6)	
H10A	0.1757	0.1626	0.4846	0.096*	
C11	0.3893 (7)	0.1268 (7)	0.6061 (10)	0.062 (5)	
H11A	0.4110	0.1683	0.5931	0.074*	
C12	0.4027 (12)	0.0851 (9)	0.5665 (12)	0.095 (8)	
H12A	0.4323	0.1003	0.5275	0.114*	
C13	0.2444 (7)	-0.0784 (7)	0.7168 (11)	0.063 (5)	
H13A	0.2185	-0.1201	0.7264	0.076*	
C14	0.4425 (7)	0.2691 (7)	0.6851 (10)	0.054 (4)	
H14A	0.4602	0.2610	0.7277	0.065*	
C15	0.2629 (13)	0.2054 (10)	0.5327 (10)	0.085 (7)	
C16	0.3002 (18)	0.2552 (15)	0.4770 (12)	0.140 (14)	
H16A	0.2817	0.2599	0.4327	0.168*	
C17	0.3649 (14)	0.2965 (12)	0.4905 (13)	0.094 (8)	
H17A	0.3881	0.3268	0.4542	0.113*	

supplementary materials

C18	0.3929 (12)	0.2912 (8)	0.5584 (11)	0.081 (7)
C19	0.3567 (8)	0.2426 (6)	0.6103 (9)	0.051 (4)
C20	0.3430 (5)	0.2139 (5)	0.8739 (7)	0.033 (3)
C21	0.3882 (6)	0.1979 (6)	0.8721 (7)	0.038 (3)
C22	0.4287 (7)	0.2118 (7)	0.9329 (9)	0.054 (4)
C23	0.4718 (8)	0.1912 (9)	0.9277 (12)	0.070 (6)
H23A	0.5000	0.1999	0.9675	0.084*
C24	0.4291 (8)	0.1453 (9)	0.8025 (11)	0.072 (5)
H24A	0.4292	0.1232	0.7592	0.086*
C25	0.2933 (11)	-0.0580 (8)	0.6650 (11)	0.081 (7)
H25A	0.3040	-0.0861	0.6460	0.097*
C26	0.2600 (6)	0.2131 (6)	0.8112 (9)	0.042 (3)
H26A	0.2338	0.2023	0.7687	0.051*
C27	0.2488 (7)	0.2413 (7)	0.8753 (9)	0.045 (4)
H27A	0.2156	0.2482	0.8747	0.054*
C28	0.2852 (8)	0.2575 (7)	0.9352 (9)	0.059 (5)
H28A	0.2784	0.2772	0.9765	0.071*
C29	0.3371 (7)	0.2455 (6)	0.9392 (8)	0.045 (3)
C30	0.3740 (9)	0.2556 (8)	0.9998 (10)	0.068 (5)
H30A	0.3661	0.2705	1.0452	0.081*
C31	0.4232 (8)	0.2451 (7)	0.9995 (9)	0.063 (5)
H31A	0.4516	0.2587	1.0399	0.076*
C32	0.2986 (8)	0.2029 (6)	0.5974 (9)	0.054 (4)
C33	0.4560 (12)	0.3308 (9)	0.5740 (13)	0.088 (8)
H33A	0.4795	0.3637	0.5411	0.105*
C34	0.3739 (9)	0.0237 (9)	0.5835 (11)	0.083 (6)
H34A	0.3847	-0.0025	0.5589	0.100*
C35	0.4820 (8)	0.3208 (7)	0.6360 (13)	0.070 (6)
H35A	0.5236	0.3461	0.6471	0.084*
C36	0.4723 (8)	0.1583 (10)	0.8642 (13)	0.084 (6)
H36A	0.5005	0.1446	0.8613	0.101*
N1	0.2665 (4)	0.0749 (4)	0.7718 (6)	0.027 (2)
N2	0.3485 (5)	0.1086 (5)	0.6585 (7)	0.040 (3)
N3	0.2656 (6)	0.1551 (5)	0.6549 (6)	0.040 (3)
N4	0.3847 (6)	0.2337 (5)	0.6735 (7)	0.041 (3)
N5	0.3875 (5)	0.1656 (5)	0.8079 (7)	0.040 (3)
N6	0.3062 (5)	0.2008 (5)	0.8083 (6)	0.031 (2)
O1	0.6667	0.3333	0.2882 (10)	0.038 (4)
O2	0.6684 (4)	0.2746 (4)	0.5635 (6)	0.047 (2)
O3	0.5416 (5)	0.3302 (5)	0.1526 (6)	0.046 (2)
O4	0.5342 (4)	0.2705 (4)	0.2961 (5)	0.040 (2)
O5	0.5375 (5)	0.4431 (5)	0.3877 (7)	0.059 (3)
O6	0.5548 (5)	0.2172 (6)	0.6463 (6)	0.063 (3)
O7	0.5918 (4)	0.3915 (4)	0.2958 (5)	0.037 (2)
O8	0.4282 (4)	0.2170 (5)	0.3920 (7)	0.056 (3)
O9	0.5345 (4)	0.3311 (4)	0.4206 (5)	0.040 (2)
O10	0.5036 (4)	0.2295 (4)	0.5095 (6)	0.049 (3)
O11	0.5687 (4)	0.1702 (4)	0.5081 (6)	0.046 (2)
O12	0.6053 (4)	0.2694 (4)	0.1776 (6)	0.040 (2)

O13	0.5982 (4)	0.2633 (4)	0.4271 (6)	0.036 (2)	
O14	0.5038 (4)	0.1676 (4)	0.3806 (6)	0.045 (2)	
O1W	0.5848 (13)	0.3187 (15)	0.7626 (19)	0.053 (9)	0.33
O2W	0.5964 (9)	0.2622 (9)	1.0192 (12)	0.013 (4)*	0.33

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0211 (7)	0.0211 (7)	0.0382 (16)	0.0105 (4)	0.000	0.000
Fe2	0.0294 (9)	0.0271 (10)	0.0303 (9)	0.0123 (8)	0.0027 (7)	-0.0021 (8)
W1	0.0322 (3)	0.0341 (3)	0.0487 (4)	0.0171 (2)	-0.0097 (3)	-0.0013 (3)
W2	0.0389 (3)	0.0400 (3)	0.0606 (5)	0.0282 (3)	-0.0024 (3)	-0.0043 (3)
W3	0.0459 (4)	0.0465 (4)	0.0512 (4)	0.0223 (3)	0.0143 (3)	0.0129 (3)
W4	0.0216 (3)	0.0361 (3)	0.0588 (4)	0.0107 (2)	0.0040 (2)	-0.0005 (3)
C1	0.038 (7)	0.033 (7)	0.039 (8)	0.020 (6)	0.012 (6)	0.008 (6)
C2	0.042 (8)	0.042 (8)	0.042 (9)	0.025 (7)	0.007 (6)	0.009 (7)
C3	0.038 (7)	0.035 (7)	0.055 (9)	0.020 (6)	-0.008 (6)	0.006 (6)
C4	0.039 (7)	0.020 (6)	0.047 (8)	0.016 (5)	0.005 (6)	0.000 (5)
C5	0.029 (6)	0.028 (7)	0.050 (8)	0.016 (5)	-0.003 (6)	0.001 (6)
C6	0.052 (8)	0.047 (8)	0.037 (8)	0.029 (7)	0.004 (7)	-0.020 (6)
C7	0.060 (9)	0.040 (8)	0.051 (10)	0.020 (7)	0.014 (7)	-0.008 (7)
C8	0.038 (8)	0.040 (7)	0.058 (10)	0.013 (6)	-0.024 (7)	-0.011 (6)
C9	0.082 (13)	0.064 (12)	0.106 (17)	0.048 (11)	-0.068 (13)	-0.046 (12)
C10	0.094 (16)	0.082 (15)	0.067 (14)	0.046 (13)	-0.038 (12)	-0.005 (11)
C11	0.061 (10)	0.033 (8)	0.082 (13)	0.017 (7)	0.034 (9)	0.010 (8)
C12	0.15 (2)	0.066 (13)	0.075 (15)	0.056 (14)	0.066 (15)	-0.001 (10)
C13	0.052 (10)	0.042 (9)	0.083 (13)	0.014 (8)	0.012 (9)	-0.012 (8)
C14	0.050 (9)	0.045 (9)	0.055 (10)	0.014 (8)	0.016 (8)	-0.010 (7)
C15	0.18 (2)	0.085 (14)	0.035 (10)	0.097 (17)	0.013 (12)	0.019 (9)
C16	0.31 (4)	0.18 (3)	0.044 (14)	0.21 (3)	0.06 (2)	0.056 (17)
C17	0.16 (2)	0.101 (19)	0.069 (16)	0.102 (19)	0.033 (16)	0.041 (13)
C18	0.15 (2)	0.056 (11)	0.056 (12)	0.067 (13)	0.049 (12)	0.015 (9)
C19	0.081 (12)	0.036 (8)	0.042 (9)	0.033 (8)	0.035 (8)	0.013 (7)
C20	0.028 (6)	0.023 (6)	0.035 (7)	0.003 (5)	0.005 (5)	0.006 (5)
C21	0.033 (7)	0.047 (8)	0.030 (7)	0.017 (6)	-0.010 (6)	-0.002 (6)
C22	0.043 (8)	0.045 (9)	0.058 (11)	0.009 (7)	0.001 (7)	0.010 (7)
C23	0.037 (9)	0.085 (13)	0.074 (14)	0.020 (9)	-0.029 (9)	0.010 (11)
C24	0.046 (9)	0.089 (13)	0.088 (14)	0.040 (10)	-0.014 (9)	-0.009 (11)
C25	0.139 (18)	0.048 (10)	0.075 (13)	0.062 (12)	0.032 (13)	-0.011 (9)
C26	0.030 (7)	0.035 (7)	0.056 (10)	0.012 (6)	-0.005 (6)	0.001 (6)
C27	0.055 (9)	0.055 (9)	0.037 (9)	0.035 (8)	-0.001 (7)	-0.006 (7)
C28	0.093 (13)	0.050 (9)	0.039 (10)	0.040 (9)	0.025 (9)	-0.008 (7)
C29	0.052 (9)	0.043 (8)	0.033 (8)	0.019 (7)	-0.011 (7)	-0.018 (6)
C30	0.085 (13)	0.059 (11)	0.059 (12)	0.036 (10)	-0.018 (10)	-0.021 (8)
C31	0.073 (12)	0.040 (8)	0.046 (10)	0.005 (8)	-0.026 (9)	-0.006 (7)
C32	0.090 (12)	0.030 (8)	0.047 (10)	0.034 (8)	0.019 (8)	0.007 (7)
C33	0.117 (19)	0.051 (11)	0.077 (16)	0.029 (13)	0.071 (14)	0.014 (11)
C34	0.083 (14)	0.073 (13)	0.095 (15)	0.039 (11)	0.026 (12)	-0.030 (11)

supplementary materials

C35	0.046 (10)	0.035 (9)	0.101 (17)	0.000 (8)	0.047 (11)	0.002 (9)
C36	0.050 (11)	0.116 (17)	0.101 (18)	0.052 (12)	-0.006 (10)	0.006 (14)
N1	0.031 (5)	0.027 (5)	0.028 (6)	0.019 (5)	0.001 (4)	0.000 (4)
N2	0.041 (7)	0.039 (6)	0.037 (7)	0.019 (5)	0.009 (5)	-0.004 (5)
N3	0.061 (8)	0.039 (6)	0.028 (6)	0.031 (6)	-0.001 (6)	-0.006 (5)
N4	0.063 (8)	0.022 (5)	0.031 (7)	0.016 (6)	0.023 (6)	0.000 (5)
N5	0.023 (5)	0.037 (6)	0.049 (7)	0.008 (5)	0.004 (5)	-0.001 (5)
N6	0.030 (6)	0.026 (5)	0.024 (6)	0.006 (5)	-0.002 (4)	-0.003 (4)
O1	0.030 (5)	0.030 (5)	0.055 (11)	0.015 (2)	0.000	0.000
O2	0.046 (6)	0.043 (6)	0.054 (7)	0.025 (5)	0.003 (5)	0.008 (5)
O3	0.048 (6)	0.048 (6)	0.048 (6)	0.028 (5)	-0.022 (5)	-0.008 (5)
O4	0.040 (5)	0.028 (5)	0.049 (6)	0.013 (4)	0.005 (4)	-0.003 (4)
O5	0.054 (6)	0.038 (6)	0.096 (9)	0.030 (5)	-0.002 (6)	-0.010 (5)
O6	0.062 (7)	0.072 (8)	0.056 (8)	0.035 (6)	0.016 (6)	0.015 (6)
O7	0.041 (5)	0.028 (5)	0.050 (6)	0.022 (4)	-0.007 (4)	-0.009 (4)
O8	0.025 (5)	0.045 (6)	0.089 (9)	0.010 (4)	0.013 (5)	-0.003 (6)
O9	0.040 (5)	0.032 (5)	0.053 (6)	0.021 (4)	0.006 (4)	-0.002 (4)
O10	0.049 (6)	0.033 (5)	0.054 (7)	0.013 (5)	0.020 (5)	-0.004 (4)
O11	0.046 (6)	0.041 (5)	0.054 (7)	0.024 (5)	0.005 (5)	0.015 (5)
O12	0.034 (5)	0.032 (5)	0.054 (7)	0.016 (4)	0.000 (4)	0.001 (4)
O13	0.036 (5)	0.026 (4)	0.044 (6)	0.013 (4)	0.002 (4)	0.002 (4)
O14	0.039 (5)	0.041 (5)	0.058 (7)	0.022 (5)	0.003 (4)	-0.005 (4)
O1W	0.048 (18)	0.08 (2)	0.05 (2)	0.041 (17)	-0.010 (14)	-0.042 (16)

Geometric parameters (Å, °)

Fe1—O13 ⁱ	1.826 (8)	C11—C12	1.42 (2)
Fe1—O13	1.826 (8)	C11—H11A	0.9300
Fe1—O13 ⁱⁱ	1.826 (8)	C12—C34	1.37 (2)
Fe1—O1	1.833 (18)	C12—H12A	0.9300
Fe2—N6	1.969 (11)	C13—C25	1.39 (2)
Fe2—N3	1.971 (12)	C13—H13A	0.9300
Fe2—N1	1.972 (10)	C14—N4	1.283 (19)
Fe2—N5	1.979 (12)	C14—C35	1.45 (2)
Fe2—N2	1.986 (11)	C14—H14A	0.9300
Fe2—N4	2.001 (11)	C15—C32	1.45 (3)
W1—O3	1.692 (9)	C15—C16	1.48 (3)
W1—O4	1.898 (9)	C16—C17	1.44 (4)
W1—O7	1.943 (8)	C16—H16A	0.9300
W1—O12 ⁱⁱ	1.950 (9)	C17—C18	1.41 (3)
W1—O12	1.950 (8)	C17—H17A	0.9300
W1—O1	2.230 (9)	C18—C33	1.41 (3)
W2—O5	1.702 (10)	C18—C19	1.42 (2)
W2—O7	1.858 (9)	C19—C32	1.31 (2)
W2—O9	1.940 (9)	C19—N4	1.37 (2)
W2—O14 ⁱⁱ	1.954 (9)	C20—C21	1.380 (17)
W2—O11 ⁱⁱ	1.963 (10)	C20—N6	1.390 (16)
W2—O13 ⁱⁱ	2.211 (8)	C20—C29	1.425 (18)

W3—O6	1.742 (11)	C21—N5	1.367 (17)
W3—O2 ⁱⁱ	1.903 (9)	C21—C22	1.377 (19)
W3—O2	1.911 (9)	C22—C23	1.42 (2)
W3—O11	1.951 (10)	C22—C31	1.47 (2)
W3—O10	1.961 (10)	C23—C36	1.37 (3)
W3—O13	2.229 (10)	C23—H23A	0.9300
W4—O8	1.678 (9)	C24—N5	1.373 (19)
W4—O9	1.896 (8)	C24—C36	1.44 (2)
W4—O4	1.904 (9)	C24—H24A	0.9300
W4—O10	1.944 (11)	C25—H25A	0.9300
W4—O14	2.001 (9)	C26—N6	1.341 (16)
W4—O13	2.221 (8)	C26—C27	1.41 (2)
C1—N1	1.319 (15)	C26—H26A	0.9300
C1—C2	1.375 (18)	C27—C28	1.30 (2)
C1—H1A	0.9300	C27—H27A	0.9300
C2—C3	1.313 (18)	C28—C29	1.48 (2)
C2—H2A	0.9300	C28—H28A	0.9300
C3—C4	1.477 (18)	C29—C30	1.33 (2)
C3—H3A	0.9300	C30—C31	1.39 (2)
C4—C5	1.378 (16)	C30—H30A	0.9300
C4—C13	1.451 (18)	C31—H31A	0.9300
C5—N1	1.360 (15)	C32—N3	1.455 (18)
C5—C6	1.409 (18)	C33—C35	1.34 (3)
C6—N2	1.363 (17)	C33—H33A	0.9300
C6—C7	1.440 (18)	C34—H34A	0.9300
C7—C25	1.36 (2)	C35—H35A	0.9300
C7—C34	1.44 (2)	C36—H36A	0.9300
C8—N3	1.279 (17)	O1—W1 ⁱ	2.230 (9)
C8—C9	1.42 (2)	O1—W1 ⁱⁱ	2.230 (9)
C8—H8A	0.9300	O2—W3 ⁱ	1.903 (9)
C9—C10	1.25 (3)	O11—W2 ⁱ	1.963 (10)
C9—H9A	0.9300	O12—W1 ⁱ	1.950 (8)
C10—C15	1.42 (3)	O13—W2 ⁱ	2.211 (8)
C10—H10A	0.9300	O14—W2 ⁱ	1.954 (9)
C11—N2	1.267 (17)		
O13 ⁱ —Fe1—O13	111.0 (3)	C34—C12—C11	122.6 (17)
O13 ⁱ —Fe1—O13 ⁱⁱ	111.0 (3)	C34—C12—H12A	118.7
O13—Fe1—O13 ⁱⁱ	111.0 (3)	C11—C12—H12A	118.7
O13 ⁱ —Fe1—O1	107.9 (3)	C25—C13—C4	117.4 (14)
O13—Fe1—O1	107.9 (3)	C25—C13—H13A	121.3
O13 ⁱⁱ —Fe1—O1	107.9 (3)	C4—C13—H13A	121.3
N6—Fe2—N3	93.0 (4)	N4—C14—C35	124.1 (18)
N6—Fe2—N1	93.4 (4)	N4—C14—H14A	118.0
N3—Fe2—N1	92.8 (4)	C35—C14—H14A	118.0
N6—Fe2—N5	82.3 (5)	C10—C15—C32	120.9 (16)
N3—Fe2—N5	174.4 (5)	C10—C15—C16	126 (2)

supplementary materials

N1—Fe2—N5	90.5 (4)	C32—C15—C16	113 (2)
N6—Fe2—N2	174.4 (5)	C17—C16—C15	121 (2)
N3—Fe2—N2	92.1 (5)	C17—C16—H16A	119.7
N1—Fe2—N2	84.0 (4)	C15—C16—H16A	119.7
N5—Fe2—N2	92.7 (5)	C18—C17—C16	120 (2)
N6—Fe2—N4	93.3 (4)	C18—C17—H17A	119.9
N3—Fe2—N4	83.7 (5)	C16—C17—H17A	119.9
N1—Fe2—N4	172.6 (4)	C17—C18—C33	122 (2)
N5—Fe2—N4	93.5 (5)	C17—C18—C19	118 (2)
N2—Fe2—N4	89.6 (4)	C33—C18—C19	120 (2)
O3—W1—O4	104.9 (5)	C32—C19—N4	118.5 (13)
O3—W1—O7	102.9 (4)	C32—C19—C18	122.6 (18)
O4—W1—O7	86.4 (4)	N4—C19—C18	118.7 (17)
O3—W1—O12 ⁱⁱ	95.3 (5)	C21—C20—N6	117.3 (11)
O4—W1—O12 ⁱⁱ	159.7 (4)	C21—C20—C29	120.5 (12)
O7—W1—O12 ⁱⁱ	87.9 (4)	N6—C20—C29	122.1 (11)
O3—W1—O12	97.3 (4)	N5—C21—C22	123.7 (13)
O4—W1—O12	90.0 (4)	N5—C21—C20	115.0 (11)
O7—W1—O12	159.7 (4)	C22—C21—C20	121.3 (13)
O12 ⁱⁱ —W1—O12	88.6 (5)	C21—C22—C23	117.4 (16)
O3—W1—O1	163.4 (5)	C21—C22—C31	118.1 (15)
O4—W1—O1	88.2 (4)	C23—C22—C31	124.4 (16)
O7—W1—O1	87.8 (4)	C36—C23—C22	120.5 (15)
O12 ⁱⁱ —W1—O1	72.1 (4)	C36—C23—H23A	119.7
O12—W1—O1	72.1 (4)	C22—C23—H23A	119.7
O5—W2—O7	103.3 (5)	N5—C24—C36	119.5 (17)
O5—W2—O9	101.8 (4)	N5—C24—H24A	120.3
O7—W2—O9	86.7 (4)	C36—C24—H24A	120.3
O5—W2—O14 ⁱⁱ	97.7 (5)	C7—C25—C13	122.2 (14)
O7—W2—O14 ⁱⁱ	92.2 (4)	C7—C25—H25A	118.9
O9—W2—O14 ⁱⁱ	160.2 (4)	C13—C25—H25A	118.9
O5—W2—O11 ⁱⁱ	95.9 (5)	N6—C26—C27	123.7 (13)
O7—W2—O11 ⁱⁱ	160.7 (4)	N6—C26—H26A	118.2
O9—W2—O11 ⁱⁱ	86.8 (4)	C27—C26—H26A	118.2
O14 ⁱⁱ —W2—O11 ⁱⁱ	87.8 (4)	C28—C27—C26	119.4 (15)
O5—W2—O13 ⁱⁱ	166.2 (5)	C28—C27—H27A	120.3
O7—W2—O13 ⁱⁱ	88.3 (4)	C26—C27—H27A	120.3
O9—W2—O13 ⁱⁱ	86.2 (3)	C27—C28—C29	121.8 (13)
O14 ⁱⁱ —W2—O13 ⁱⁱ	74.0 (3)	C27—C28—H28A	119.1
O11 ⁱⁱ —W2—O13 ⁱⁱ	73.1 (3)	C29—C28—H28A	119.1
O6—W3—O2 ⁱⁱ	103.8 (5)	C30—C29—C20	118.0 (14)
O6—W3—O2	102.5 (5)	C30—C29—C28	126.6 (14)
O2 ⁱⁱ —W3—O2	85.6 (5)	C20—C29—C28	115.0 (12)
O6—W3—O11	96.5 (5)	C29—C30—C31	124.1 (16)
O2 ⁱⁱ —W3—O11	159.7 (4)	C29—C30—H30A	117.9

O2—W3—O11	90.1 (4)	C31—C30—H30A	117.9
O6—W3—O10	96.9 (5)	C30—C31—C22	117.2 (14)
O2 ⁱⁱ —W3—O10	90.0 (4)	C30—C31—H31A	121.4
O2—W3—O10	160.6 (4)	C22—C31—H31A	121.4
O11—W3—O10	87.5 (4)	C19—C32—C15	125.1 (16)
O6—W3—O13	165.8 (5)	C19—C32—N3	118.0 (15)
O2 ⁱⁱ —W3—O13	87.1 (4)	C15—C32—N3	116.7 (16)
O2—W3—O13	87.2 (4)	C35—C33—C18	120.0 (17)
O11—W3—O13	73.0 (3)	C35—C33—H33A	120.0
O10—W3—O13	73.7 (4)	C18—C33—H33A	120.0
O8—W4—O9	103.7 (4)	C12—C34—C7	117.0 (15)
O8—W4—O4	104.3 (5)	C12—C34—H34A	121.5
O9—W4—O4	87.5 (4)	C7—C34—H34A	121.5
O8—W4—O10	95.1 (5)	C33—C35—C14	117.1 (18)
O9—W4—O10	90.8 (4)	C33—C35—H35A	121.5
O4—W4—O10	160.3 (4)	C14—C35—H35A	121.5
O8—W4—O14	96.0 (4)	C23—C36—C24	119.6 (17)
O9—W4—O14	160.3 (4)	C23—C36—H36A	120.2
O4—W4—O14	88.3 (4)	C24—C36—H36A	120.2
O10—W4—O14	86.8 (4)	C1—N1—C5	116.9 (10)
O8—W4—O13	164.7 (4)	C1—N1—Fe2	131.6 (8)
O9—W4—O13	87.6 (3)	C5—N1—Fe2	111.5 (8)
O4—W4—O13	86.2 (4)	C11—N2—C6	119.4 (12)
O10—W4—O13	74.2 (4)	C11—N2—Fe2	130.2 (10)
O14—W4—O13	72.9 (3)	C6—N2—Fe2	110.2 (8)
N1—C1—C2	124.0 (12)	C8—N3—C32	118.3 (13)
N1—C1—H1A	118.0	C8—N3—Fe2	132.7 (10)
C2—C1—H1A	118.0	C32—N3—Fe2	108.4 (10)
C3—C2—C1	121.8 (14)	C14—N4—C19	120.3 (13)
C3—C2—H2A	119.1	C14—N4—Fe2	128.8 (12)
C1—C2—H2A	119.1	C19—N4—Fe2	110.5 (10)
C2—C3—C4	116.7 (12)	C21—N5—C24	119.2 (13)
C2—C3—H3A	121.7	C21—N5—Fe2	113.5 (9)
C4—C3—H3A	121.7	C24—N5—Fe2	127.2 (11)
C5—C4—C13	120.5 (13)	C26—N6—C20	117.7 (11)
C5—C4—C3	117.5 (11)	C26—N6—Fe2	130.1 (9)
C13—C4—C3	122.0 (12)	C20—N6—Fe2	111.7 (8)
N1—C5—C4	123.0 (12)	Fe1—O1—W1 ⁱ	119.9 (4)
N1—C5—C6	116.5 (11)	Fe1—O1—W1 ⁱⁱ	119.9 (4)
C4—C5—C6	120.5 (12)	W1 ⁱ —O1—W1 ⁱⁱ	97.4 (5)
N2—C6—C5	117.7 (11)	Fe1—O1—W1	119.9 (4)
N2—C6—C7	123.9 (13)	W1 ⁱ —O1—W1	97.4 (5)
C5—C6—C7	118.1 (13)	W1 ⁱⁱ —O1—W1	97.4 (5)
C25—C7—C6	120.5 (14)	W3 ⁱ —O2—W3	152.4 (5)
C25—C7—C34	124.2 (15)	W1—O4—W4	150.8 (5)
C6—C7—C34	115.3 (14)	W2—O7—W1	151.3 (5)
N3—C8—C9	121.7 (16)	W4—O9—W2	150.3 (5)

supplementary materials

N3—C8—H8A	119.2	W4—O10—W3	115.8 (4)
C9—C8—H8A	119.2	W3—O11—W2 ⁱ	116.6 (4)
C10—C9—C8	126.2 (19)	W1 ⁱ —O12—W1	118.3 (5)
C10—C9—H9A	116.9	Fe1—O13—W2 ⁱ	121.3 (4)
C8—C9—H9A	116.9	Fe1—O13—W4	120.8 (4)
C9—C10—C15	116.0 (18)	W2 ⁱ —O13—W4	97.7 (3)
C9—C10—H10A	122.0	Fe1—O13—W3	118.3 (5)
C15—C10—H10A	122.0	W2 ⁱ —O13—W3	97.2 (3)
N2—C11—C12	121.5 (14)	W4—O13—W3	96.0 (3)
N2—C11—H11A	119.2	W2 ⁱ —O14—W4	115.2 (4)
C12—C11—H11A	119.2		

Symmetry codes: (i) $-y+1, x-y, z$; (ii) $-x+y+1, -x+1, z$.

Fig. 1

