

Diaquabis(benzyloxyacetato)copper(II)

Sheng-Li Sun,^a Chun-Liang Chen,^a Chang-Sheng Gu,^{b*}
Weng-Dong Song^b and Xiao-Min Hao^b

^aMonitoring Center of Marine Resources and the Environment, Guangdong Ocean University, Zhanjiang 524088, People's Republic of China, and ^bDepartment of Applied Chemistry, Guangdong Ocean University, Zhanjiang 524088, People's Republic of China

Correspondence e-mail: liujiwei0706@163.com


Received 7 April 2008; accepted 17 April 2008

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.034; wR factor = 0.083; data-to-parameter ratio = 16.5.

In the title mononuclear complex, $[\text{Cu}(\text{C}_9\text{H}_9\text{O}_3)_2(\text{H}_2\text{O})_2]$, the Cu^{II} ion, located on an inversion center, is hexacoordinated by four O atoms from two benzyloxyacetate ligands [$\text{Cu}-\text{O} = 1.9420$ (14) and 2.2922 (14) \AA] and two water molecules [$\text{Cu}-\text{O} = 2.0157$ (15) \AA] in a distorted octahedral geometry. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into layers parallel to the bc plane.

Related literature

For general background, see: Eddaoudi *et al.* (2005).

**Experimental***Crystal data*

$[\text{Cu}(\text{C}_9\text{H}_9\text{O}_3)_2(\text{H}_2\text{O})_2]$	$V = 929.21$ (6) \AA^3
$M_r = 429.91$	$Z = 2$
Monoclinic, $P2_1/c$	$\text{Mo } K\alpha$ radiation
$a = 11.8847$ (4) \AA	$\mu = 1.22\text{ mm}^{-1}$
$b = 7.1509$ (2) \AA	$T = 296$ (2) K
$c = 11.6564$ (5) \AA	$0.32 \times 0.24 \times 0.18\text{ mm}$
$\beta = 110.283$ (3) $^\circ$	

Data collection

Bruker P4/APEXII diffractometer	7991 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	2144 independent reflections
$(SADABS; Sheldrick, 1996)$	1606 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.711$, $T_{\max} = 0.803$	$R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.082$	$\Delta\rho_{\max} = 0.40\text{ e } \text{\AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\min} = -0.57\text{ e } \text{\AA}^{-3}$
2144 reflections	
130 parameters	
3 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1W—H1W1 \cdots O2 ⁱ	0.839 (9)	1.978 (12)	2.796 (2)	165 (2)
O1W—H1W2 \cdots O2 ⁱⁱ	0.83 (2)	1.964 (10)	2.788 (2)	173 (2)

Symmetry codes: (i) $-x + 1$, $-y + 1$, $-z + 1$; (ii) x , $-y + \frac{1}{2}$, $z - \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by Guangdong Ocean University (Project Nos. 0612178 and 0612179) and Zhanjiang City Technology Tender (Project No. 0810014).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2398).

References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Eddaoudi, M., Chen, B., O'Keefe, M. & Yaghi, O. M. (2005). *J. Am. Chem. Soc.* **127**, 1504–1510.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m691 [doi:10.1107/S1600536808010593]

Diaquabis(benzyloxyacetato)copper(II)

S.-L. Sun, C.-L. Chen, C.-S. Gu, W.-D. Song and X.-M. Hao

Comment

Current interests in supramolecular chemistry are rapidly expanding for their intriguing architectures and potential applications (Eddaoudi *et al.*, 2005). The organic aromatic carboxylate ligand, benzyloxyacetate, has various coordination modes and can link metal centres through carboxylate groups or/and benzyloxy group into different extended architectures. Therefore, benzyloxyacetate can be considered as a good candidate to construct various metal-organic complexes. Herein we report the crystal structure of the title mononuclear complex of benzyloxyacetate, $[\text{Cu}(\text{C}_9\text{H}_9\text{O}_3)_2(\text{H}_2\text{O})_2]$, (I).

As illustrated in Fig. 1, the Cu^{II} ion lies on an inversion center and displays an octahedral geometry defined by four carboxylate O atoms from two different benzyloxyacetate ligands and two water molecules. The $\text{Cu}—\text{O}$ and $\text{Cu}—\text{Ow}$ bond lengths are 1.942 (1), 2.292 (1) and 2.016 (2) Å, respectively. The characteristic $C—O$ (carboxylate) bond lengths suggest electron localization of the carboxylate groups of the anionic ligands. In the crystal structure, intermolecular hydrogen bonds (Table 1) give rise to a supramolecular structure.

Experimental

The ligand, benzyloxyacetic acid was commercially available and used without further purification. The title complex was prepared by the addition of $\text{Cu}(\text{Ac})_2 \cdot \text{H}_2\text{O}$ (4.00 g, 20 mmol) to a hot aqueous solution of benzyloxyacetic acid (1.66 g, 10 mmol); the pH was adjusted to 6 with 0.1*M* sodium hydroxide. The solution was allowed to evaporate at room temperature. Blue prismatic crystals were separated from the filtered solution after several days. C&H analysis. Calc. for $\text{C}_{18}\text{H}_{22}\text{CuO}_8$: C 50.28, H 5.16%. Found: C 50.26, H 5.17%.

Refinement

The C-bound H atoms were placed in calculated positions, with $\text{C}—\text{H} = 0.93$ or 0.97 Å, and were refined in the riding-model approximation, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The H atoms of the water molecule were located in a difference Fourier map and refined with bond restrict $\text{O}—\text{H} = 0.84$ (2) Å in the riding-model approximation, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

Figures



Fig. 1. The molecular structure of (I) showing the atomic numbering and 30% probability displacement ellipsoids [symmetry code: (i) $-x+1, -y, -z+1$].

supplementary materials

Diaquabis(benzyloxyacetato)copper(II)

Crystal data

[Cu(C ₉ H ₉ O ₃) ₂ (H ₂ O) ₂]	$F_{000} = 446$
$M_r = 429.91$	$D_x = 1.536 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 11.8847 (4) \text{ \AA}$	Cell parameters from 7991 reflections
$b = 7.1509 (2) \text{ \AA}$	$\theta = 1.8\text{--}27.6^\circ$
$c = 11.6564 (5) \text{ \AA}$	$\mu = 1.22 \text{ mm}^{-1}$
$\beta = 110.283 (3)^\circ$	$T = 296 (2) \text{ K}$
$V = 929.21 (6) \text{ \AA}^3$	Prism, blue
$Z = 2$	$0.32 \times 0.24 \times 0.18 \text{ mm}$

Data collection

Bruker P4 diffractometer	2144 independent reflections
Radiation source: fine-focus sealed tube	1606 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.035$
Detector resolution: 10.000 pixels mm ⁻¹	$\theta_{\text{max}} = 27.6^\circ$
$T = 296(2) \text{ K}$	$\theta_{\text{min}} = 1.8^\circ$
ω scans	$h = -15 \rightarrow 15$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$k = -9 \rightarrow 8$
$T_{\text{min}} = 0.711, T_{\text{max}} = 0.803$	$l = -15 \rightarrow 10$
7991 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.082$	$w = 1/[\sigma^2(F_o^2) + (0.0429P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2144 reflections	$\Delta\rho_{\text{max}} = 0.40 \text{ e \AA}^{-3}$
130 parameters	$\Delta\rho_{\text{min}} = -0.57 \text{ e \AA}^{-3}$
3 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.5000	0.0000	0.5000	0.02860 (13)
O1	0.47417 (14)	0.2249 (2)	0.58090 (13)	0.0345 (4)
O2	0.53762 (17)	0.4957 (2)	0.67150 (14)	0.0417 (4)
O3	0.66880 (13)	0.1742 (2)	0.52517 (14)	0.0407 (4)
O1W	0.42562 (15)	0.1187 (2)	0.33393 (14)	0.0374 (4)
C1	0.5503 (2)	0.3554 (3)	0.61482 (17)	0.0325 (5)
C2	0.6644 (2)	0.3444 (3)	0.5852 (2)	0.0376 (5)
C3	0.7506 (2)	0.1723 (4)	0.4603 (2)	0.0442 (6)
C4	0.8798 (2)	0.1745 (3)	0.54208 (19)	0.0341 (5)
C5	0.9166 (2)	0.0908 (4)	0.6556 (2)	0.0427 (6)
C6	1.0355 (3)	0.0908 (4)	0.7283 (2)	0.0510 (7)
C7	1.1196 (2)	0.1751 (4)	0.6891 (2)	0.0513 (7)
C8	1.0844 (2)	0.2568 (4)	0.5758 (3)	0.0507 (7)
C9	0.9650 (2)	0.2583 (4)	0.5025 (2)	0.0430 (6)
H1W1	0.425 (2)	0.2360 (13)	0.335 (2)	0.056*
H1W2	0.457 (2)	0.075 (3)	0.286 (2)	0.056*
H2A	0.6675	0.4486	0.5331	0.045*
H2B	0.7333	0.3530	0.6601	0.045*
H3A	0.7352	0.2803	0.4067	0.053*
H3B	0.7361	0.0612	0.4094	0.053*
H5A	0.8604	0.0339	0.6833	0.051*
H6A	1.0593	0.0333	0.8045	0.061*
H7A	1.1998	0.1768	0.7392	0.062*
H8A	1.1414	0.3115	0.5482	0.061*
H9A	0.9416	0.3157	0.4262	0.052*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0348 (2)	0.0229 (2)	0.03073 (19)	-0.00185 (16)	0.01464 (16)	-0.00319 (15)
O1	0.0435 (9)	0.0261 (8)	0.0393 (8)	-0.0028 (7)	0.0212 (7)	-0.0052 (7)
O2	0.0643 (11)	0.0267 (9)	0.0400 (8)	-0.0002 (8)	0.0254 (8)	-0.0066 (7)
O1W	0.0465 (10)	0.0316 (9)	0.0357 (8)	0.0020 (8)	0.0160 (7)	0.0004 (7)
C1	0.0444 (13)	0.0287 (12)	0.0245 (9)	0.0028 (10)	0.0122 (10)	0.0021 (9)
C2	0.0400 (13)	0.0300 (13)	0.0423 (12)	-0.0057 (10)	0.0136 (11)	-0.0086 (10)
O3	0.0369 (9)	0.0375 (9)	0.0537 (9)	-0.0061 (7)	0.0234 (8)	-0.0147 (8)
C3	0.0414 (14)	0.0559 (17)	0.0389 (12)	-0.0013 (12)	0.0188 (11)	-0.0086 (11)
C4	0.0382 (12)	0.0335 (12)	0.0347 (11)	0.0017 (10)	0.0178 (10)	-0.0028 (9)
C5	0.0493 (16)	0.0419 (14)	0.0427 (12)	0.0061 (12)	0.0233 (12)	0.0054 (11)
C6	0.0590 (18)	0.0537 (17)	0.0393 (13)	0.0205 (14)	0.0158 (13)	0.0053 (12)
C7	0.0400 (14)	0.0482 (17)	0.0584 (15)	0.0070 (13)	0.0077 (13)	-0.0118 (13)
C8	0.0413 (15)	0.0456 (16)	0.0726 (18)	-0.0036 (12)	0.0289 (14)	-0.0005 (14)
C9	0.0453 (14)	0.0432 (15)	0.0459 (13)	0.0038 (11)	0.0225 (12)	0.0055 (11)

supplementary materials

Geometric parameters (\AA , $^\circ$)

Cu1—O1	1.9420 (14)	C3—C4	1.500 (3)
Cu1—O3	2.2922 (14)	C3—H3A	0.9700
Cu1—O1W	2.0157 (15)	C3—H3B	0.9700
O1—C1	1.264 (3)	C4—C5	1.379 (3)
O2—C1	1.239 (2)	C4—C9	1.386 (3)
Cu1—O1 ⁱ	1.9420 (14)	C5—C6	1.372 (4)
Cu1—O3 ⁱ	2.2922 (14)	C5—H5A	0.9300
Cu1—O1W ⁱ	2.0157 (15)	C6—C7	1.374 (4)
O3—C3	1.424 (2)	C6—H6A	0.9300
O1W—H1W1	0.839 (9)	C7—C8	1.370 (4)
O1W—H1W2	0.83 (2)	C7—H7A	0.9300
C1—C2	1.512 (3)	C8—C9	1.380 (4)
C2—O3	1.414 (2)	C8—H8A	0.9300
C2—H2A	0.9700	C9—H9A	0.9300
C2—H2B	0.9700		
O1 ⁱ —Cu1—O1	180.00 (7)	C1—C2—H2B	109.6
O1 ⁱ —Cu1—O3	103.51 (6)	C2—O3—C3	114.88 (17)
O1—Cu1—O3	76.49 (6)	C2—O3—Cu1	110.48 (12)
O1 ⁱ —Cu1—O1W	88.45 (6)	C3—O3—Cu1	130.94 (13)
O1—Cu1—O1W	91.55 (6)	C4—C3—H3A	108.9
O3—Cu1—O3 ⁱ	180.0	C4—C3—H3B	108.9
O1W—Cu1—O3	88.11 (6)	C4—C5—H5A	119.7
O1W ⁱ —Cu1—O3	91.89 (6)	C4—C9—H9A	119.9
O1W—Cu1—O1W ⁱ	180.0	C5—C4—C9	118.8 (2)
Cu1—O1W—H1W1	114.0 (18)	C5—C4—C3	121.3 (2)
Cu1—O1W—H1W2	109.5 (18)	C5—C6—C7	120.4 (2)
O1 ⁱ —Cu1—O3 ⁱ	76.49 (6)	C5—C6—H6A	119.8
O1—Cu1—O3 ⁱ	103.51 (6)	C6—C5—C4	120.6 (2)
O1 ⁱ —Cu1—O1W ⁱ	91.55 (6)	C6—C5—H5A	119.7
O1—Cu1—O1W ⁱ	88.45 (6)	C6—C7—H7A	120.2
O1—C1—C2	119.44 (18)	C7—C6—H6A	119.8
O2—C1—O1	123.9 (2)	C7—C8—C9	120.3 (2)
O2—C1—C2	116.6 (2)	C7—C8—H8A	119.9
O3—C2—C1	110.35 (18)	C8—C7—C6	119.6 (2)
O3—C2—H2A	109.6	C8—C7—H7A	120.2
O3—C2—H2B	109.6	C8—C9—C4	120.3 (2)
O3—C3—C4	113.51 (18)	C8—C9—H9A	119.9
O3—C3—H3A	108.9	C9—C8—H8A	119.9
O3—C3—H3B	108.9	C9—C4—C3	119.9 (2)
O1W—Cu1—O3 ⁱ	91.89 (6)	H1W1—O1W—H1W2	113.6 (15)
O1W ⁱ —Cu1—O3 ⁱ	88.11 (6)	H2A—C2—H2B	108.1
C1—O1—Cu1	123.14 (13)	H3A—C3—H3B	107.7
C1—C2—H2A	109.6		

Cu1—O1—C1—O2	−176.53 (15)	O1W—Cu1—O3—C2	92.76 (14)
Cu1—O1—C1—C2	3.9 (3)	O1W ⁱ —Cu1—O3—C2	−87.24 (14)
Cu1—O3—C3—C4	−134.22 (17)	O1W—Cu1—O3—C3	−63.95 (19)
O1 ⁱ —Cu1—O3—C2	−179.29 (13)	O1W ⁱ —Cu1—O3—C3	116.05 (19)
O1—Cu1—O3—C2	0.71 (13)	C1—C2—O3—C3	161.48 (18)
O1 ⁱ —Cu1—O3—C3	24.0 (2)	C1—C2—O3—Cu1	0.7 (2)
O1—Cu1—O3—C3	−156.0 (2)	C2—O3—C3—C4	69.9 (3)
O1—C1—C2—O3	−2.8 (3)	C3—C4—C5—C6	178.6 (2)
O2—C1—C2—O3	177.60 (17)	C3—C4—C9—C8	−178.3 (2)
O3—Cu1—O1—C1	−2.51 (15)	C4—C5—C6—C7	0.3 (4)
O3 ⁱ —Cu1—O1—C1	177.49 (15)	C5—C4—C9—C8	0.2 (3)
O3—C3—C4—C5	31.9 (3)	C5—C6—C7—C8	−1.2 (4)
O3—C3—C4—C9	−149.7 (2)	C6—C7—C8—C9	1.5 (4)
O1W—Cu1—O1—C1	−90.19 (16)	C7—C8—C9—C4	−1.0 (4)
O1W ⁱ —Cu1—O1—C1	89.81 (16)	C9—C4—C5—C6	0.1 (3)

Symmetry codes: (i) $-x+1, -y, -z+1$.

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1W—H1W1···O2 ⁱⁱ	0.839 (9)	1.978 (12)	2.796 (2)	165 (2)
O1W—H1W2···O2 ⁱⁱⁱ	0.83 (2)	1.964 (10)	2.788 (2)	173 (2)

Symmetry codes: (ii) $-x+1, -y+1, -z+1$; (iii) $x, -y+1/2, z-1/2$.

supplementary materials

Fig. 1

