

Hexaaquacadmium(II) bis[4-(2-hydroxybenzylideneamino)benzenesulfonate] dihydrate

Xi-Shi Tai,^{a*} Jun Xu,^b Yi-Min Feng^a and Zu-Pei Liang^a

^aDepartment of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China, and ^bWeifang Institute of Supervision and Inspection of Product Quality, Weifang 261031, People's Republic of China
Correspondence e-mail: taixishi@zu.edu.cn


Received 18 April 2008; accepted 18 April 2008

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.028; wR factor = 0.076; data-to-parameter ratio = 13.6.

In the title compound, $[\text{Cd}(\text{H}_2\text{O})_6](\text{C}_{13}\text{H}_{10}\text{NO}_4\text{S})_2 \cdot 2\text{H}_2\text{O}$, the Cd atom (site symmetry $\bar{1}$) adopts a regular octahedral coordination and the anion is stabilized by an intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond. $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds involving the coordinated and uncoordinated water molecules lead to a three-dimensional network.

Related literature

For related literature, see: Tai *et al.* (2008).



Experimental

Crystal data

$[\text{Cd}(\text{H}_2\text{O})_6](\text{C}_{13}\text{H}_{10}\text{NO}_4\text{S})_2 \cdot 2\text{H}_2\text{O}$

$M_r = 809.09$

Monoclinic, $P2_1/c$

$a = 18.464 (2)\text{ \AA}$

$b = 6.1488 (8)\text{ \AA}$

$c = 14.5701 (12)\text{ \AA}$

$\beta = 92.226 (2)^\circ$

$V = 1652.9 (3)\text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 0.86\text{ mm}^{-1}$

$T = 298 (2)\text{ K}$

$0.48 \times 0.45 \times 0.18\text{ mm}$

Data collection

Bruker SMART CCD diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2000)

$T_{\min} = 0.682$, $T_{\max} = 0.860$

7936 measured reflections

2904 independent reflections

2447 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$

$wR(F^2) = 0.075$

$S = 1.06$

2904 reflections

214 parameters

H-atom parameters constrained

$\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.60\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

Cd1–O5	2.2684 (19)	Cd1–O6	2.2862 (17)
Cd1–O7	2.2826 (18)		

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4–H4 \cdots N1	0.82	1.89	2.611 (3)	147
O5–H5A \cdots O1 ⁱ	0.85	2.07	2.909 (3)	170
O5–H5B \cdots O8 ⁱⁱ	0.85	1.92	2.750 (3)	167
O6–H6A \cdots O8 ⁱⁱⁱ	0.85	1.93	2.778 (3)	177
O6–H6B \cdots O2 ^{iv}	0.85	1.97	2.802 (3)	166
O7–H7A \cdots O1	0.85	1.95	2.793 (3)	174
O7–H7B \cdots O3 ⁱ	0.85	1.91	2.757 (3)	172
O8–H8A \cdots O2	0.85	1.90	2.750 (3)	176
O8–H8B \cdots O3 ⁱ	0.85	2.00	2.851 (3)	176

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (grant No. 20671073), the National Natural Science Foundation of Shandong (grant No. Y2007B60), the Science and Technology Foundation of Weifang and Weifang University for research grants.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2722).

References

- Bruker (2000). *SMART, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tai, X.-S., Feng, Y.-M. & Zhang, H.-X. (2008). *Acta Cryst. E* **64**, m502.

supplementary materials

Acta Cryst. (2008). E64, m694 [doi:10.1107/S1600536808010878]

Hexaaquacadmium(II) bis[4-(2-hydroxybenzylideneamino)benzenesulfonate] dihydrate

X.-S. Tai, J. Xu, Y.-M. Feng and Z.-P. Liang

Comment

As part of our ongoing studies of the synthesis and coordination chemistry of Schiff-base ligands (e.g. Tai *et al.*, 2008), we now report the synthesis and structure of the title compound, (I), (Fig. 1), in which the organic species does not coordinate to the metal and a hydrated molecular salt arises.

The Cd atom (site symmetry $\bar{1}$) in (I) is bonded to six water molecules (Table 1). The anion is stabilised by an intramolecular O—H···N hydrogen bond (Table 2), which perhaps correlates with the fact that the aromatic rings are almost co-planar [dihedral angle = 4.09 (14) $^\circ$].

The water molecules, both bound and unbound, participate in O—H···O hydrogen bonds to link the component species into a three-dimensional network.

Experimental

1 mmol of cadmium nitrate was added to a solution of salicylaldehyde-4-aminobenzene sulfonic acid (1 mmol) in 10 ml of 95% ethanol. The mixture was stirred for 2 h at refluxing temperature. Evaporating some ethanol, clear blocks of (I) were obtained after one weeks.

Refinement

The H atoms were placed geometrically (C—H = 0.93–0.96 Å, O—H = 0.85 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$.

Figures




Fig. 1. The molecular structure of (I) showing 50% displacement ellipsoids (arbitrary spheres for the H atoms). The hydrogen bonds are indicated by double-dashed lines. Symmetry code: (i) 1-x, 1-y, 1-z.

Hexaaquacadmium(II) bis[4-(2-hydroxybenzylideneamino)benzenesulfonate] dihydrate

Crystal data

[Cd(H₂O)₆](C₁₃H₁₀NO₄S)₂·2H₂O

$F_{000} = 828$

$M_r = 809.09$

$D_x = 1.626 \text{ Mg m}^{-3}$

supplementary materials

Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 18.464 (2) \text{ \AA}$	Cell parameters from 4564 reflections
$b = 6.1488 (8) \text{ \AA}$	$\theta = 2.6\text{--}27.9^\circ$
$c = 14.5701 (12) \text{ \AA}$	$\mu = 0.86 \text{ mm}^{-1}$
$\beta = 92.226 (2)^\circ$	$T = 298 (2) \text{ K}$
$V = 1652.9 (3) \text{ \AA}^3$	Block, colourless
$Z = 2$	$0.48 \times 0.45 \times 0.18 \text{ mm}$

Data collection

Bruker SMART CCD diffractometer	2904 independent reflections
Radiation source: fine-focus sealed tube	2447 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.035$
$T = 298(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
ω scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	$h = -21 \rightarrow 21$
$T_{\text{min}} = 0.682$, $T_{\text{max}} = 0.860$	$k = -7 \rightarrow 5$
7936 measured reflections	$l = -17 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H-atom parameters constrained
$wR(F^2) = 0.075$	$w = 1/[\sigma^2(F_o^2) + (0.0376P)^2 + 0.6281P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2904 reflections	$\Delta\rho_{\text{max}} = 0.28 \text{ e \AA}^{-3}$
214 parameters	$\Delta\rho_{\text{min}} = -0.60 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.5000	0.5000	0.5000	0.03115 (11)
N1	0.00853 (11)	0.8935 (4)	0.61988 (16)	0.0392 (5)
O1	0.35962 (10)	0.9727 (3)	0.59697 (14)	0.0419 (5)
O2	0.34686 (10)	0.9949 (3)	0.76142 (14)	0.0424 (5)
O3	0.33264 (10)	1.3176 (3)	0.66883 (14)	0.0425 (5)
O4	-0.09042 (11)	0.6152 (4)	0.56383 (17)	0.0655 (7)
H4	-0.0493	0.6618	0.5742	0.098*
O5	0.50034 (11)	0.1696 (3)	0.56929 (14)	0.0472 (5)
H5A	0.4573	0.1274	0.5779	0.057*
H5B	0.5235	0.1750	0.6208	0.057*
O6	0.46644 (10)	0.3400 (3)	0.36324 (12)	0.0425 (5)
H6A	0.4997	0.3570	0.3251	0.051*
H6B	0.4279	0.3987	0.3412	0.051*
O7	0.38005 (10)	0.5588 (3)	0.52336 (13)	0.0415 (5)
H7A	0.3745	0.6886	0.5418	0.050*
H7B	0.3663	0.4736	0.5652	0.050*
O8	0.42645 (10)	0.6168 (3)	0.76420 (13)	0.0462 (5)
H8A	0.4035	0.7366	0.7646	0.055*
H8B	0.4000	0.5230	0.7362	0.055*
S1	0.32404 (3)	1.08214 (11)	0.67158 (5)	0.03098 (16)
C1	0.23015 (14)	1.0294 (4)	0.65692 (18)	0.0296 (6)
C2	0.18127 (14)	1.1822 (5)	0.6840 (2)	0.0463 (8)
H2	0.1976	1.3125	0.7098	0.056*
C3	0.10786 (15)	1.1428 (5)	0.6730 (2)	0.0523 (9)
H3	0.0749	1.2470	0.6912	0.063*
C4	0.08304 (14)	0.9485 (5)	0.63495 (19)	0.0343 (6)
C5	0.13250 (15)	0.7954 (5)	0.6086 (2)	0.0447 (7)
H5	0.1163	0.6647	0.5830	0.054*
C6	0.20627 (14)	0.8343 (5)	0.6199 (2)	0.0440 (7)
H6	0.2395	0.7296	0.6027	0.053*
C7	-0.04225 (15)	1.0304 (5)	0.6320 (2)	0.0406 (7)
H7	-0.0302	1.1709	0.6507	0.049*
C8	-0.11807 (14)	0.9738 (4)	0.61762 (19)	0.0368 (7)
C9	-0.13909 (15)	0.7700 (5)	0.5839 (2)	0.0437 (7)
C10	-0.21291 (15)	0.7219 (5)	0.5712 (2)	0.0522 (8)
H10	-0.2273	0.5861	0.5491	0.063*
C11	-0.26382 (16)	0.8743 (6)	0.5911 (2)	0.0521 (8)
H11	-0.3127	0.8403	0.5826	0.063*
C12	-0.24418 (16)	1.0779 (6)	0.6235 (2)	0.0529 (8)
H12	-0.2793	1.1808	0.6361	0.064*
C13	-0.17160 (15)	1.1260 (5)	0.6369 (2)	0.0464 (7)
H13	-0.1580	1.2623	0.6592	0.056*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.03180 (16)	0.03028 (17)	0.03128 (16)	0.00014 (11)	0.00026 (11)	-0.00045 (11)
N1	0.0266 (12)	0.0439 (14)	0.0467 (14)	-0.0041 (11)	-0.0031 (10)	-0.0035 (11)
O1	0.0316 (10)	0.0456 (12)	0.0490 (12)	-0.0008 (8)	0.0082 (9)	-0.0114 (9)
O2	0.0331 (10)	0.0496 (13)	0.0438 (12)	-0.0010 (8)	-0.0079 (9)	0.0034 (9)
O3	0.0378 (10)	0.0300 (10)	0.0600 (13)	-0.0074 (8)	0.0050 (9)	-0.0035 (9)
O4	0.0424 (12)	0.0513 (14)	0.102 (2)	-0.0018 (11)	-0.0095 (12)	-0.0244 (14)
O5	0.0534 (12)	0.0412 (12)	0.0463 (12)	-0.0071 (10)	-0.0058 (9)	0.0082 (9)
O6	0.0426 (11)	0.0443 (11)	0.0399 (11)	0.0034 (9)	-0.0062 (9)	-0.0043 (9)
O7	0.0358 (10)	0.0384 (11)	0.0508 (12)	0.0016 (9)	0.0067 (9)	-0.0007 (9)
O8	0.0505 (12)	0.0381 (12)	0.0492 (12)	0.0031 (9)	-0.0094 (9)	-0.0032 (9)
S1	0.0249 (3)	0.0302 (3)	0.0378 (4)	-0.0031 (3)	0.0007 (3)	-0.0016 (3)
C1	0.0258 (13)	0.0322 (15)	0.0306 (13)	-0.0012 (11)	0.0004 (10)	-0.0005 (11)
C2	0.0330 (15)	0.0418 (17)	0.064 (2)	-0.0024 (13)	-0.0032 (14)	-0.0210 (15)
C3	0.0283 (14)	0.052 (2)	0.076 (2)	0.0050 (14)	-0.0011 (14)	-0.0259 (17)
C4	0.0285 (14)	0.0411 (16)	0.0332 (14)	-0.0034 (12)	-0.0013 (11)	-0.0016 (12)
C5	0.0329 (14)	0.0346 (16)	0.066 (2)	-0.0045 (13)	-0.0012 (13)	-0.0125 (14)
C6	0.0294 (14)	0.0326 (16)	0.070 (2)	-0.0018 (12)	0.0036 (13)	-0.0113 (14)
C7	0.0344 (15)	0.0415 (17)	0.0458 (17)	-0.0083 (13)	0.0000 (13)	0.0011 (13)
C8	0.0274 (14)	0.0446 (18)	0.0384 (15)	-0.0035 (12)	-0.0001 (11)	0.0035 (12)
C9	0.0354 (15)	0.0475 (18)	0.0477 (17)	-0.0032 (14)	-0.0036 (13)	0.0003 (14)
C10	0.0398 (17)	0.055 (2)	0.061 (2)	-0.0140 (15)	-0.0127 (14)	0.0020 (16)
C11	0.0293 (15)	0.076 (3)	0.0501 (19)	-0.0080 (16)	-0.0062 (13)	0.0125 (17)
C12	0.0331 (16)	0.069 (2)	0.057 (2)	0.0102 (16)	0.0052 (14)	0.0098 (17)
C13	0.0389 (16)	0.0487 (19)	0.0517 (18)	0.0001 (14)	0.0025 (13)	0.0026 (14)

Geometric parameters (\AA , $^\circ$)

Cd1—O5	2.2684 (19)	C1—C2	1.371 (4)
Cd1—O5 ⁱ	2.2684 (19)	C1—C6	1.380 (4)
Cd1—O7	2.2826 (18)	C2—C3	1.380 (4)
Cd1—O7 ⁱ	2.2826 (18)	C2—H2	0.9300
Cd1—O6 ⁱ	2.2862 (17)	C3—C4	1.388 (4)
Cd1—O6	2.2862 (17)	C3—H3	0.9300
N1—C7	1.278 (4)	C4—C5	1.377 (4)
N1—C4	1.425 (3)	C5—C6	1.386 (4)
O1—S1	1.4565 (19)	C5—H5	0.9300
O2—S1	1.462 (2)	C6—H6	0.9300
O3—S1	1.4570 (19)	C7—C8	1.450 (4)
O4—C9	1.349 (4)	C7—H7	0.9300
O4—H4	0.8200	C8—C9	1.395 (4)
O5—H5A	0.8499	C8—C13	1.398 (4)
O5—H5B	0.8501	C9—C10	1.400 (4)
O6—H6A	0.8500	C10—C11	1.366 (4)
O6—H6B	0.8501	C10—H10	0.9300

O7—H7A	0.8500	C11—C12	1.381 (5)
O7—H7B	0.8500	C11—H11	0.9300
O8—H8A	0.8499	C12—C13	1.379 (4)
O8—H8B	0.8499	C12—H12	0.9300
S1—C1	1.768 (3)	C13—H13	0.9300
O5—Cd1—O5 ⁱ	180.0	C1—C2—C3	120.1 (3)
O5—Cd1—O7	93.52 (7)	C1—C2—H2	120.0
O5 ⁱ —Cd1—O7	86.48 (7)	C3—C2—H2	120.0
O5—Cd1—O7 ⁱ	86.48 (7)	C2—C3—C4	120.3 (3)
O5 ⁱ —Cd1—O7 ⁱ	93.52 (7)	C2—C3—H3	119.8
O7—Cd1—O7 ⁱ	180.0	C4—C3—H3	119.8
O5—Cd1—O6 ⁱ	90.09 (7)	C5—C4—C3	119.2 (2)
O5 ⁱ —Cd1—O6 ⁱ	89.91 (7)	C5—C4—N1	116.2 (2)
O7—Cd1—O6 ⁱ	91.93 (7)	C3—C4—N1	124.6 (3)
O7 ⁱ —Cd1—O6 ⁱ	88.07 (7)	C4—C5—C6	120.5 (3)
O5—Cd1—O6	89.91 (7)	C4—C5—H5	119.7
O5 ⁱ —Cd1—O6	90.09 (7)	C6—C5—H5	119.7
O7—Cd1—O6	88.07 (7)	C1—C6—C5	119.6 (3)
O7 ⁱ —Cd1—O6	91.93 (7)	C1—C6—H6	120.2
O6 ⁱ —Cd1—O6	180.0	C5—C6—H6	120.2
C7—N1—C4	122.1 (2)	N1—C7—C8	122.1 (3)
C9—O4—H4	109.5	N1—C7—H7	119.0
Cd1—O5—H5A	110.6	C8—C7—H7	119.0
Cd1—O5—H5B	110.5	C9—C8—C13	118.9 (3)
H5A—O5—H5B	108.8	C9—C8—C7	121.4 (3)
Cd1—O6—H6A	109.8	C13—C8—C7	119.7 (3)
Cd1—O6—H6B	110.0	O4—C9—C8	122.1 (2)
H6A—O6—H6B	108.4	O4—C9—C10	118.4 (3)
Cd1—O7—H7A	109.0	C8—C9—C10	119.5 (3)
Cd1—O7—H7B	109.2	C11—C10—C9	120.1 (3)
H7A—O7—H7B	107.9	C11—C10—H10	120.0
H8A—O8—H8B	108.3	C9—C10—H10	120.0
O1—S1—O3	112.68 (12)	C10—C11—C12	121.4 (3)
O1—S1—O2	112.09 (12)	C10—C11—H11	119.3
O3—S1—O2	111.19 (12)	C12—C11—H11	119.3
O1—S1—C1	107.08 (12)	C13—C12—C11	118.9 (3)
O3—S1—C1	106.63 (11)	C13—C12—H12	120.5
O2—S1—C1	106.74 (12)	C11—C12—H12	120.5
C2—C1—C6	120.2 (2)	C12—C13—C8	121.2 (3)
C2—C1—S1	119.5 (2)	C12—C13—H13	119.4
C6—C1—S1	120.2 (2)	C8—C13—H13	119.4
O1—S1—C1—C2	-148.3 (2)	S1—C1—C6—C5	-179.8 (2)
O3—S1—C1—C2	-27.4 (3)	C4—C5—C6—C1	-0.8 (5)
O2—S1—C1—C2	91.5 (2)	C4—N1—C7—C8	-179.3 (3)
O1—S1—C1—C6	32.8 (3)	N1—C7—C8—C9	-4.2 (4)
O3—S1—C1—C6	153.7 (2)	N1—C7—C8—C13	176.4 (3)

supplementary materials

O2—S1—C1—C6	−87.4 (2)	C13—C8—C9—O4	−180.0 (3)
C6—C1—C2—C3	−1.0 (5)	C7—C8—C9—O4	0.6 (4)
S1—C1—C2—C3	−179.9 (3)	C13—C8—C9—C10	−0.7 (4)
C1—C2—C3—C4	0.2 (5)	C7—C8—C9—C10	179.9 (3)
C2—C3—C4—C5	0.3 (5)	O4—C9—C10—C11	179.7 (3)
C2—C3—C4—N1	−179.3 (3)	C8—C9—C10—C11	0.5 (5)
C7—N1—C4—C5	−172.3 (3)	C9—C10—C11—C12	0.3 (5)
C7—N1—C4—C3	7.3 (5)	C10—C11—C12—C13	−0.8 (5)
C3—C4—C5—C6	0.0 (5)	C11—C12—C13—C8	0.5 (5)
N1—C4—C5—C6	179.6 (3)	C9—C8—C13—C12	0.2 (4)
C2—C1—C6—C5	1.3 (4)	C7—C8—C13—C12	179.6 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , °)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O4—H4 \cdots N1	0.82	1.89	2.611 (3)
O5—H5A \cdots O1 ⁱⁱ	0.85	2.07	2.909 (3)
O5—H5B \cdots O8 ⁱⁱⁱ	0.85	1.92	2.750 (3)
O6—H6A \cdots O8 ⁱ	0.85	1.93	2.778 (3)
O6—H6B \cdots O2 ^{iv}	0.85	1.97	2.802 (3)
O7—H7A \cdots O1	0.85	1.95	2.793 (3)
O7—H7B \cdots O3 ⁱⁱ	0.85	1.91	2.757 (3)
O8—H8A \cdots O2	0.85	1.90	2.750 (3)
O8—H8B \cdots O3 ⁱⁱ	0.85	2.00	2.851 (3)

Symmetry codes: (ii) $x, y-1, z$; (iii) $-x+1, y-1/2, -z+3/2$; (i) $-x+1, -y+1, -z+1$; (iv) $x, -y+3/2, z-1/2$.

Fig. 1

