

## 2-Amino-4-(4-chlorophenyl)-6-ferrocenylpyridine-3-carbonitrile

Jinpeng Zhang,<sup>a</sup> Shu Yan,<sup>b</sup> Zhen He,<sup>a</sup> Lichun Xu<sup>a</sup> and Shuiping Huang<sup>a\*</sup>

<sup>a</sup>Department of Public Health, Xuzhou Medical College, Xuzhou 221000, People's Republic of China, and <sup>b</sup>College of Chemistry and Chemical Engineering, Xuzhou Normal University, Xuzhou 221116, People's Republic of China

Correspondence e-mail: laotu2001@263.net


Received 28 March 2008; accepted 1 April 2008

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.006$  Å;  $R$  factor = 0.046;  $wR$  factor = 0.093; data-to-parameter ratio = 13.4.

In the molecule of the title compound,  $[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{17}\text{H}_{11}\text{ClN}_3)]$ , the dihedral angles between the two five-membered rings and between the two six-membered rings are  $3.28$  (4) and  $51.33$  (4)°, respectively. In the crystal structure, intermolecular  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bonds link the molecules into centrosymmetric dimers.

### Related literature

For general background, see: Dombrowski *et al.* (1986); Alyoubi (2000); Desai & Shah (2003); Murata *et al.* (2004).



### Experimental

#### Crystal data

$[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{17}\text{H}_{11}\text{ClN}_3)]$	$V = 1853.7$ (4) Å <sup>3</sup>
$M_r = 413.68$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 12.1517$ (13) Å	$\mu = 0.97$ mm <sup>-1</sup>
$b = 7.4214$ (11) Å	$T = 298$ (2) K
$c = 20.742$ (2) Å	$0.20 \times 0.15 \times 0.09$ mm
$\beta = 97.691$ (2)°	

#### Data collection

Bruker SMART CCD area-detector diffractometer	8876 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	3260 independent reflections
$T_{\min} = 0.830$ , $T_{\max} = 0.918$	2298 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.041$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	244 parameters
$wR(F^2) = 0.093$	H-atom parameters constrained
$S = 1.09$	$\Delta\rho_{\text{max}} = 0.44$ e Å <sup>-3</sup>
3260 reflections	$\Delta\rho_{\text{min}} = -0.43$ e Å <sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2B}\cdots\text{N3}^i$	0.86	2.28	3.047 (5)	149

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINTE* (Bruker, 1999); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (grant No. 20672090) and the Natural Science Foundation of Jiangsu Province (grant No. BK2006033) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2443).

### References

- Alyoubi, A. O. (2000). *Spectrochim. Acta Part A*, **56**, 2397–2404.  
 Bruker (1998). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (1999). *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Desai, J. M. & Shah, V. H. (2003). *Indian J. Chem. Sect. B*, **42**, 382–385.  
 Dombrowski, K. E., Baldwin, W. & Sheats, J. E. (1986). *J. Organomet. Chem.* **302**, 281–306.  
 Murata, T., Shimada, M., Sakakibara, S., Yoshino, T., Masuda, T., Shintani, T., Sato, H., Koriyama, Y., Fukushima, K., Nunami, N., Yamauchi, M., Fuchikami, K., Komura, H., Watanabe, A., Ziegelbauer, K. B., Bacon, K. B. & Lowinger, T. B. (2004). *Bioorg. Med. Chem. Lett.* **14**, 4019–4022.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2008). E64, m730 [ doi:10.1107/S1600536808008714 ]

## 2-Amino-4-(4-chlorophenyl)-6-ferrocenylpyridine-3-carbonitrile

J. Zhang, S. Yan, Z. He, L. Xu and S. Huang

### Comment

Metalloenes are known to exhibit a wide range of biological activity. Among them, ferrocene has attracted special attention since it is neutral, chemically stable, non-toxic and able to cross cell membranes (Dombrowski *et al.*, 1986). In fact, it is now well established that the incorporation of ferrocene units into organic molecules introduces significant and new properties in these materials. In addition, it has been demonstrated that molecules containing cyanopyridine moiety may be able to work as ligands towards transition-metal ions (Alyoubi, 2000), new drugs (Murata *et al.*, 2004; Desai & Shah, 2003) and significant intermediates for the synthesis of important vitamins such as nicotinic acids and nicotinamides. For these reasons, the synthesis of new compounds containing cyanopyridine derivatives is strongly desired. We report herein the crystal structure of the title compound, (I).

In the molecule of (I), (Fig. 1), rings A (N1/C1-C5), B (C17-C22), C (C6-C10) and D (C11-C15) are, of course, planar. The dihedral angles between them are A/B = 51.33 (4)°, A/C = 13.20 (3)°, A/D = 16.32 (4)°, B/C = 46.38 (3)°, B/D = 44.47 (3)° and C/D = 3.28 (4)°. So, rings C and D are nearly parallel to each other.

In the crystal structure, intermolecular N-H...N hydrogen bonds (Table 1) link the molecules into centrosymmetric dimers (Fig. 2), in which they may be effective in the stabilization of the structure.

### Experimental

Compound (I) was prepared by the reaction of 4-chlorobenzaldehyde (2 mmol), malononitrile (2 mmol), acetylferrocene (2 mmol) and ammonium acetate (4 mmol) in water (2 ml). Single crystals of (I) suitable for X-ray analysis were obtained by slow evaporation of an aqueous ethanol solution (95%) (yield; 95%, m.p. 548-550 K). IR (cm<sup>-1</sup>): 3457, 3354, 2211; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 4.10 (5H, s, ferrocenyl), 4.50 (2H, s, ferrocenyl), 5.04 (2H, s, ferrocenyl), 6.81 (2H, brs, NH<sub>2</sub>), 6.95 (1H, s, ArH), 7.64 (2H, d, J = 8.4 Hz, ArH), 7.68 (2H, d, J = 8.4 Hz, ArH), 7.87 (2H, brs, NH<sub>2</sub>), 7.88–8.01 (4H, m, ArH), 11.85 (1H, s, NH).

### Refinement

H atoms were positioned geometrically, with N-H = 0.86 Å (for NH<sub>2</sub>) and C-H = 0.93 Å for aromatic H, and constrained to ride on their parent atoms with U<sub>iso</sub>(H) = 1.2U<sub>eq</sub>(C,N).

## Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. A partial packing diagram of (I). Hydrogen bonds are shown as dashed lines. Hydrogen atoms not involved in hydrogen bonding are omitted for clarity.

## 2-Amino-4-(4-chlorophenyl)-6-ferrocenylpyridine-3-carbonitrile

### Crystal data

[Fe(C<sub>5</sub>H<sub>5</sub>)(C<sub>17</sub>H<sub>11</sub>ClN<sub>3</sub>)]

$M_r = 413.68$

Monoclinic,  $P2_1/n$

Hall symbol: -P 2yn

$a = 12.1517 (13) \text{ \AA}$

$b = 7.4214 (11) \text{ \AA}$

$c = 20.742 (2) \text{ \AA}$

$\beta = 97.691 (2)^\circ$

$V = 1853.7 (4) \text{ \AA}^3$

$Z = 4$

$F_{000} = 848$

$D_x = 1.482 \text{ Mg m}^{-3}$

Melting point = 548–550 K

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2405 reflections

$\theta = 2.9\text{--}26.2^\circ$

$\mu = 0.97 \text{ mm}^{-1}$

$T = 298 (2) \text{ K}$

Block, red

$0.20 \times 0.15 \times 0.09 \text{ mm}$

### Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298(2) \text{ K}$

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.830$ ,  $T_{\max} = 0.918$

8876 measured reflections

3260 independent reflections

2298 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.041$

$\theta_{\text{max}} = 25.0^\circ$

$\theta_{\text{min}} = 1.8^\circ$

$h = -14 \rightarrow 14$

$k = -8 \rightarrow 6$

$l = -24 \rightarrow 24$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.046$	H-atom parameters constrained
$wR(F^2) = 0.093$	$w = 1/[\sigma^2(F_o^2) + (0.0178P)^2 + 2.3504P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
3260 reflections	$(\Delta/\sigma)_{\max} < 0.001$
244 parameters	$\Delta\rho_{\max} = 0.44 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.43 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.38223 (4)	0.42499 (7)	0.10732 (3)	0.03424 (16)
Cl1	0.81792 (10)	1.44271 (17)	0.38672 (6)	0.0694 (4)
N1	0.3585 (2)	0.5432 (4)	0.27711 (13)	0.0335 (7)
N2	0.3644 (2)	0.4443 (4)	0.38140 (14)	0.0450 (8)
H2A	0.3154	0.3649	0.3672	0.054*
H2B	0.3892	0.4491	0.4222	0.054*
N3	0.5500 (3)	0.6983 (5)	0.48341 (17)	0.0652 (12)
C1	0.4019 (3)	0.5604 (5)	0.33992 (16)	0.0321 (8)
C2	0.4807 (3)	0.6949 (5)	0.36089 (16)	0.0301 (8)
C3	0.5168 (3)	0.8132 (5)	0.31540 (17)	0.0307 (8)
C4	0.4739 (3)	0.7885 (5)	0.25086 (17)	0.0309 (8)
H4	0.4977	0.8613	0.2189	0.037*
C5	0.3950 (3)	0.6543 (5)	0.23368 (16)	0.0292 (8)
C6	0.3441 (3)	0.6283 (5)	0.16610 (16)	0.0287 (8)
C7	0.3802 (3)	0.6998 (5)	0.10848 (18)	0.0366 (9)
H7	0.4408	0.7755	0.1070	0.044*
C8	0.3080 (3)	0.6355 (5)	0.05402 (17)	0.0396 (10)
H8	0.3125	0.6617	0.0106	0.048*

## supplementary materials

---

C9	0.2277 (3)	0.5243 (5)	0.07769 (17)	0.0390 (10)
H9	0.1701	0.4648	0.0523	0.047*
C10	0.2493 (3)	0.5186 (5)	0.14592 (17)	0.0345 (9)
H10	0.2088	0.4543	0.1732	0.041*
C11	0.5039 (4)	0.2760 (6)	0.1590 (2)	0.0615 (13)
H11	0.5385	0.3005	0.2008	0.074*
C12	0.5390 (3)	0.3359 (6)	0.1002 (2)	0.0582 (13)
H12	0.6003	0.4081	0.0964	0.070*
C13	0.4640 (4)	0.2662 (6)	0.0484 (2)	0.0562 (12)
H13	0.4672	0.2835	0.0042	0.067*
C14	0.3841 (4)	0.1665 (6)	0.0753 (3)	0.0594 (13)
H14	0.3249	0.1057	0.0519	0.071*
C15	0.4071 (4)	0.1726 (6)	0.1426 (3)	0.0648 (14)
H15	0.3658	0.1178	0.1717	0.078*
C16	0.5217 (3)	0.7025 (5)	0.42883 (19)	0.0409 (10)
C17	0.5934 (3)	0.9656 (5)	0.33533 (17)	0.0308 (8)
C18	0.5737 (3)	1.0847 (6)	0.38359 (19)	0.0469 (10)
H18	0.5127	1.0663	0.4055	0.056*
C19	0.6424 (3)	1.2308 (6)	0.4002 (2)	0.0489 (11)
H19	0.6276	1.3099	0.4328	0.059*
C20	0.7325 (3)	1.2576 (5)	0.36813 (19)	0.0429 (10)
C21	0.7559 (3)	1.1393 (6)	0.3206 (2)	0.0487 (11)
H21	0.8182	1.1567	0.2998	0.058*
C22	0.6861 (3)	0.9946 (6)	0.30423 (19)	0.0450 (10)
H22	0.7015	0.9155	0.2718	0.054*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Fe1	0.0363 (3)	0.0281 (3)	0.0382 (3)	0.0077 (3)	0.0046 (2)	-0.0010 (3)
Cl1	0.0680 (7)	0.0544 (8)	0.0859 (9)	-0.0359 (6)	0.0103 (6)	-0.0084 (7)
N1	0.0387 (16)	0.0311 (18)	0.0301 (15)	-0.0057 (15)	0.0022 (13)	0.0036 (15)
N2	0.0533 (19)	0.049 (2)	0.0304 (16)	-0.0262 (18)	-0.0027 (14)	0.0062 (18)
N3	0.085 (3)	0.067 (3)	0.039 (2)	-0.033 (2)	-0.0095 (19)	0.010 (2)
C1	0.0330 (18)	0.033 (2)	0.0308 (19)	-0.0026 (18)	0.0045 (15)	0.0014 (19)
C2	0.0306 (19)	0.029 (2)	0.0293 (19)	-0.0047 (16)	0.0005 (16)	-0.0001 (17)
C3	0.0280 (18)	0.027 (2)	0.038 (2)	-0.0002 (16)	0.0048 (16)	-0.0007 (17)
C4	0.0320 (19)	0.029 (2)	0.032 (2)	-0.0036 (16)	0.0054 (16)	0.0039 (17)
C5	0.0295 (18)	0.0260 (19)	0.0326 (19)	0.0050 (16)	0.0056 (16)	0.0004 (17)
C6	0.0323 (19)	0.027 (2)	0.0264 (18)	0.0038 (16)	0.0022 (15)	0.0028 (16)
C7	0.042 (2)	0.026 (2)	0.042 (2)	0.0062 (19)	0.0067 (18)	0.006 (2)
C8	0.050 (2)	0.042 (2)	0.0275 (19)	0.011 (2)	0.0043 (18)	0.0011 (19)
C9	0.039 (2)	0.040 (2)	0.036 (2)	0.0075 (18)	-0.0011 (17)	-0.0050 (19)
C10	0.0318 (19)	0.035 (2)	0.036 (2)	0.0041 (17)	0.0018 (16)	-0.0006 (18)
C11	0.065 (3)	0.054 (3)	0.061 (3)	0.031 (3)	-0.008 (3)	0.001 (3)
C12	0.039 (2)	0.052 (3)	0.084 (4)	0.018 (2)	0.006 (2)	-0.009 (3)
C13	0.058 (3)	0.049 (3)	0.064 (3)	0.014 (2)	0.017 (2)	-0.017 (3)
C14	0.065 (3)	0.033 (3)	0.079 (4)	0.007 (2)	0.006 (3)	-0.011 (3)

C15	0.075 (3)	0.038 (3)	0.083 (4)	0.017 (3)	0.017 (3)	0.015 (3)
C16	0.048 (2)	0.036 (2)	0.037 (2)	-0.0157 (19)	0.0013 (19)	0.006 (2)
C17	0.0309 (18)	0.026 (2)	0.0358 (19)	-0.0052 (16)	0.0041 (15)	0.0030 (17)
C18	0.043 (2)	0.044 (2)	0.057 (3)	-0.014 (2)	0.0189 (19)	-0.012 (2)
C19	0.053 (3)	0.040 (2)	0.056 (3)	-0.011 (2)	0.013 (2)	-0.012 (2)
C20	0.043 (2)	0.039 (2)	0.046 (2)	-0.014 (2)	0.0060 (19)	0.000 (2)
C21	0.041 (2)	0.049 (3)	0.060 (3)	-0.013 (2)	0.019 (2)	0.002 (2)
C22	0.045 (2)	0.045 (2)	0.046 (2)	-0.008 (2)	0.0123 (19)	-0.004 (2)

*Geometric parameters (Å, °)*

Fe1—C15	2.019 (5)	C7—C8	1.417 (5)
Fe1—C10	2.020 (3)	C7—H7	0.9300
Fe1—C14	2.031 (4)	C8—C9	1.415 (5)
Fe1—C6	2.032 (3)	C8—H8	0.9300
Fe1—C11	2.033 (4)	C9—C10	1.405 (5)
Fe1—C9	2.034 (4)	C9—H9	0.9300
Fe1—C7	2.040 (4)	C10—H10	0.9300
Fe1—C12	2.040 (4)	C11—C15	1.408 (6)
Fe1—C13	2.048 (4)	C11—C12	1.415 (6)
Fe1—C8	2.052 (4)	C11—H11	0.9300
C11—C20	1.734 (4)	C12—C13	1.412 (6)
N1—C5	1.339 (4)	C12—H12	0.9300
N1—C1	1.344 (4)	C13—C14	1.395 (6)
N2—C1	1.340 (4)	C13—H13	0.9300
N2—H2A	0.8600	C14—C15	1.385 (6)
N2—H2B	0.8600	C14—H14	0.9300
N3—C16	1.139 (4)	C15—H15	0.9300
C1—C2	1.411 (5)	C17—C18	1.380 (5)
C2—C3	1.401 (5)	C17—C22	1.388 (5)
C2—C16	1.432 (5)	C18—C19	1.384 (5)
C3—C4	1.382 (5)	C18—H18	0.9300
C3—C17	1.488 (5)	C19—C20	1.369 (5)
C4—C5	1.396 (5)	C19—H19	0.9300
C4—H4	0.9300	C20—C21	1.377 (5)
C5—C6	1.467 (4)	C21—C22	1.383 (5)
C6—C10	1.426 (5)	C21—H21	0.9300
C6—C7	1.429 (5)	C22—H22	0.9300
C15—Fe1—C10	105.34 (18)	C6—C7—Fe1	69.2 (2)
C15—Fe1—C14	40.01 (18)	C8—C7—H7	125.8
C10—Fe1—C14	119.93 (18)	C6—C7—H7	125.8
C15—Fe1—C6	120.45 (18)	Fe1—C7—H7	126.4
C10—Fe1—C6	41.22 (13)	C9—C8—C7	107.6 (3)
C14—Fe1—C6	155.60 (18)	C9—C8—Fe1	69.1 (2)
C15—Fe1—C11	40.67 (18)	C7—C8—Fe1	69.3 (2)
C10—Fe1—C11	122.83 (18)	C9—C8—H8	126.2
C14—Fe1—C11	67.62 (19)	C7—C8—H8	126.2
C6—Fe1—C11	107.21 (16)	Fe1—C8—H8	127.0
C15—Fe1—C9	122.21 (19)	C10—C9—C8	108.8 (3)

## supplementary materials

---

C10—Fe1—C9	40.55 (13)	C10—C9—Fe1	69.2 (2)
C14—Fe1—C9	107.10 (17)	C8—C9—Fe1	70.4 (2)
C6—Fe1—C9	68.69 (14)	C10—C9—H9	125.6
C11—Fe1—C9	158.94 (19)	C8—C9—H9	125.6
C15—Fe1—C7	157.7 (2)	Fe1—C9—H9	126.4
C10—Fe1—C7	68.88 (15)	C9—C10—C6	108.3 (3)
C14—Fe1—C7	161.62 (19)	C9—C10—Fe1	70.3 (2)
C6—Fe1—C7	41.10 (13)	C6—C10—Fe1	69.86 (19)
C11—Fe1—C7	123.14 (18)	C9—C10—H10	125.9
C9—Fe1—C7	68.23 (15)	C6—C10—H10	125.9
C15—Fe1—C12	68.25 (19)	Fe1—C10—H10	125.6
C10—Fe1—C12	160.75 (17)	C15—C11—C12	107.5 (4)
C14—Fe1—C12	67.58 (19)	C15—C11—Fe1	69.1 (2)
C6—Fe1—C12	125.06 (16)	C12—C11—Fe1	69.9 (2)
C11—Fe1—C12	40.67 (17)	C15—C11—H11	126.2
C9—Fe1—C12	158.22 (17)	C12—C11—H11	126.2
C7—Fe1—C12	109.75 (18)	Fe1—C11—H11	126.3
C15—Fe1—C13	67.7 (2)	C13—C12—C11	107.6 (4)
C10—Fe1—C13	155.78 (16)	C13—C12—Fe1	70.1 (2)
C14—Fe1—C13	40.00 (17)	C11—C12—Fe1	69.4 (2)
C6—Fe1—C13	162.41 (16)	C13—C12—H12	126.2
C11—Fe1—C13	67.98 (19)	C11—C12—H12	126.2
C9—Fe1—C13	122.00 (16)	Fe1—C12—H12	125.9
C7—Fe1—C13	126.16 (18)	C14—C13—C12	107.5 (4)
C12—Fe1—C13	40.39 (16)	C14—C13—Fe1	69.3 (2)
C15—Fe1—C8	159.28 (19)	C12—C13—Fe1	69.5 (2)
C10—Fe1—C8	68.55 (15)	C14—C13—H13	126.2
C14—Fe1—C8	124.52 (18)	C12—C13—H13	126.2
C6—Fe1—C8	68.82 (13)	Fe1—C13—H13	126.5
C11—Fe1—C8	159.14 (19)	C15—C14—C13	109.2 (5)
C9—Fe1—C8	40.52 (14)	C15—C14—Fe1	69.5 (3)
C7—Fe1—C8	40.53 (14)	C13—C14—Fe1	70.7 (3)
C12—Fe1—C8	123.74 (18)	C15—C14—H14	125.4
C13—Fe1—C8	109.26 (17)	C13—C14—H14	125.4
C5—N1—C1	118.1 (3)	Fe1—C14—H14	126.0
C1—N2—H2A	120.0	C14—C15—C11	108.1 (5)
C1—N2—H2B	120.0	C14—C15—Fe1	70.5 (3)
H2A—N2—H2B	120.0	C11—C15—Fe1	70.2 (3)
N2—C1—N1	116.0 (3)	C14—C15—H15	126.0
N2—C1—C2	122.2 (3)	C11—C15—H15	126.0
N1—C1—C2	121.8 (3)	Fe1—C15—H15	125.0
C3—C2—C1	119.8 (3)	N3—C16—C2	175.2 (4)
C3—C2—C16	122.6 (3)	C18—C17—C22	117.9 (3)
C1—C2—C16	117.6 (3)	C18—C17—C3	121.7 (3)
C4—C3—C2	117.2 (3)	C22—C17—C3	120.4 (3)
C4—C3—C17	120.7 (3)	C17—C18—C19	121.7 (4)
C2—C3—C17	122.0 (3)	C17—C18—H18	119.2
C3—C4—C5	119.9 (3)	C19—C18—H18	119.2
C3—C4—H4	120.0	C20—C19—C18	119.2 (4)



C5—C4—H4	120.0	C20—C19—H19	120.4
N1—C5—C4	123.1 (3)	C18—C19—H19	120.4
N1—C5—C6	115.2 (3)	C19—C20—C21	120.7 (4)
C4—C5—C6	121.7 (3)	C19—C20—C11	120.0 (3)
C10—C6—C7	107.0 (3)	C21—C20—C11	119.3 (3)
C10—C6—C5	125.2 (3)	C20—C21—C22	119.5 (4)
C7—C6—C5	127.8 (3)	C20—C21—H21	120.3
C10—C6—Fe1	68.9 (2)	C22—C21—H21	120.3
C7—C6—Fe1	69.73 (19)	C21—C22—C17	121.0 (4)
C5—C6—Fe1	124.7 (2)	C21—C22—H22	119.5
C8—C7—C6	108.4 (3)	C17—C22—H22	119.5
C8—C7—Fe1	70.2 (2)		
C5—N1—C1—N2	-178.8 (3)	C13—Fe1—C10—C9	-52.3 (5)
C5—N1—C1—C2	2.2 (5)	C8—Fe1—C10—C9	37.1 (2)
N2—C1—C2—C3	-179.8 (3)	C15—Fe1—C10—C6	119.0 (2)
N1—C1—C2—C3	-0.9 (5)	C14—Fe1—C10—C6	159.6 (2)
N2—C1—C2—C16	0.9 (5)	C11—Fe1—C10—C6	78.3 (3)
N1—C1—C2—C16	179.8 (3)	C9—Fe1—C10—C6	-119.1 (3)
C1—C2—C3—C4	-1.4 (5)	C7—Fe1—C10—C6	-38.3 (2)
C16—C2—C3—C4	177.8 (3)	C12—Fe1—C10—C6	51.3 (6)
C1—C2—C3—C17	175.5 (3)	C13—Fe1—C10—C6	-171.4 (4)
C16—C2—C3—C17	-5.3 (5)	C8—Fe1—C10—C6	-81.9 (2)
C2—C3—C4—C5	2.3 (5)	C10—Fe1—C11—C15	74.5 (3)
C17—C3—C4—C5	-174.6 (3)	C14—Fe1—C11—C15	-37.6 (3)
C1—N1—C5—C4	-1.3 (5)	C6—Fe1—C11—C15	117.0 (3)
C1—N1—C5—C6	-179.9 (3)	C9—Fe1—C11—C15	41.7 (6)
C3—C4—C5—N1	-1.1 (5)	C7—Fe1—C11—C15	159.3 (3)
C3—C4—C5—C6	177.5 (3)	C12—Fe1—C11—C15	-118.8 (4)
N1—C5—C6—C10	10.9 (5)	C13—Fe1—C11—C15	-81.0 (3)
C4—C5—C6—C10	-167.8 (3)	C8—Fe1—C11—C15	-167.5 (4)
N1—C5—C6—C7	-166.7 (3)	C15—Fe1—C11—C12	118.8 (4)
C4—C5—C6—C7	14.6 (6)	C10—Fe1—C11—C12	-166.7 (3)
N1—C5—C6—Fe1	-76.5 (4)	C14—Fe1—C11—C12	81.2 (3)
C4—C5—C6—Fe1	104.9 (4)	C6—Fe1—C11—C12	-124.2 (3)
C15—Fe1—C6—C10	-78.2 (3)	C9—Fe1—C11—C12	160.5 (4)
C14—Fe1—C6—C10	-46.9 (5)	C7—Fe1—C11—C12	-81.9 (3)
C11—Fe1—C6—C10	-120.5 (2)	C13—Fe1—C11—C12	37.8 (3)
C9—Fe1—C6—C10	37.6 (2)	C8—Fe1—C11—C12	-48.6 (6)
C7—Fe1—C6—C10	118.5 (3)	C15—C11—C12—C13	-0.8 (5)
C12—Fe1—C6—C10	-161.7 (2)	Fe1—C11—C12—C13	-59.9 (3)
C13—Fe1—C6—C10	168.2 (5)	C15—C11—C12—Fe1	59.1 (3)
C8—Fe1—C6—C10	81.2 (2)	C15—Fe1—C12—C13	80.8 (3)
C15—Fe1—C6—C7	163.4 (2)	C10—Fe1—C12—C13	154.6 (5)
C10—Fe1—C6—C7	-118.5 (3)	C14—Fe1—C12—C13	37.4 (3)
C14—Fe1—C6—C7	-165.4 (4)	C6—Fe1—C12—C13	-166.5 (3)
C11—Fe1—C6—C7	121.0 (3)	C11—Fe1—C12—C13	118.7 (4)
C9—Fe1—C6—C7	-80.9 (2)	C9—Fe1—C12—C13	-42.5 (6)
C12—Fe1—C6—C7	79.9 (3)	C7—Fe1—C12—C13	-123.0 (3)
C13—Fe1—C6—C7	49.8 (6)	C8—Fe1—C12—C13	-80.0 (3)

## supplementary materials

---

C8—Fe1—C6—C7	-37.3 (2)	C15—Fe1—C12—C11	-37.9 (3)
C15—Fe1—C6—C5	40.8 (4)	C10—Fe1—C12—C11	35.9 (7)
C10—Fe1—C6—C5	118.9 (4)	C14—Fe1—C12—C11	-81.3 (3)
C14—Fe1—C6—C5	72.1 (5)	C6—Fe1—C12—C11	74.8 (3)
C11—Fe1—C6—C5	-1.6 (4)	C9—Fe1—C12—C11	-161.2 (4)
C9—Fe1—C6—C5	156.5 (3)	C7—Fe1—C12—C11	118.3 (3)
C7—Fe1—C6—C5	-122.6 (4)	C13—Fe1—C12—C11	-118.7 (4)
C12—Fe1—C6—C5	-42.7 (4)	C8—Fe1—C12—C11	161.3 (3)
C13—Fe1—C6—C5	-72.8 (6)	C11—C12—C13—C14	0.4 (5)
C8—Fe1—C6—C5	-159.8 (3)	Fe1—C12—C13—C14	-59.1 (3)
C10—C6—C7—C8	0.4 (4)	C11—C12—C13—Fe1	59.5 (3)
C5—C6—C7—C8	178.3 (3)	C15—Fe1—C13—C14	36.9 (3)
Fe1—C6—C7—C8	59.5 (3)	C10—Fe1—C13—C14	-40.8 (6)
C10—C6—C7—Fe1	-59.1 (2)	C6—Fe1—C13—C14	158.3 (5)
C5—C6—C7—Fe1	118.8 (4)	C11—Fe1—C13—C14	81.0 (3)
C15—Fe1—C7—C8	-160.1 (4)	C9—Fe1—C13—C14	-78.2 (3)
C10—Fe1—C7—C8	-81.3 (2)	C7—Fe1—C13—C14	-163.3 (3)
C14—Fe1—C7—C8	41.0 (6)	C12—Fe1—C13—C14	119.0 (4)
C6—Fe1—C7—C8	-119.7 (3)	C8—Fe1—C13—C14	-121.2 (3)
C11—Fe1—C7—C8	162.5 (2)	C15—Fe1—C13—C12	-82.1 (3)
C9—Fe1—C7—C8	-37.6 (2)	C10—Fe1—C13—C12	-159.8 (4)
C12—Fe1—C7—C8	119.2 (2)	C14—Fe1—C13—C12	-119.0 (4)
C13—Fe1—C7—C8	76.9 (3)	C6—Fe1—C13—C12	39.3 (7)
C15—Fe1—C7—C6	-40.4 (5)	C11—Fe1—C13—C12	-38.1 (3)
C10—Fe1—C7—C6	38.39 (19)	C9—Fe1—C13—C12	162.8 (3)
C14—Fe1—C7—C6	160.7 (5)	C7—Fe1—C13—C12	77.7 (3)
C11—Fe1—C7—C6	-77.8 (3)	C8—Fe1—C13—C12	119.8 (3)
C9—Fe1—C7—C6	82.1 (2)	C12—C13—C14—C15	0.2 (5)
C12—Fe1—C7—C6	-121.1 (2)	Fe1—C13—C14—C15	-59.0 (3)
C13—Fe1—C7—C6	-163.4 (2)	C12—C13—C14—Fe1	59.2 (3)
C8—Fe1—C7—C6	119.7 (3)	C10—Fe1—C14—C15	-77.8 (3)
C6—C7—C8—C9	-0.2 (4)	C6—Fe1—C14—C15	-44.1 (5)
Fe1—C7—C8—C9	58.6 (3)	C11—Fe1—C14—C15	38.3 (3)
C6—C7—C8—Fe1	-58.8 (2)	C9—Fe1—C14—C15	-120.1 (3)
C15—Fe1—C8—C9	39.2 (6)	C7—Fe1—C14—C15	167.7 (5)
C10—Fe1—C8—C9	-37.2 (2)	C12—Fe1—C14—C15	82.4 (3)
C14—Fe1—C8—C9	75.2 (3)	C13—Fe1—C14—C15	120.2 (4)
C6—Fe1—C8—C9	-81.6 (2)	C8—Fe1—C14—C15	-161.1 (3)
C11—Fe1—C8—C9	-164.4 (4)	C15—Fe1—C14—C13	-120.2 (4)
C7—Fe1—C8—C9	-119.3 (3)	C10—Fe1—C14—C13	162.0 (3)
C12—Fe1—C8—C9	159.6 (2)	C6—Fe1—C14—C13	-164.3 (3)
C13—Fe1—C8—C9	117.1 (2)	C11—Fe1—C14—C13	-82.0 (3)
C15—Fe1—C8—C7	158.6 (5)	C9—Fe1—C14—C13	119.7 (3)
C10—Fe1—C8—C7	82.2 (2)	C7—Fe1—C14—C13	47.5 (7)
C14—Fe1—C8—C7	-165.5 (3)	C12—Fe1—C14—C13	-37.8 (3)
C6—Fe1—C8—C7	37.8 (2)	C8—Fe1—C14—C13	78.6 (3)
C11—Fe1—C8—C7	-45.0 (6)	C13—C14—C15—C11	-0.7 (5)
C9—Fe1—C8—C7	119.3 (3)	Fe1—C14—C15—C11	-60.4 (3)
C12—Fe1—C8—C7	-81.0 (3)	C13—C14—C15—Fe1	59.7 (3)

C13—Fe1—C8—C7	-123.6 (2)	C12—C11—C15—C14	0.9 (5)
C7—C8—C9—C10	-0.1 (4)	Fe1—C11—C15—C14	60.6 (3)
Fe1—C8—C9—C10	58.7 (3)	C12—C11—C15—Fe1	-59.6 (3)
C7—C8—C9—Fe1	-58.8 (3)	C10—Fe1—C15—C14	118.6 (3)
C15—Fe1—C9—C10	75.2 (3)	C6—Fe1—C15—C14	160.5 (3)
C14—Fe1—C9—C10	116.3 (3)	C11—Fe1—C15—C14	-118.5 (4)
C6—Fe1—C9—C10	-38.2 (2)	C9—Fe1—C15—C14	77.9 (3)
C11—Fe1—C9—C10	44.4 (6)	C7—Fe1—C15—C14	-169.8 (4)
C7—Fe1—C9—C10	-82.5 (2)	C12—Fe1—C15—C14	-80.6 (3)
C12—Fe1—C9—C10	-171.5 (4)	C13—Fe1—C15—C14	-36.9 (3)
C13—Fe1—C9—C10	157.5 (2)	C8—Fe1—C15—C14	48.8 (6)
C8—Fe1—C9—C10	-120.1 (3)	C10—Fe1—C15—C11	-122.9 (3)
C15—Fe1—C9—C8	-164.7 (2)	C14—Fe1—C15—C11	118.5 (4)
C10—Fe1—C9—C8	120.1 (3)	C6—Fe1—C15—C11	-80.9 (3)
C14—Fe1—C9—C8	-123.5 (2)	C9—Fe1—C15—C11	-163.6 (3)
C6—Fe1—C9—C8	81.9 (2)	C7—Fe1—C15—C11	-51.3 (6)
C11—Fe1—C9—C8	164.5 (4)	C12—Fe1—C15—C11	37.9 (3)
C7—Fe1—C9—C8	37.6 (2)	C13—Fe1—C15—C11	81.7 (3)
C12—Fe1—C9—C8	-51.3 (5)	C8—Fe1—C15—C11	167.4 (4)
C13—Fe1—C9—C8	-82.4 (3)	C4—C3—C17—C18	126.9 (4)
C8—C9—C10—C6	0.4 (4)	C2—C3—C17—C18	-49.8 (5)
Fe1—C9—C10—C6	59.8 (2)	C4—C3—C17—C22	-51.3 (5)
C8—C9—C10—Fe1	-59.4 (3)	C2—C3—C17—C22	131.9 (4)
C7—C6—C10—C9	-0.5 (4)	C22—C17—C18—C19	1.1 (6)
C5—C6—C10—C9	-178.5 (3)	C3—C17—C18—C19	-177.2 (4)
Fe1—C6—C10—C9	-60.1 (2)	C17—C18—C19—C20	-0.2 (6)
C7—C6—C10—Fe1	59.6 (2)	C18—C19—C20—C21	-1.1 (6)
C5—C6—C10—Fe1	-118.4 (3)	C18—C19—C20—C11	178.5 (3)
C15—Fe1—C10—C9	-122.0 (3)	C19—C20—C21—C22	1.5 (6)
C14—Fe1—C10—C9	-81.3 (3)	C11—C20—C21—C22	-178.0 (3)
C6—Fe1—C10—C9	119.1 (3)	C20—C21—C22—C17	-0.7 (6)
C11—Fe1—C10—C9	-162.6 (3)	C18—C17—C22—C21	-0.6 (6)
C7—Fe1—C10—C9	80.8 (2)	C3—C17—C22—C21	177.7 (3)
C12—Fe1—C10—C9	170.4 (5)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N2-H2B\cdots N3^i$	0.86	2.28	3.047 (5)	149

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .

Fig. 1




Fig. 2

