

5,3'-Dihydroxy-7,4'-dimethoxyflavanone from *Artemisia sphaerocephala* Kraschen

Sumei Yao* and Weixia Qing

Medical College of Henan University, Henan University, Kaifeng 475004, People's Republic of China
Correspondence e-mail: ysum@yahoo.cn


Received 6 April 2008; accepted 6 April 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003 \text{ \AA}$; R factor = 0.029; wR factor = 0.078; data-to-parameter ratio = 7.5.

The title compound, $C_{17}H_{16}O_6$, was isolated from the Chinese Tibetan medicinal plant *Artemisia sphaerocephala* Kraschen. The molecular conformation is consolidated by two intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. A further intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond leads to chains along [010] in the crystal structure.

Related literature

For background, see: Zhao *et al.* (2007).



Experimental

Crystal data

$C_{17}H_{16}O_6$
 $M_r = 316.30$
Monoclinic, $P2_1$

$a = 5.4234(12) \text{ \AA}$
 $b = 9.293(2) \text{ \AA}$
 $c = 14.940(3) \text{ \AA}$

$\beta = 91.039(2)^\circ$
 $V = 752.9(3) \text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 0.11 \text{ mm}^{-1}$
 $T = 296(2) \text{ K}$
 $0.22 \times 0.18 \times 0.12 \text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2001)
 $T_{\min} = 0.977$, $T_{\max} = 0.987$

7958 measured reflections
1581 independent reflections
1489 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.077$
 $S = 1.06$
1581 reflections
212 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.11 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.14 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H3 \cdots O2	0.82	1.85	2.579 (2)	148
O6—H5 \cdots O5	0.82	2.18	2.640 (2)	116
O6—H5 \cdots O4 ⁱ	0.82	2.29	2.892 (2)	130

Symmetry code: (i) $-x, y + \frac{1}{2}, -z$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT-Plus* (Bruker, 2001); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *PLATON*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2716).

References

- Bruker (2001). *SAINT-Plus* (Version 6.45), *SMART* and *SADABS*. Bruker AXS, Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
Zhao, D. B., Li, L. X., Liu, X. H., Li, M. J. & Wang, W. L. (2007). *Chin. Chem. Lett.* **18**, 551–557.

supplementary materials

Acta Cryst. (2008). E64, o822 [doi:10.1107/S1600536808009355]

5,3'-Dihydroxy-7,4'-dimethoxyflavanone from *Artemisia sphaerocephala* Kraschen

S. Yao and W. Qing

Comment

As part of the ongoing investigations of the Chinese Tibetan medicinal plant *Artemisia sphaerocephala* Kraschen (Zhao *et al.*, 2007), we now report the isolation and structure of the flavone-related title compound, (I).

Compound (I) consists of three-ring system, including a phenyl ring and a benzopyrone fused ring (Fig. 1). The C–O bond distances range from 1.239 (2) to 1.453 (2) Å, in which C4–O2 [1.239 (2) Å] is typical for a C=O double-bond. The S(6) ring of O1/C2/C3/C4/C9/C10 in (I) is nonplanar, characterized by a O1–C2–C3–C4 torsion angle of 52.9 (2) °. Atom C2 is chiral, but the absolute structure of (I) could not be established from the present experiment. The dihedral angle between the aromatic ring planes is 77.26 (9) °.

Two intramolecular O-H···O hydrogen bonds (Table 1) help to establish the molecular conformation, both constructing S(6) rings. In addition, an intermolecular O-H···O link leads to [010] chains in the crystal (Fig. 2).

Experimental

The air-dried whole plant (5.1 kg) was ground into powder and extracted three times with 95% EtOH for 3 h each time. The concentrated extract was dispersed in water and partitioned successively with petroleum ether, CHCl₃, EtOAc and n-BuOH. The chloroform fraction (115 g) was subjected to silics gel with petroleum ether-acetone (9:1) to yield two fractions (Frs. 1–2). Fraction 2 (50 g) was subjected to silics gel with petroleum ether-EtOAc (50:1, 45:1, 40:1, 30:1) to yield six fractions. After a week, the crude title compound (20 mg) was crystallized from the fourth fraction. After recrystallization from petroleum ether-EtOAc, colourless blocks of (I) arose, with a melting point of 475 K. The molecular formula, C₁₇H₁₆O₆, was established by ESIMS m/z:316(M⁺). Spectroscopic analysis, ¹H NMR (400 MHz, DMSO-d₆) δ: 12.15 (1H, s), 6.08 (1H, d, J=2.2 Hz), 6.05 (1H, d, J=2.2 Hz), 5.47 (1H, dd, J=12.6 Hz, 3.0 Hz), 3.20 (1H, dd, J=15.5 Hz, 12.6 Hz), 2.82 (1H, dd, J=15.5 Hz, 3.0 Hz); ¹³C NMR (400 MHz, DMSO-d₆) δ: 197.0 (C-4), 168.0 (C-7), 163.8 (C-5), 163.1 (C-9), 147.8 (C-4'), 146.7 (C-3'), 131.8 (C-1'), 117.9 (C-6'), 113.5 (C-5'), 111.3 (C-2'), 102.9 (C-10), 94.6 (C-6), 93.7 (C-8), 79.0 (C-2), 42.6 (C-3), 55.4 (4'-OCH₃), 55.3 (7-OCH₃).

Refinement

Anomalous dispersion was negligible and Friedel pairs were merged before refinement.

The H atoms were geometrically placed (C–H = 0.93–0.98 Å, O—H = 0.82 Å), and refined as riding with U_{iso}(H)=1.2U_{eq}(C) or U_{iso}(H)=1.5U_{eq}(O or methyl C).

supplementary materials

Figures




Fig. 1. The molecular structure of (I), with displacement ellipsoids for the non-hydrogen atoms drawn at the 40% probability level. Hydrogen bonds are shown as double dashed lines.




Fig. 2. Fragment of the one-dimensional chain structure of (I) with hydrogen bonds shown as dashed lines.

5,3'-Dihydroxy-7,4'-dimethoxyflavanone

Crystal data

C ₁₇ H ₁₆ O ₆	$F_{000} = 332$
$M_r = 316.30$	$D_x = 1.395 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
Hall symbol: P 2y b	$\lambda = 0.71073 \text{ \AA}$
$a = 5.4234(12) \text{ \AA}$	Cell parameters from 4047 reflections
$b = 9.293(2) \text{ \AA}$	$\theta = 2.6\text{--}27.5^\circ$
$c = 14.940(3) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$\beta = 91.039(2)^\circ$	$T = 296(2) \text{ K}$
$V = 752.9(3) \text{ \AA}^3$	Block, colorless
$Z = 2$	$0.22 \times 0.18 \times 0.12 \text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer	1581 independent reflections
Radiation source: fine-focus sealed tube	1489 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.019$
$T = 296(2) \text{ K}$	$\theta_{\text{max}} = 26.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.6^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$h = -6 \rightarrow 6$
$T_{\text{min}} = 0.977, T_{\text{max}} = 0.987$	$k = -11 \rightarrow 11$
7958 measured reflections	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H-atom parameters constrained
$wR(F^2) = 0.077$	$w = 1/[\sigma^2(F_o^2) + (0.0485P)^2 + 0.0531P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\max} < 0.001$
1581 reflections	$\Delta\rho_{\max} = 0.11 \text{ e } \text{\AA}^{-3}$
212 parameters	$\Delta\rho_{\min} = -0.14 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.3993 (2)	0.73089 (16)	0.11833 (7)	0.0443 (3)
O2	0.8292 (3)	0.9824 (2)	0.28305 (11)	0.0697 (5)
O3	0.6055 (3)	0.8761 (2)	0.41875 (10)	0.0703 (5)
H3	0.7086	0.9231	0.3919	0.105*
O4	-0.0461 (3)	0.54437 (19)	0.35407 (9)	0.0532 (4)
O5	0.5698 (3)	0.79981 (18)	-0.28802 (9)	0.0582 (4)
O6	0.2587 (3)	0.94212 (19)	-0.18692 (9)	0.0595 (4)
H5	0.2732	0.9377	-0.2414	0.089*
C1'	0.6333 (3)	0.7757 (2)	-0.01135 (12)	0.0396 (4)
C2	0.6447 (3)	0.7700 (2)	0.08935 (12)	0.0407 (4)
H2	0.7605	0.6942	0.1079	0.049*
C2'	0.8030 (4)	0.7024 (2)	-0.06221 (12)	0.0469 (5)
H2'	0.9271	0.6494	-0.0338	0.056*
C3	0.7244 (4)	0.9111 (3)	0.13314 (13)	0.0509 (5)
H3A	0.8946	0.9310	0.1185	0.061*
H3B	0.6235	0.9889	0.1094	0.061*
C3'	0.7904 (4)	0.7070 (2)	-0.15510 (13)	0.0508 (5)
H3'	0.9056	0.6574	-0.1886	0.061*
C4	0.7001 (4)	0.9053 (2)	0.23342 (13)	0.0485 (5)
C4'	0.6070 (4)	0.7852 (2)	-0.19750 (12)	0.0434 (4)
C5	0.4695 (4)	0.7986 (2)	0.35954 (12)	0.0472 (5)

supplementary materials

C5'	0.4376 (3)	0.8613 (2)	-0.14645 (12)	0.0415 (4)
C6	0.2845 (4)	0.7110 (3)	0.39144 (12)	0.0490 (5)
H6	0.2560	0.7047	0.4525	0.059*
C6'	0.4503 (3)	0.8559 (2)	-0.05446 (12)	0.0439 (4)
H6'	0.3359	0.9062	-0.0210	0.053*
C7	0.1420 (3)	0.6325 (2)	0.33055 (12)	0.0416 (4)
C7'	0.7267 (6)	0.7210 (3)	-0.34504 (14)	0.0701 (7)
H7'A	0.8943	0.7509	-0.3350	0.105*
H7'B	0.6798	0.7386	-0.4063	0.105*
H7'C	0.7120	0.6201	-0.3324	0.105*
C8	0.1841 (3)	0.6390 (2)	0.23879 (12)	0.0410 (4)
H8	0.0882	0.5852	0.1989	0.049*
C9	0.3691 (3)	0.7260 (2)	0.20815 (11)	0.0373 (4)
C10	0.5161 (3)	0.8091 (2)	0.26731 (12)	0.0413 (4)
C11	-0.1037 (5)	0.5297 (3)	0.44657 (14)	0.0660 (7)
H11A	0.0353	0.4893	0.4784	0.099*
H11B	-0.2436	0.4673	0.4525	0.099*
H11C	-0.1413	0.6225	0.4710	0.099*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0462 (7)	0.0580 (8)	0.0285 (6)	-0.0111 (6)	0.0003 (5)	0.0007 (6)
O2	0.0826 (11)	0.0731 (11)	0.0531 (9)	-0.0365 (10)	-0.0075 (8)	-0.0084 (8)
O3	0.0893 (12)	0.0821 (12)	0.0389 (7)	-0.0298 (10)	-0.0109 (7)	-0.0116 (8)
O4	0.0558 (8)	0.0709 (10)	0.0330 (7)	-0.0121 (7)	0.0052 (6)	0.0056 (7)
O5	0.0790 (10)	0.0625 (9)	0.0332 (7)	0.0170 (8)	0.0059 (6)	-0.0006 (7)
O6	0.0639 (9)	0.0755 (11)	0.0391 (8)	0.0270 (8)	-0.0006 (7)	0.0045 (8)
C1'	0.0429 (9)	0.0412 (10)	0.0346 (9)	-0.0019 (8)	0.0026 (7)	0.0034 (8)
C2	0.0407 (9)	0.0457 (10)	0.0357 (9)	-0.0013 (8)	-0.0004 (7)	0.0048 (8)
C2'	0.0465 (10)	0.0504 (11)	0.0438 (10)	0.0109 (9)	0.0036 (8)	0.0073 (9)
C3	0.0557 (11)	0.0530 (12)	0.0440 (11)	-0.0138 (10)	0.0009 (9)	0.0023 (9)
C3'	0.0552 (11)	0.0531 (12)	0.0446 (11)	0.0140 (10)	0.0117 (9)	0.0014 (10)
C4	0.0525 (11)	0.0484 (11)	0.0443 (10)	-0.0106 (10)	-0.0054 (9)	-0.0026 (9)
C4'	0.0561 (10)	0.0409 (10)	0.0334 (9)	0.0030 (9)	0.0069 (8)	0.0024 (8)
C5	0.0568 (11)	0.0521 (12)	0.0326 (9)	-0.0008 (10)	-0.0075 (8)	-0.0051 (8)
C5'	0.0443 (9)	0.0430 (10)	0.0372 (9)	0.0055 (8)	0.0010 (8)	0.0040 (8)
C6	0.0597 (11)	0.0605 (12)	0.0267 (8)	-0.0003 (10)	0.0001 (8)	-0.0005 (9)
C6'	0.0465 (10)	0.0484 (10)	0.0370 (9)	0.0089 (9)	0.0075 (8)	0.0013 (8)
C7	0.0422 (10)	0.0491 (11)	0.0335 (9)	0.0010 (8)	0.0014 (7)	0.0047 (8)
C7'	0.1016 (19)	0.0707 (16)	0.0385 (11)	0.0154 (15)	0.0164 (11)	-0.0066 (12)
C8	0.0409 (9)	0.0494 (10)	0.0326 (9)	-0.0038 (8)	-0.0039 (7)	-0.0016 (8)
C9	0.0405 (9)	0.0424 (10)	0.0290 (8)	0.0004 (8)	-0.0028 (7)	0.0007 (8)
C10	0.0450 (9)	0.0443 (10)	0.0346 (9)	-0.0016 (8)	-0.0031 (7)	-0.0011 (8)
C11	0.0706 (15)	0.0886 (18)	0.0393 (11)	-0.0062 (14)	0.0133 (10)	0.0150 (12)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.3555 (19)	C3'—C4'	1.377 (3)
-------	-------------	---------	-----------

O1—C2	1.453 (2)	C3'—H3'	0.9300
O2—C4	1.239 (2)	C4—C10	1.439 (3)
O3—C5	1.350 (2)	C4'—C5'	1.397 (3)
O3—H3	0.8200	C5—C6	1.383 (3)
O4—C7	1.359 (2)	C5—C10	1.409 (3)
O4—C11	1.429 (2)	C5'—C6'	1.376 (2)
O5—C4'	1.370 (2)	C6—C7	1.390 (3)
O5—C7'	1.419 (3)	C6—H6	0.9300
O6—C5'	1.360 (2)	C6'—H6'	0.9300
O6—H5	0.8200	C7—C8	1.395 (2)
C1'—C2'	1.383 (3)	C7'—H7'A	0.9600
C1'—C6'	1.390 (3)	C7'—H7'B	0.9600
C1'—C2	1.506 (2)	C7'—H7'C	0.9600
C2—C3	1.525 (3)	C8—C9	1.373 (3)
C2—H2	0.9800	C8—H8	0.9300
C2'—C3'	1.389 (3)	C9—C10	1.409 (2)
C2'—H2'	0.9300	C11—H11A	0.9600
C3—C4	1.507 (3)	C11—H11B	0.9600
C3—H3A	0.9700	C11—H11C	0.9600
C3—H3B	0.9700		
C9—O1—C2	115.53 (13)	C6—C5—C10	121.46 (17)
C5—O3—H3	109.5	O6—C5'—C6'	119.12 (16)
C7—O4—C11	119.06 (17)	O6—C5'—C4'	120.52 (16)
C4'—O5—C7'	117.54 (18)	C6'—C5'—C4'	120.37 (17)
C5'—O6—H5	109.5	C5—C6—C7	118.79 (17)
C2'—C1'—C6'	119.09 (16)	C5—C6—H6	120.6
C2'—C1'—C2	121.06 (16)	C7—C6—H6	120.6
C6'—C1'—C2	119.86 (16)	C5'—C6'—C1'	120.31 (17)
O1—C2—C1'	106.57 (13)	C5'—C6'—H6'	119.8
O1—C2—C3	109.97 (16)	C1'—C6'—H6'	119.8
C1'—C2—C3	113.88 (16)	O4—C7—C6	123.92 (16)
O1—C2—H2	108.8	O4—C7—C8	114.67 (16)
C1'—C2—H2	108.8	C6—C7—C8	121.40 (17)
C3—C2—H2	108.8	O5—C7'—H7'A	109.5
C1'—C2'—C3'	120.83 (17)	O5—C7'—H7'B	109.5
C1'—C2'—H2'	119.6	H7'A—C7'—H7'B	109.5
C3'—C2'—H2'	119.6	O5—C7'—H7'C	109.5
C4—C3—C2	111.48 (16)	H7'A—C7'—H7'C	109.5
C4—C3—H3A	109.3	H7'B—C7'—H7'C	109.5
C2—C3—H3A	109.3	C9—C8—C7	119.13 (17)
C4—C3—H3B	109.3	C9—C8—H8	120.4
C2—C3—H3B	109.3	C7—C8—H8	120.4
H3A—C3—H3B	108.0	O1—C9—C8	116.84 (16)
C4'—C3'—C2'	119.86 (17)	O1—C9—C10	121.70 (16)
C4'—C3'—H3'	120.1	C8—C9—C10	121.44 (15)
C2'—C3'—H3'	120.1	C5—C10—C9	117.77 (17)
O2—C4—C10	122.49 (18)	C5—C10—C4	121.68 (17)
O2—C4—C3	121.02 (19)	C9—C10—C4	120.51 (16)
C10—C4—C3	116.47 (17)	O4—C11—H11A	109.5

supplementary materials


O5—C4'—C3'	126.74 (17)	O4—C11—H11B	109.5
O5—C4'—C5'	113.72 (17)	H11A—C11—H11B	109.5
C3'—C4'—C5'	119.53 (16)	O4—C11—H11C	109.5
O3—C5—C6	118.64 (17)	H11A—C11—H11C	109.5
O3—C5—C10	119.90 (18)	H11B—C11—H11C	109.5
C9—O1—C2—C1'	−177.36 (16)	C2'—C1'—C6'—C5'	0.5 (3)
C9—O1—C2—C3	−53.5 (2)	C2—C1'—C6'—C5'	−179.73 (18)
C2'—C1'—C2—O1	−128.91 (19)	C11—O4—C7—C6	−0.2 (3)
C6'—C1'—C2—O1	51.3 (2)	C11—O4—C7—C8	179.76 (19)
C2'—C1'—C2—C3	109.7 (2)	C5—C6—C7—O4	−179.2 (2)
C6'—C1'—C2—C3	−70.1 (2)	C5—C6—C7—C8	0.8 (3)
C6'—C1'—C2'—C3'	−0.8 (3)	O4—C7—C8—C9	179.50 (18)
C2—C1'—C2'—C3'	179.46 (18)	C6—C7—C8—C9	−0.6 (3)
O1—C2—C3—C4	52.9 (2)	C2—O1—C9—C8	−155.13 (17)
C1'—C2—C3—C4	172.44 (16)	C2—O1—C9—C10	26.5 (3)
C1'—C2'—C3'—C4'	0.0 (3)	C7—C8—C9—O1	−178.69 (17)
C2—C3—C4—O2	154.3 (2)	C7—C8—C9—C10	−0.3 (3)
C2—C3—C4—C10	−27.3 (3)	O3—C5—C10—C9	180.00 (19)
C7'—O5—C4'—C3'	3.6 (3)	C6—C5—C10—C9	−0.6 (3)
C7'—O5—C4'—C5'	−177.3 (2)	O3—C5—C10—C4	−2.5 (3)
C2'—C3'—C4'—O5	−179.8 (2)	C6—C5—C10—C4	176.9 (2)
C2'—C3'—C4'—C5'	1.1 (3)	O1—C9—C10—C5	179.19 (17)
O5—C4'—C5'—O6	−0.8 (3)	C8—C9—C10—C5	0.9 (3)
C3'—C4'—C5'—O6	178.4 (2)	O1—C9—C10—C4	1.6 (3)
O5—C4'—C5'—C6'	179.39 (18)	C8—C9—C10—C4	−176.67 (19)
C3'—C4'—C5'—C6'	−1.4 (3)	O2—C4—C10—C5	0.9 (3)
O3—C5—C6—C7	179.2 (2)	C3—C4—C10—C5	−177.44 (19)
C10—C5—C6—C7	−0.2 (3)	O2—C4—C10—C9	178.4 (2)
O6—C5'—C6'—C1'	−179.19 (19)	C3—C4—C10—C9	0.0 (3)
C4'—C5'—C6'—C1'	0.6 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O3—H3—O2	0.82	1.85	2.579 (2)	148
O6—H5—O5	0.82	2.18	2.640 (2)	116
O6—H5—O4 ⁱ	0.82	2.29	2.892 (2)	130

Symmetry codes: (i) $-x, y+1/2, -z$.

Fig. 1



supplementary materials

Fig. 2

