

Poly[penta- μ -aqua- μ_6 -methylene-disulfonato- μ_5 -methylenedisulfonato-tetrasodium(I)]

Dan-Dan Cao and Zai-Chao Zhang*

Department of Chemistry, Huaiyin Teachers College, 111 West Changjiang Road, Huaian 223300, Jiangsu, People's Republic of China
Correspondence e-mail: overloadzz@hotmail.com


Received 1 May 2008; accepted 10 May 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(S-C) = 0.002$ Å; R factor = 0.031; wR factor = 0.093; data-to-parameter ratio = 13.6.

The title compound, $[\text{Na}_4(\text{CH}_2\text{O}_6\text{S}_2)_2(\text{H}_2\text{O})_5]_n$, was crystallized from an aqueous solution. The sodium ions are surrounded and bridged by O atoms from coordinated water molecules and sulfonate ions in a three-dimensional neutral network. The crystal structure is also stabilized by an intricate system of hydrogen bonds.

Related literature

The supramolecular chemistry of the sulfonate group in extended solids constructed by cooperative coordination and other weak intermolecular interactions, as well as the structural and functional properties of Ba^{2+} and Ag^+ sulfonates, has been reviewed by Côté & Shimizu (2003). For a review of the structural chemistry and properties of metal arenesulfonates, see: Cai (2004). For related literature, see: Li *et al.* (2008); Mi *et al.* (2007); Videnova-Adrabsinska (2007).



Experimental

Crystal data

$[\text{Na}_4(\text{CH}_2\text{O}_6\text{S}_2)_2(\text{H}_2\text{O})_5]$	$\gamma = 87.227$ (2)°
$M_r = 530.33$	$V = 860.19$ (11) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.7758$ (7) Å	Mo $K\alpha$ radiation
$b = 9.5339$ (7) Å	$\mu = 0.74$ mm ⁻¹
$c = 10.7878$ (8) Å	$T = 296$ (2) K
$\alpha = 81.425$ (2)°	$0.30 \times 0.25 \times 0.25$ mm
$\beta = 74.545$ (2)°	

Data collection

Bruker SMART APEX2 diffractometer	10702 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2000)	3321 independent reflections
$T_{\min} = 0.81$, $T_{\max} = 0.84$	3050 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	244 parameters
$wR(F^2) = 0.093$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.47$ e Å ⁻³
3321 reflections	$\Delta\rho_{\text{min}} = -0.56$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O13—H13A \cdots O8 ⁱ	0.97	1.93	2.868 (2)	163
O13—H13B \cdots O5 ⁱⁱ	0.97	1.87	2.824 (2)	166
O14—H14A \cdots O9 ⁱⁱⁱ	0.97	2.00	2.871 (2)	148
O15—H15B \cdots O3 ^{iv}	0.97	1.97	2.877 (2)	155
O16—H16A \cdots O9 ⁱⁱⁱ	0.97	1.87	2.794 (2)	158
O16—H16B \cdots O12	0.97	2.18	3.034 (3)	147
O17—H17A \cdots O4 ^v	0.97	2.17	2.879 (3)	129
O17—H17B \cdots O12 ^{vi}	0.97	2.10	2.918 (3)	141

Symmetry codes: (i) $x-1, y, z$; (ii) $x, y-1, z$; (iii) $-x+2, -y+1, -z+1$; (iv) $x+1, y, z$; (v) $-x, -y+2, -z+2$; (vi) $x-1, y+1, z$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2003).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2114).

References

- Bruker (2000). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Cai, J.-W. (2004). *Coord. Chem. Rev.* **248**, 1061–1083.
Côté, A. P. & Shimizu, G. K. H. (2003). *Coord. Chem. Rev.* **245**, 49–60.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Li, M., Xiang, J.-F., Chen, S.-P., Wu, S.-M., Yuan, L.-J., Li, H., He, H.-J. & Sun, J.-T. (2008). *J. Coord. Chem.* **61**, 372–383.
Mi, L.-W., Hou, H.-W., Song, Z.-Y., Han, H.-Y., Xu, H., Fan, Y.-T. & Ng, S.-W. (2007). *J. Cryst. Growth Des.* **7**, 2553–2561.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
Videnova-Adrabsinska, V. (2007). *Coord. Chem. Rev.* **251**, 1987–2016.

supplementary materials

Acta Cryst. (2008). E64, m815 [doi:10.1107/S1600536808014037]

Poly[penta- μ -aqua- μ_6 -methylenedisulfonato- μ_5 -methylenedisulfonato-tetrasodium(I)]

D.-D. Cao and Z.-C. Zhang

Comment

Due to the weak coordination strength of monosulfonate ions, most metal complexes of these ligands obtained from aqueous solution are water-coordinated metal sulfonate salts, and the coordination chemistry of the sulfonate ion has been less well investigated in comparison with other organic acidato anions such as carbonates and phosphonates (Côté & Shimizu, 2003). However, by employing disulfonates, which can provide multiple potentially chelating coordination sites, stable networks sustained by sulfonate-metal interactions can be obtained with various dimensionalities (Cai, 2004; Li *et al.*, 2008; Mi *et al.*, 2007; Videnova-Adrabinska, 2007).

The present structural study of the title compound $\{\text{Na}_4(\text{mds})_2(\text{H}_2\text{O})_5\}_n$ (I) reveals the existence of sulfonate-sodium interactions. The asymmetric unit consists of four sodium ions, five coordinated μ_2 -H₂O molecules and two CH₃(SO₃)₂²⁻ ligands (Fig. 1). Of the four sodium ions, Na1, Na2 and Na4 are six-coordinated. Na3 is coordinated by five oxygen atoms (O2^{vi}, O13, O14, O16 and O16^{iv}). The Na—O bond lengths fall in the range of 2.334 (2) to 2.5609 (18) Å. All sulfonate oxygens are deprotonated. Three sulfonate oxygen atoms (O5, O9 and O12) are not coordinated to sodium atoms and do instead act as hydrogen bonding acceptors towards coordinated water molecules. All other hydrogen bonds are from water molecules towards Na coordinated sulfonate oxygen atoms (Table 1).

Each sodium atom is bridged to an equivalent neighbour, forming $\{\text{Na}_2\text{O}_2\}$ dimers with inversion centers in the middle. O1 and O7 connect the dimers of Na1 and Na2 into an infinite chain. Na1-Na2 chains are joined to dimers of Na3 and Na4 through intricate bridges of oxygens from sulfonate groups and coordinated waters, and O—S—O connectivities, thus generating a three-dimensional network (Fig. 2).

Experimental

An 1 mol L⁻¹ NaOH aqueous solution was dropped into one of mdsH₂ (0.14 g, 1 mmol in 20 ml H₂O) until the pH value of the solution reached 7 to 9. Slow evaporation for several days yielded colorless prismatic crystals of the title compound. Anal. Calcd for C₂H₁₄O₁₇Na₄S₄ (530.34): C 4.51, H 2.69%; Found: C 4.53, H 2.66%.

Refinement

All the non-hydrogen atoms were located from the Fourier maps, and were refined anisotropically. H atoms from μ_2 -H₂O can be suitably placed in calculated positions, so all the H atoms, including those attached to carbon atoms and from water molecules, were positioned geometrically, and the isotropic vibration parameters related to the atoms which they are bonded to with $U_{\text{iso}} = 1.2 U_{\text{eq}}$.

Figures




Fig. 1. The asymmetric unit of I with atom labels and 50% probability displacement ellipsoids for non-H atoms. H atoms attached to carbon atoms are omitted for clarity. Symmetry codes: (i) $-1+x, y, z$; (ii) $1+x, y, z$; (iii) $-x, 2-y, 2-z$. (iv) $1-x, 1-y, 1-z$; (v) $1-x, 1-y, 2-z$; (vi) $1-x, 2-y, 1-z$; (vii) $1-x, 2-y, 2-z$; (viii) $2-x, 2-y, 1-z$;




Fig. 2. Packing diagram of compound I. Na—O bond is showed by black line, H atoms are omitted for clarity.

Poly[penta- μ -aqua- μ_6 -methylenedisulfonato- μ_5 -methylenedisulfonato- tetrasodium(I)]

Crystal data

$[\text{Na}_4(\text{CH}_2\text{O}_6\text{S}_2)_2(\text{H}_2\text{O})_5]$

$M_r = 530.33$

Triclinic, PT

Hall symbol: $-P\ 1$

$a = 8.7758\ (7)\ \text{\AA}$

$b = 9.5339\ (7)\ \text{\AA}$

$c = 10.7878\ (8)\ \text{\AA}$

$\alpha = 81.425\ (2)^\circ$

$\beta = 74.545\ (2)^\circ$

$\gamma = 87.227\ (2)^\circ$

$V = 860.19\ (11)\ \text{\AA}^3$

$Z = 2$

$F_{000} = 540$

$D_x = 2.048\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 202 reflections

$\theta = 2.6\text{--}21.1^\circ$

$\mu = 0.74\ \text{mm}^{-1}$

$T = 296\ (2)\ \text{K}$

Prismlike, colourless

$0.30 \times 0.25 \times 0.25\ \text{mm}$

Data collection

Bruker SMART APEX2
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 296\ (2)\ \text{K}$

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2000)

$T_{\min} = 0.81, T_{\max} = 0.84$

10702 measured reflections

3321 independent reflections

3050 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\text{max}} = 26.0^\circ$

$\theta_{\text{min}} = 2.0^\circ$

$h = -10 \rightarrow 10$

$k = -11 \rightarrow 11$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.030$	$w = 1/[\sigma^2(F_o^2) + (0.0428P)^2 + 0.7023P]$
$wR(F^2) = 0.093$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\max} = 0.001$
3321 reflections	$\Delta\rho_{\max} = 0.47 \text{ e } \text{\AA}^{-3}$
244 parameters	$\Delta\rho_{\min} = -0.56 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F^* = F(1 + 0.002x F^2(\sin(2\theta))^{-0.25})$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.00025 (5)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4436 (2)	1.0561 (2)	0.8451 (2)	0.0189 (4)
H1A	0.4297	0.9743	0.9121	0.023*
H1B	0.5440	1.0983	0.8394	0.023*
C2	0.9094 (3)	0.4391 (2)	0.8842 (2)	0.0217 (4)
H2A	1.0010	0.3786	0.8863	0.026*
H2B	0.8798	0.4796	0.9647	0.026*
Na1	0.55562 (10)	0.66480 (9)	0.86815 (9)	0.0253 (2)
Na2	0.85901 (11)	0.87496 (9)	0.59357 (9)	0.0268 (2)
Na3	0.54875 (13)	0.67712 (11)	0.52620 (12)	0.0421 (3)
Na4	0.08149 (10)	0.88263 (9)	0.88189 (8)	0.0240 (2)
O1	0.57149 (19)	0.87897 (16)	0.69116 (15)	0.0261 (4)
O2	0.5103 (2)	1.11297 (17)	0.59433 (15)	0.0278 (4)
O3	0.29960 (18)	0.94639 (19)	0.70181 (15)	0.0287 (4)
O4	0.3169 (2)	1.21026 (19)	1.01825 (16)	0.0334 (4)
O5	0.3122 (2)	1.30241 (17)	0.79628 (18)	0.0353 (4)
O6	0.13991 (19)	1.11171 (17)	0.91904 (17)	0.0290 (4)

supplementary materials

O7	0.82527 (18)	0.65404 (17)	0.73482 (15)	0.0259 (4)
O8	1.06688 (19)	0.66657 (17)	0.80084 (17)	0.0286 (4)
O9	1.0542 (2)	0.51641 (18)	0.64174 (16)	0.0325 (4)
O10	0.6110 (2)	0.41828 (18)	0.9018 (2)	0.0366 (4)
O11	0.7435 (2)	0.21996 (18)	0.99171 (18)	0.0352 (4)
O12	0.7916 (2)	0.27615 (19)	0.75861 (17)	0.0367 (4)
O13	0.3962 (2)	0.59164 (17)	0.74448 (17)	0.0316 (4)
H13A	0.2921	0.6356	0.7642	0.038*
H13B	0.3854	0.4893	0.7581	0.038*
O14	0.8268 (2)	0.7545 (2)	0.42866 (17)	0.0361 (4)
H14A	0.8993	0.6746	0.4159	0.043*
H14B	0.8388	0.8170	0.3470	0.043*
O15	1.1401 (2)	0.88559 (19)	0.51526 (17)	0.0338 (4)
H15A	1.1832	0.8177	0.4560	0.041*
H15B	1.1927	0.8759	0.5846	0.041*
O16	0.6570 (2)	0.43211 (19)	0.54436 (17)	0.0355 (4)
H16A	0.7614	0.4243	0.4860	0.043*
H16B	0.6587	0.3945	0.6326	0.043*
O17	-0.1158 (2)	0.97729 (18)	0.77543 (16)	0.0308 (4)
H17A	-0.2173	0.9697	0.8396	0.037*
H17B	-0.0951	1.0776	0.7471	0.037*
S1	0.45658 (6)	0.99467 (5)	0.69417 (5)	0.01723 (13)
S2	0.29091 (6)	1.18145 (5)	0.89699 (5)	0.02035 (14)
S3	0.96766 (6)	0.58056 (5)	0.75349 (5)	0.01822 (14)
S4	0.75141 (6)	0.33016 (5)	0.88177 (5)	0.01945 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0163 (10)	0.0213 (10)	0.0196 (9)	-0.0008 (8)	-0.0062 (8)	-0.0015 (8)
C2	0.0201 (11)	0.0206 (10)	0.0249 (10)	-0.0059 (8)	-0.0097 (9)	0.0031 (8)
Na1	0.0236 (5)	0.0240 (5)	0.0298 (5)	-0.0020 (3)	-0.0086 (4)	-0.0049 (4)
Na2	0.0269 (5)	0.0246 (5)	0.0278 (5)	-0.0050 (4)	-0.0061 (4)	-0.0010 (4)
Na3	0.0352 (6)	0.0290 (5)	0.0592 (7)	-0.0041 (4)	-0.0151 (5)	0.0090 (5)
Na4	0.0222 (5)	0.0225 (4)	0.0275 (4)	-0.0026 (3)	-0.0064 (4)	-0.0035 (3)
O1	0.0236 (8)	0.0227 (8)	0.0298 (8)	0.0048 (6)	-0.0047 (6)	-0.0026 (6)
O2	0.0333 (9)	0.0226 (8)	0.0217 (8)	0.0005 (7)	-0.0009 (7)	0.0034 (6)
O3	0.0179 (8)	0.0449 (10)	0.0253 (8)	-0.0059 (7)	-0.0055 (6)	-0.0102 (7)
O4	0.0336 (10)	0.0408 (10)	0.0289 (9)	-0.0089 (8)	-0.0056 (7)	-0.0161 (8)
O5	0.0451 (11)	0.0193 (8)	0.0410 (10)	0.0027 (7)	-0.0139 (8)	0.0013 (7)
O6	0.0177 (8)	0.0302 (9)	0.0412 (9)	-0.0024 (7)	-0.0055 (7)	-0.0149 (7)
O7	0.0199 (8)	0.0258 (8)	0.0296 (8)	0.0012 (6)	-0.0064 (6)	0.0031 (6)
O8	0.0241 (8)	0.0224 (8)	0.0401 (9)	-0.0078 (6)	-0.0080 (7)	-0.0050 (7)
O9	0.0320 (9)	0.0296 (9)	0.0299 (9)	-0.0016 (7)	0.0049 (7)	-0.0088 (7)
O10	0.0195 (8)	0.0249 (9)	0.0626 (12)	-0.0013 (7)	-0.0085 (8)	-0.0010 (8)
O11	0.0371 (10)	0.0292 (9)	0.0376 (9)	-0.0137 (8)	-0.0151 (8)	0.0151 (7)
O12	0.0471 (11)	0.0332 (10)	0.0309 (9)	-0.0113 (8)	-0.0073 (8)	-0.0097 (7)
O13	0.0273 (9)	0.0237 (8)	0.0458 (10)	-0.0008 (7)	-0.0128 (8)	-0.0051 (7)

O14	0.0411 (11)	0.0366 (10)	0.0339 (9)	0.0084 (8)	-0.0144 (8)	-0.0096 (8)
O15	0.0280 (9)	0.0371 (10)	0.0331 (9)	0.0018 (7)	-0.0063 (7)	0.0011 (7)
O16	0.0296 (9)	0.0405 (10)	0.0330 (9)	-0.0025 (8)	-0.0060 (7)	0.0020 (8)
O17	0.0318 (9)	0.0290 (9)	0.0328 (9)	-0.0004 (7)	-0.0097 (7)	-0.0061 (7)
S1	0.0149 (3)	0.0180 (3)	0.0175 (2)	-0.00013 (19)	-0.00291 (19)	-0.00080 (19)
S2	0.0192 (3)	0.0183 (3)	0.0247 (3)	-0.0022 (2)	-0.0057 (2)	-0.0062 (2)
S3	0.0165 (3)	0.0157 (3)	0.0210 (3)	-0.00201 (19)	-0.00272 (19)	-0.00110 (19)
S4	0.0192 (3)	0.0153 (3)	0.0240 (3)	-0.00369 (19)	-0.0073 (2)	0.00080 (19)

Geometric parameters (Å, °)

C1—S2	1.783 (2)	Na4—O17	2.3942 (19)
C1—S1	1.785 (2)	Na4—O11 ⁱⁱ	2.3972 (19)
C1—H1A	0.9700	Na4—O6 ^{viii}	2.4881 (18)
C1—H1B	0.9700	Na4—Na4 ^{viii}	3.6050 (17)
C2—S4	1.780 (2)	Na4—Na2 ^{vii}	4.1000 (13)
C2—S3	1.785 (2)	O1—S1	1.4562 (16)
C2—H2A	0.9700	O2—S1	1.4398 (16)
C2—H2B	0.9700	O2—Na3 ^v	2.3339 (18)
Na1—O4 ⁱ	2.3382 (19)	O3—S1	1.4512 (16)
Na1—O13	2.3644 (19)	O4—S2	1.4528 (17)
Na1—O10	2.3709 (19)	O4—Na1 ⁱ	2.3382 (19)
Na1—O7	2.4260 (18)	O5—S2	1.4437 (18)
Na1—O10 ⁱⁱ	2.552 (2)	O6—S2	1.4577 (17)
Na1—O1	2.5607 (18)	O6—Na4 ^{viii}	2.4881 (18)
Na1—O11 ⁱⁱ	2.913 (2)	O7—S3	1.4492 (16)
Na1—S4 ⁱⁱ	3.2699 (10)	O8—S3	1.4569 (16)
Na1—Na3	3.6898 (15)	O8—Na4 ⁱⁱⁱ	2.3763 (18)
Na1—Na2	3.7975 (13)	O9—S3	1.4460 (16)
Na1—Na1 ⁱⁱ	3.8822 (18)	O10—S4	1.4432 (17)
Na2—O14	2.3389 (19)	O10—Na1 ⁱⁱ	2.552 (2)
Na2—O17 ⁱⁱⁱ	2.3816 (19)	O11—S4	1.4515 (17)
Na2—O15	2.3866 (19)	O11—Na4 ⁱⁱ	2.3972 (19)
Na2—O7	2.3893 (18)	O11—Na1 ⁱⁱ	2.913 (2)
Na2—O15 ^{iv}	2.4057 (19)	O12—S4	1.4462 (17)
Na2—O1	2.4623 (18)	O13—H13A	0.9700
Na2—Na2 ^{iv}	3.5152 (18)	O13—H13B	0.9700
Na2—Na3	3.6688 (15)	O14—H14A	0.9700
Na2—Na4 ⁱⁱⁱ	4.1000 (13)	O14—H14B	0.9700
Na3—O2 ^v	2.3339 (18)	O15—Na2 ^{iv}	2.4057 (19)
Na3—O13	2.420 (2)	O15—H15A	0.9700
Na3—O16 ^{vi}	2.465 (2)	O15—H15B	0.9700
Na3—O16	2.479 (2)	O16—Na3 ^{vi}	2.465 (2)
Na3—O14	2.486 (2)	O16—H16A	0.9700

supplementary materials

Na3—O1	2.856 (2)	O16—H16B	0.9700
Na3—Na3 ^{vi}	3.683 (2)	O17—Na2 ^{vii}	2.3816 (19)
Na4—O3	2.3635 (18)	O17—H17A	0.9700
Na4—O8 ^{vii}	2.3763 (18)	O17—H17B	0.9700
Na4—O6	2.3816 (18)	S4—Na1 ⁱⁱ	3.2699 (10)
S2—C1—S1	117.32 (11)	O13—Na3—Na2	100.89 (6)
S2—C1—H1A	108.0	O16 ^{vi} —Na3—Na2	172.47 (6)
S1—C1—H1A	108.0	O16—Na3—Na2	101.22 (6)
S2—C1—H1B	108.0	O14—Na3—Na2	39.03 (5)
S1—C1—H1B	108.0	O1—Na3—Na2	42.08 (4)
H1A—C1—H1B	107.2	O2 ^v —Na3—Na3 ^{vi}	123.29 (7)
S4—C2—S3	117.32 (12)	O13—Na3—Na3 ^{vi}	81.18 (5)
S4—C2—H2A	108.0	O16 ^{vi} —Na3—Na3 ^{vi}	41.99 (5)
S3—C2—H2A	108.0	O16—Na3—Na3 ^{vi}	41.70 (5)
S4—C2—H2B	108.0	O14—Na3—Na3 ^{vi}	116.43 (7)
S3—C2—H2B	108.0	O1—Na3—Na3 ^{vi}	151.94 (6)
H2A—C2—H2B	107.2	Na2—Na3—Na3 ^{vi}	142.60 (5)
O4 ⁱ —Na1—O13	166.18 (8)	O2 ^v —Na3—Na1	121.20 (6)
O4 ⁱ —Na1—O10	111.23 (8)	O13—Na3—Na1	38.99 (5)
O13—Na1—O10	82.42 (7)	O16 ^{vi} —Na3—Na1	124.37 (6)
O4 ⁱ —Na1—O7	80.23 (6)	O16—Na3—Na1	85.67 (5)
O13—Na1—O7	105.45 (7)	O14—Na3—Na1	96.42 (5)
O10—Na1—O7	78.58 (6)	O1—Na3—Na1	43.79 (4)
O4 ⁱ —Na1—O10 ⁱⁱ	81.40 (7)	Na2—Na3—Na1	62.14 (3)
O13—Na1—O10 ⁱⁱ	100.65 (7)	Na3 ^{vi} —Na3—Na1	109.09 (4)
O10—Na1—O10 ⁱⁱ	75.97 (7)	O3—Na4—O8 ^{vii}	88.53 (6)
O7—Na1—O10 ⁱⁱ	140.45 (7)	O3—Na4—O6	78.83 (6)
O4 ⁱ —Na1—O1	91.18 (7)	O8 ^{vii} —Na4—O6	167.28 (7)
O13—Na1—O1	78.18 (6)	O3—Na4—O17	95.65 (7)
O10—Na1—O1	142.97 (7)	O8 ^{vii} —Na4—O17	87.86 (6)
O7—Na1—O1	76.66 (6)	O6—Na4—O17	92.01 (6)
O10 ⁱⁱ —Na1—O1	138.45 (6)	O3—Na4—O11 ⁱⁱ	90.43 (7)
O4 ⁱ —Na1—O11 ⁱⁱ	88.36 (6)	O8 ^{vii} —Na4—O11 ⁱⁱ	92.18 (7)
O13—Na1—O11 ⁱⁱ	82.51 (6)	O6—Na4—O11 ⁱⁱ	89.29 (7)
O10—Na1—O11 ⁱⁱ	120.57 (6)	O17—Na4—O11 ⁱⁱ	173.92 (7)
O7—Na1—O11 ⁱⁱ	160.46 (6)	O3—Na4—O6 ^{viii}	163.22 (7)
O10 ⁱⁱ —Na1—O11 ⁱⁱ	51.33 (5)	O8 ^{vii} —Na4—O6 ^{viii}	108.07 (6)
O1—Na1—O11 ⁱⁱ	87.86 (5)	O6—Na4—O6 ^{viii}	84.51 (6)
O4 ⁱ —Na1—S4 ⁱⁱ	85.44 (5)	O17—Na4—O6 ^{viii}	82.85 (6)
O13—Na1—S4 ⁱⁱ	90.84 (5)	O11 ⁱⁱ —Na4—O6 ^{viii}	91.36 (7)
O10—Na1—S4 ⁱⁱ	97.33 (5)	O3—Na4—Na4 ^{viii}	122.20 (6)
O7—Na1—S4 ⁱⁱ	162.36 (5)	O8 ^{vii} —Na4—Na4 ^{viii}	149.15 (6)

O10 ⁱⁱ —Na1—S4 ⁱⁱ	25.03 (4)	O6—Na4—Na4 ^{viii}	43.39 (4)
O1—Na1—S4 ⁱⁱ	114.07 (5)	O17—Na4—Na4 ^{viii}	86.40 (5)
O11 ⁱⁱ —Na1—S4 ⁱⁱ	26.35 (3)	O11 ⁱⁱ —Na4—Na4 ^{viii}	90.47 (6)
O4 ⁱ —Na1—Na3	135.93 (6)	O6 ^{viii} —Na4—Na4 ^{viii}	41.12 (4)
O13—Na1—Na3	40.08 (5)	O3—Na4—Na2 ^{vii}	81.93 (5)
O10—Na1—Na3	95.38 (6)	O8 ^{vii} —Na4—Na2 ^{vii}	60.20 (5)
O7—Na1—Na3	71.14 (5)	O6—Na4—Na2 ^{vii}	115.90 (5)
O10 ⁱⁱ —Na1—Na3	140.70 (6)	O17—Na4—Na2 ^{vii}	30.76 (4)
O1—Na1—Na3	50.52 (4)	O11 ⁱⁱ —Na4—Na2 ^{vii}	151.34 (6)
O11 ⁱⁱ —Na1—Na3	108.04 (5)	O6 ^{viii} —Na4—Na2 ^{vii}	103.86 (5)
S4 ⁱⁱ —Na1—Na3	126.48 (3)	Na4 ^{viii} —Na4—Na2 ^{vii}	117.00 (4)
O4 ⁱ —Na1—Na2	78.15 (5)	S1—O1—Na2	131.01 (10)
O13—Na1—Na2	98.51 (5)	S1—O1—Na1	125.55 (9)
O10—Na1—Na2	114.45 (6)	Na2—O1—Na1	98.21 (6)
O7—Na1—Na2	37.60 (4)	S1—O1—Na3	114.45 (9)
O10 ⁱⁱ —Na1—Na2	159.32 (5)	Na2—O1—Na3	86.90 (5)
O1—Na1—Na2	39.92 (4)	Na1—O1—Na3	85.68 (5)
O11 ⁱⁱ —Na1—Na2	124.52 (4)	S1—O2—Na3 ^v	148.85 (11)
S4 ⁱⁱ —Na1—Na2	147.72 (3)	S1—O3—Na4	131.30 (9)
Na3—Na1—Na2	58.66 (3)	S2—O4—Na1 ⁱ	150.62 (11)
O4 ⁱ —Na1—Na1 ⁱⁱ	97.06 (6)	S2—O6—Na4	130.70 (10)
O13—Na1—Na1 ⁱⁱ	92.35 (5)	S2—O6—Na4 ^{viii}	123.77 (9)
O10—Na1—Na1 ⁱⁱ	39.63 (5)	Na4—O6—Na4 ^{viii}	95.49 (6)
O7—Na1—Na1 ⁱⁱ	112.71 (5)	S3—O7—Na2	116.86 (9)
O10 ⁱⁱ —Na1—Na1 ⁱⁱ	36.33 (4)	S3—O7—Na1	134.13 (9)
O1—Na1—Na1 ⁱⁱ	168.41 (6)	Na2—O7—Na1	104.11 (6)
O11 ⁱⁱ —Na1—Na1 ⁱⁱ	84.25 (4)	S3—O8—Na4 ⁱⁱⁱ	144.64 (11)
S4 ⁱⁱ —Na1—Na1 ⁱⁱ	58.81 (2)	S4—O10—Na1	135.88 (11)
Na3—Na1—Na1 ⁱⁱ	124.47 (4)	S4—O10—Na1 ⁱⁱ	106.52 (10)
Na2—Na1—Na1 ⁱⁱ	150.21 (4)	Na1—O10—Na1 ⁱⁱ	104.03 (7)
O14—Na2—O17 ⁱⁱⁱ	174.53 (8)	S4—O11—Na4 ⁱⁱ	136.83 (11)
O14—Na2—O15	94.48 (7)	S4—O11—Na1 ⁱⁱ	90.67 (9)
O17 ⁱⁱⁱ —Na2—O15	87.48 (7)	Na4 ⁱⁱ —O11—Na1 ⁱⁱ	117.01 (7)
O14—Na2—O7	88.36 (7)	Na1—O13—Na3	100.93 (7)
O17 ⁱⁱⁱ —Na2—O7	86.24 (6)	Na1—O13—H13A	111.6
O15—Na2—O7	101.40 (7)	Na3—O13—H13A	111.6
O14—Na2—O15 ^{iv}	99.06 (7)	Na1—O13—H13B	111.6
O17 ⁱⁱⁱ —Na2—O15 ^{iv}	86.17 (7)	Na3—O13—H13B	111.6
O15—Na2—O15 ^{iv}	85.64 (7)	H13A—O13—H13B	109.4
O7—Na2—O15 ^{iv}	169.40 (7)	Na2—O14—Na3	98.94 (7)
O14—Na2—O1	91.38 (7)	Na2—O14—H14A	112.0
O17 ⁱⁱⁱ —Na2—O1	86.73 (6)	Na3—O14—H14A	112.0

supplementary materials

O15—Na2—O1	174.12 (7)	Na2—O14—H14B	112.0
O7—Na2—O1	79.24 (6)	Na3—O14—H14B	112.0
O15 ^{iv} —Na2—O1	92.95 (6)	H14A—O14—H14B	109.7
O14—Na2—Na2 ^{iv}	99.26 (6)	Na2—O15—Na2 ^{iv}	94.36 (7)
O17 ⁱⁱⁱ —Na2—Na2 ^{iv}	85.67 (6)	Na2—O15—H15A	112.9
O15—Na2—Na2 ^{iv}	43.03 (5)	Na2 ^{iv} —O15—H15A	112.9
O7—Na2—Na2 ^{iv}	143.79 (6)	Na2—O15—H15B	112.9
O15 ^{iv} —Na2—Na2 ^{iv}	42.61 (5)	Na2 ^{iv} —O15—H15B	112.9
O1—Na2—Na2 ^{iv}	135.28 (6)	H15A—O15—H15B	110.3
O14—Na2—Na3	42.03 (5)	Na3 ^{vi} —O16—Na3	96.30 (7)
O17 ⁱⁱⁱ —Na2—Na3	134.64 (6)	Na3 ^{vi} —O16—H16A	112.5
O15—Na2—Na3	134.80 (6)	Na3—O16—H16A	112.5
O7—Na2—Na3	71.87 (5)	Na3 ^{vi} —O16—H16B	112.5
O15 ^{iv} —Na2—Na3	108.79 (6)	Na3—O16—H16B	112.5
O1—Na2—Na3	51.02 (5)	H16A—O16—H16B	110.0
Na2 ^{iv} —Na2—Na3	134.34 (4)	Na2 ^{vii} —O17—Na4	118.30 (8)
O14—Na2—Na1	96.33 (6)	Na2 ^{vii} —O17—H17A	107.7
O17 ⁱⁱⁱ —Na2—Na1	78.85 (5)	Na4—O17—H17A	107.7
O15—Na2—Na1	137.49 (5)	Na2 ^{vii} —O17—H17B	107.7
O7—Na2—Na1	38.29 (4)	Na4—O17—H17B	107.7
O15 ^{iv} —Na2—Na1	132.48 (5)	H17A—O17—H17B	107.1
O1—Na2—Na1	41.87 (4)	O2—S1—O3	113.99 (10)
Na2 ^{iv} —Na2—Na1	164.30 (4)	O2—S1—O1	113.19 (9)
Na3—Na2—Na1	59.20 (3)	O3—S1—O1	112.55 (10)
O14—Na2—Na4 ⁱⁱⁱ	146.42 (6)	O2—S1—C1	106.32 (10)
O17 ⁱⁱⁱ —Na2—Na4 ⁱⁱⁱ	30.94 (5)	O3—S1—C1	106.12 (9)
O15—Na2—Na4 ⁱⁱⁱ	66.79 (5)	O1—S1—C1	103.62 (10)
O7—Na2—Na4 ⁱⁱⁱ	69.72 (5)	O5—S2—O4	114.42 (11)
O15 ^{iv} —Na2—Na4 ⁱⁱⁱ	106.56 (5)	O5—S2—O6	112.59 (11)
O1—Na2—Na4 ⁱⁱⁱ	108.29 (5)	O4—S2—O6	111.20 (10)
Na2 ^{iv} —Na2—Na4 ⁱⁱⁱ	85.84 (3)	O5—S2—C1	107.49 (10)
Na3—Na2—Na4 ⁱⁱⁱ	139.39 (3)	O4—S2—C1	102.79 (10)
Na1—Na2—Na4 ⁱⁱⁱ	82.22 (3)	O6—S2—C1	107.57 (10)
O2 ^v —Na3—O13	123.40 (7)	O9—S3—O7	113.00 (10)
O2 ^v —Na3—O16 ^{vi}	86.03 (7)	O9—S3—O8	112.97 (10)
O13—Na3—O16 ^{vi}	85.41 (7)	O7—S3—O8	112.99 (10)
O2 ^v —Na3—O16	152.16 (8)	O9—S3—C2	106.55 (10)
O13—Na3—O16	81.48 (6)	O7—S3—C2	107.55 (10)
O16 ^{vi} —Na3—O16	83.70 (7)	O8—S3—C2	102.91 (10)
O2 ^v —Na3—O14	83.66 (7)	O10—S4—O12	113.60 (12)
O13—Na3—O14	134.23 (8)	O10—S4—O11	111.27 (11)
O16 ^{vi} —Na3—O14	136.85 (8)	O12—S4—O11	113.07 (11)

O16—Na3—O14	86.32 (7)	O10—S4—C2	106.45 (10)
O2 ^v —Na3—O1	79.21 (6)	O12—S4—C2	107.98 (11)
O13—Na3—O1	71.68 (6)	O11—S4—C2	103.70 (10)
O16 ^{vi} —Na3—O1	138.69 (7)	O10—S4—Na1 ⁱⁱ	48.45 (8)
O16—Na3—O1	124.33 (7)	O12—S4—Na1 ⁱⁱ	137.74 (8)
O14—Na3—O1	79.74 (6)	O11—S4—Na1 ⁱⁱ	62.98 (8)
O2 ^v —Na3—Na2	87.04 (5)	C2—S4—Na1 ⁱⁱ	113.81 (8)

Symmetry codes: (i) $-x+1, -y+2, -z+2$; (ii) $-x+1, -y+1, -z+2$; (iii) $x+1, y, z$; (iv) $-x+2, -y+2, -z+1$; (v) $-x+1, -y+2, -z+1$; (vi) $-x+1, -y+1, -z+1$; (vii) $x-1, y, z$; (viii) $-x, -y+2, -z+2$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O13—H13A \cdots O8 ^{vii}	0.97	1.93	2.868 (2)	163
O13—H13B \cdots O5 ^{ix}	0.97	1.87	2.824 (2)	166
O14—H14A \cdots O9 ^x	0.97	2.00	2.871 (2)	148
O15—H15B \cdots O3 ⁱⁱⁱ	0.97	1.97	2.877 (2)	155
O16—H16A \cdots O9 ^x	0.97	1.87	2.794 (2)	158
O16—H16B \cdots O12	0.97	2.18	3.034 (3)	147
O17—H17A \cdots O4 ^{viii}	0.97	2.17	2.879 (3)	129
O17—H17B \cdots O12 ^{xi}	0.97	2.10	2.918 (3)	141

Symmetry codes: (vii) $x-1, y, z$; (ix) $x, y-1, z$; (x) $-x+2, -y+1, -z+1$; (iii) $x+1, y, z$; (viii) $-x, -y+2, -z+2$; (xi) $x-1, y+1, z$.

Fig. 1




Fig. 2

