

5-Ethyl-2-methyl-3-phenylsulfonyl-1-benzofuran

Hong Dae Choi,^a Pil Ja Seo,^a Byeng Wha Son^b and Uk Lee^{b*}

^aDepartment of Chemistry, Dongeui University, San 24 Kaya-dong, Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea

Correspondence e-mail: uklee@pknu.ac.kr


Received 3 April 2008; accepted 1 May 2008

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.057; wR factor = 0.149; data-to-parameter ratio = 16.6.

The title compound, $\text{C}_{17}\text{H}_{16}\text{O}_3\text{S}$, was prepared by the oxidation of 5-ethyl-2-methyl-3-phenylsulfanyl-1-benzofuran with 3-chloroperoxybenzoic acid. The phenyl ring makes a dihedral angle of $75.94(8)^\circ$ with the plane of the benzofuran fragment. The crystal structure is stabilized by $\pi-\pi$ interactions between the furan rings of neighbouring molecules [centroid–centroid distance = $3.620(4)\text{ \AA}$]. In addition, the crystal structure exhibits C–H··· π interactions.

Related literature

For the crystal structures of similar 3-phenylsulfonyl-1-benzofuran compounds, see: Choi *et al.* (2008a,b).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{16}\text{O}_3\text{S}$	$V = 1463.98(12)\text{ \AA}^3$
$M_r = 300.36$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 8.7009(4)\text{ \AA}$	$\mu = 0.23\text{ mm}^{-1}$
$b = 8.2019(4)\text{ \AA}$	$T = 173(2)\text{ K}$
$c = 20.682(1)\text{ \AA}$	$0.40 \times 0.20 \times 0.20\text{ mm}$
$\beta = 97.301(1)^\circ$	

Data collection

Bruker SMART CCD diffractometer	3194 independent reflections
Absorption correction: none	2264 reflections with $I > 2\sigma(I)$
8693 measured reflections	$R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$	192 parameters
$wR(F^2) = 0.148$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.42\text{ e \AA}^{-3}$
3194 reflections	$\Delta\rho_{\text{min}} = -0.49\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C15—H15C···Cg2 ⁱ	0.98	2.80	3.592 (4)	139
C16—H16B···Cg2 ⁱⁱ	0.98	3.21	3.903 (4)	128

Symmetry codes: (i) $-x + 1, -y + 2, -z$; (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CF2194).

References

- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2001). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2008a). *Acta Cryst. E64*, o793.
Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2008b). *Acta Cryst. E64*, o794.
Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o1016 [doi:10.1107/S1600536808012877]

5-Ethyl-2-methyl-3-phenylsulfonyl-1-benzofuran

H. D. Choi, P. J. Seo, B. W. Son and U. Lee

Comment

As a part of our ongoing studies on the synthesis and structure of 3-phenyl-sulfonyl-1-benzofuran analogues, the crystal structure of 5-bromo-2-methyl-3-phenylsulfonyl-1-benzofuran (Choi *et al.*, 2008a) and 2,5,7-trimethyl-3-phenylsulfonyl-1-benzofuran (Choi *et al.*, 2008b) have been described in the literature. Here we report the crystal structure of the title compound, 5-ethyl-2-methyl-3-phenylsulfonyl-1-benzofuran (Fig. 1).

The benzofuran unit is essentially planar, with a mean deviation of 0.007 Å from the least-squares plane defined by the nine constituent atoms. The phenyl ring (C9—C14) makes a dihedral angle of 75.94 (8)° with the plane of the benzofuran fragment. The crystal packing (Fig. 2) is stabilized by aromatic π – π stacking interactions between the furan rings of neighbouring molecules. The $Cg1 \cdots Cg1^i$ distance is 3.620 (4) Å ($Cg1$ is the centroid of the O1/C8/C1/C2/C7 furan ring, symmetry code as in Fig. 2). The molecular packing is further stabilized by C—H \cdots π interactions (Table 1 and Fig. 2); one between a methyl H atom and the benzene ring of the benzofuran unit, *i.e.* C15—H15C \cdots $Cg2^i$, a second between a methylene H atom of the ethyl group and the benzene ring of the benzofuran unit, *i.e.* C16—H16B \cdots $Cg2^{ii}$, respectively. In both cases the benzene ring ($Cg2$) is involved ($Cg2$ is the centroid of the C2—7 benzene ring, symmetry code as in Fig. 2).

Experimental

3-Chloroperoxybenzoic acid (77%, 471 mg, 2.1 mmol) was added in small portions to a stirred solution of 5-ethyl-2-methyl-3-phenylsulfonyl-1-benzofuran (268 mg, 1.0 mmol) in dichloromethane (30 ml) at 273 K. After being stirred for 4 h at room temperature, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated in vacuum. The residue was purified by column chromatography (hexane-ethyl acetate, 2:1 *v/v*) to afford the title compound as a colorless solid [yield 80%, m.p. 396–397 K; R_f = 0.66 (hexane-ethyl acetate, 2:1 *v/v*)]. Single crystals suitable for X-ray diffraction were prepared by evaporation of a solution of the title compound in benzene at room temperature. Spectroscopic analysis: ^1H NMR (CDCl_3 , 400 MHz) δ 1.26 (t, J = 7.68 Hz, 3H), 2.74 (q, J = 7.72 Hz, 2H), 2.79 (s, 3H), 7.14 (d, J = 8.44 Hz and 1.48 Hz, 1H), 7.32 (d, J = 8.44 Hz, 1H), 7.48–7.61 (m, 3H), 7.69 (s, 1H), 8.01 (d, J = 6.96 Hz, 2H); EI—MS 300 [M^+].

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.95 Å for aromatic H atoms, 0.99 Å for methylene H atoms and 0.98 Å for methyl H atoms, respectively, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic and methylene H atoms, and $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing displacement ellipsoids drawn at the 50% probability level.




Fig. 2. π — π and C—H \cdots π interactions (dotted lines) in the title compound. Cg denotes the ring centroids. [Symmetry code: (i) $-x + 1, -y + 2, -z$; (ii) $-x + 3/2, y + 1/2, -z + 1/2$; (iii) $-x + 3/2, y - 1/2, -z + 1/2$.]

5-Ethyl-2-methyl-3-phenylsulfonyl-1-benzofuran

Crystal data

$C_{17}H_{16}O_3S$	$F_{000} = 632$
$M_r = 300.36$	$D_x = 1.363 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P_2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 8.7009 (4) \text{ \AA}$	Cell parameters from 2779 reflections
$b = 8.2019 (4) \text{ \AA}$	$\theta = 2.4\text{--}27.9^\circ$
$c = 20.682 (1) \text{ \AA}$	$\mu = 0.23 \text{ mm}^{-1}$
$\beta = 97.301 (1)^\circ$	$T = 173 (2) \text{ K}$
$V = 1463.98 (12) \text{ \AA}^3$	Block, colorless
$Z = 4$	$0.40 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Bruker SMART CCD diffractometer	3194 independent reflections
Radiation source: fine-focus sealed tube	2264 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.048$
Detector resolution: 10.0 pixels mm^{-1}	$\theta_{\text{max}} = 27.0^\circ$
$T = 173(2) \text{ K}$	$\theta_{\text{min}} = 2.4^\circ$
φ and ω scans	$h = -8 \rightarrow 11$
Absorption correction: none	$k = -8 \rightarrow 10$
8693 measured reflections	$l = -24 \rightarrow 26$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.056$	H-atom parameters constrained
$wR(F^2) = 0.148$	$w = 1/[\sigma^2(F_o^2) + (0.0725P)^2 + 0.989P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\max} < 0.001$
3194 reflections	$\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$
192 parameters	$\Delta\rho_{\min} = -0.49 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S	0.22072 (7)	0.75221 (8)	0.07999 (3)	0.01957 (19)
O1	0.6098 (2)	0.7989 (3)	0.00433 (10)	0.0340 (5)
O2	0.1125 (2)	0.7474 (2)	0.02135 (9)	0.0302 (5)
O3	0.1953 (2)	0.8653 (2)	0.13092 (9)	0.0256 (4)
C1	0.4056 (3)	0.7908 (3)	0.06010 (12)	0.0206 (6)
C2	0.5331 (3)	0.8631 (3)	0.10261 (13)	0.0219 (6)
C3	0.5561 (3)	0.9250 (3)	0.16565 (13)	0.0253 (6)
H3	0.4738	0.9264	0.1918	0.030*
C4	0.7023 (3)	0.9852 (4)	0.18992 (15)	0.0318 (7)
C5	0.8208 (3)	0.9848 (4)	0.15001 (17)	0.0362 (8)
H5	0.9195	1.0266	0.1671	0.043*
C6	0.8006 (3)	0.9267 (4)	0.08721 (16)	0.0361 (8)
H6	0.8817	0.9286	0.0605	0.043*
C7	0.6548 (3)	0.8648 (4)	0.06503 (14)	0.0286 (6)
C8	0.4583 (3)	0.7531 (3)	0.00219 (13)	0.0283 (6)
C9	0.2322 (3)	0.5547 (3)	0.11444 (12)	0.0184 (5)
C10	0.3091 (3)	0.5341 (3)	0.17728 (13)	0.0246 (6)
H10	0.3552	0.6245	0.2010	0.030*
C11	0.3172 (3)	0.3800 (4)	0.20437 (13)	0.0280 (6)
H11	0.3697	0.3637	0.2470	0.034*
C12	0.2487 (3)	0.2487 (3)	0.16934 (14)	0.0294 (6)
H12	0.2542	0.1429	0.1882	0.035*

supplementary materials

C13	0.1722 (3)	0.2713 (3)	0.10688 (14)	0.0280 (6)
H13	0.1255	0.1809	0.0833	0.034*
C14	0.1637 (3)	0.4242 (3)	0.07899 (13)	0.0236 (6)
H14	0.1119	0.4400	0.0362	0.028*
C15	0.3901 (4)	0.6744 (4)	-0.05890 (14)	0.0389 (8)
H15A	0.2823	0.6446	-0.0556	0.058*
H15B	0.4494	0.5762	-0.0664	0.058*
H15C	0.3933	0.7503	-0.0953	0.058*
C16	0.7384 (4)	1.0484 (5)	0.25879 (18)	0.0546 (10)
H16A	0.8332	0.9924	0.2791	0.066*
H16B	0.7639	1.1657	0.2561	0.066*
C17	0.6211 (5)	1.0323 (6)	0.30328 (18)	0.0597 (11)
H17A	0.5272	1.0912	0.2855	0.090*
H17B	0.6612	1.0782	0.3459	0.090*
H17C	0.5965	0.9168	0.3083	0.090*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S	0.0181 (3)	0.0187 (3)	0.0212 (3)	0.0015 (2)	-0.0004 (2)	0.0000 (3)
O1	0.0302 (11)	0.0407 (13)	0.0338 (12)	0.0061 (9)	0.0137 (9)	0.0073 (10)
O2	0.0274 (10)	0.0313 (11)	0.0295 (10)	0.0013 (8)	-0.0058 (8)	0.0035 (9)
O3	0.0231 (10)	0.0222 (10)	0.0323 (11)	0.0004 (8)	0.0063 (8)	-0.0056 (8)
C1	0.0222 (13)	0.0177 (13)	0.0225 (13)	0.0017 (10)	0.0049 (10)	0.0029 (11)
C2	0.0177 (13)	0.0195 (13)	0.0289 (14)	0.0024 (10)	0.0045 (11)	0.0070 (11)
C3	0.0202 (14)	0.0259 (15)	0.0299 (15)	-0.0016 (11)	0.0037 (11)	0.0014 (12)
C4	0.0241 (15)	0.0291 (16)	0.0404 (17)	-0.0061 (12)	-0.0025 (12)	0.0039 (13)
C5	0.0175 (14)	0.0372 (18)	0.052 (2)	-0.0044 (12)	-0.0009 (13)	0.0117 (16)
C6	0.0208 (15)	0.0384 (18)	0.051 (2)	0.0047 (12)	0.0116 (14)	0.0179 (16)
C7	0.0264 (15)	0.0277 (15)	0.0329 (15)	0.0067 (12)	0.0082 (12)	0.0084 (13)
C8	0.0306 (15)	0.0293 (15)	0.0256 (14)	0.0069 (12)	0.0058 (12)	0.0089 (13)
C9	0.0171 (12)	0.0192 (13)	0.0191 (13)	-0.0002 (10)	0.0037 (10)	-0.0006 (10)
C10	0.0263 (14)	0.0255 (15)	0.0212 (13)	-0.0015 (11)	-0.0001 (11)	-0.0037 (11)
C11	0.0324 (16)	0.0306 (16)	0.0198 (13)	0.0005 (12)	-0.0008 (11)	0.0012 (12)
C12	0.0373 (16)	0.0208 (14)	0.0317 (15)	0.0006 (12)	0.0104 (13)	0.0044 (13)
C13	0.0333 (15)	0.0224 (15)	0.0289 (15)	-0.0044 (11)	0.0062 (12)	-0.0055 (12)
C14	0.0258 (14)	0.0239 (14)	0.0209 (13)	-0.0016 (11)	0.0023 (11)	-0.0019 (11)
C15	0.052 (2)	0.0423 (19)	0.0232 (15)	0.0102 (16)	0.0090 (14)	-0.0009 (14)
C16	0.0393 (19)	0.066 (3)	0.058 (2)	-0.0246 (18)	0.0013 (17)	-0.017 (2)
C17	0.059 (2)	0.077 (3)	0.042 (2)	-0.018 (2)	-0.0003 (18)	-0.019 (2)

Geometric parameters (\AA , $^\circ$)

S—O2	1.4386 (19)	C9—C10	1.395 (3)
S—O3	1.4417 (19)	C10—C11	1.381 (4)
S—C1	1.740 (3)	C10—H10	0.950
S—C9	1.767 (3)	C11—C12	1.389 (4)
O1—C8	1.367 (3)	C11—H11	0.950
O1—C7	1.377 (4)	C12—C13	1.388 (4)

C1—C8	1.371 (4)	C12—H12	0.950
C1—C2	1.451 (4)	C13—C14	1.379 (4)
C2—C3	1.390 (4)	C13—H13	0.950
C2—C7	1.391 (4)	C14—H14	0.950
C3—C4	1.397 (4)	C15—H15A	0.980
C3—H3	0.950	C15—H15B	0.980
C4—C5	1.400 (4)	C15—H15C	0.980
C4—C16	1.511 (5)	C16—C17	1.464 (5)
C5—C6	1.374 (5)	C16—H16A	0.990
C5—H5	0.950	C16—H16B	0.990
C6—C7	1.389 (4)	C17—H17A	0.980
C6—H6	0.950	C17—H17B	0.980
C8—C15	1.474 (4)	C17—H17C	0.980
C9—C14	1.388 (4)		
O2—S—O3	119.24 (12)	C11—C10—C9	118.9 (2)
O2—S—C1	109.38 (12)	C11—C10—H10	120.6
O3—S—C1	106.76 (12)	C9—C10—H10	120.6
O2—S—C9	108.19 (12)	C10—C11—C12	120.1 (3)
O3—S—C9	107.54 (11)	C10—C11—H11	119.9
C1—S—C9	104.81 (12)	C12—C11—H11	119.9
C8—O1—C7	107.4 (2)	C11—C12—C13	120.3 (3)
C8—C1—C2	107.9 (2)	C11—C12—H12	119.8
C8—C1—S	126.0 (2)	C13—C12—H12	119.8
C2—C1—S	126.13 (19)	C14—C13—C12	120.3 (3)
C3—C2—C7	119.4 (2)	C14—C13—H13	119.9
C3—C2—C1	136.5 (2)	C12—C13—H13	119.9
C7—C2—C1	104.1 (2)	C13—C14—C9	118.9 (2)
C2—C3—C4	118.8 (3)	C13—C14—H14	120.5
C2—C3—H3	120.6	C9—C14—H14	120.5
C4—C3—H3	120.6	C8—C15—H15A	109.5
C3—C4—C5	119.5 (3)	C8—C15—H15B	109.5
C3—C4—C16	122.0 (3)	H15A—C15—H15B	109.5
C5—C4—C16	118.4 (3)	C8—C15—H15C	109.5
C6—C5—C4	122.9 (3)	H15A—C15—H15C	109.5
C6—C5—H5	118.6	H15B—C15—H15C	109.5
C4—C5—H5	118.6	C17—C16—C4	118.9 (3)
C5—C6—C7	116.2 (3)	C17—C16—H16A	107.6
C5—C6—H6	121.9	C4—C16—H16A	107.6
C7—C6—H6	121.9	C17—C16—H16B	107.6
O1—C7—C6	125.9 (3)	C4—C16—H16B	107.6
O1—C7—C2	110.9 (2)	H16A—C16—H16B	107.0
C6—C7—C2	123.2 (3)	C16—C17—H17A	109.5
O1—C8—C1	109.7 (2)	C16—C17—H17B	109.5
O1—C8—C15	115.4 (2)	H17A—C17—H17B	109.5
C1—C8—C15	134.8 (3)	C16—C17—H17C	109.5
C14—C9—C10	121.5 (2)	H17A—C17—H17C	109.5
C14—C9—S	119.7 (2)	H17B—C17—H17C	109.5
C10—C9—S	118.85 (19)		

supplementary materials


O2—S—C1—C8	−26.7 (3)	C1—C2—C7—C6	179.3 (3)
O3—S—C1—C8	−157.0 (2)	C7—O1—C8—C1	0.8 (3)
C9—S—C1—C8	89.1 (3)	C7—O1—C8—C15	−178.2 (2)
O2—S—C1—C2	155.7 (2)	C2—C1—C8—O1	−1.1 (3)
O3—S—C1—C2	25.4 (3)	S—C1—C8—O1	−179.00 (19)
C9—S—C1—C2	−88.5 (2)	C2—C1—C8—C15	177.7 (3)
C8—C1—C2—C3	−179.7 (3)	S—C1—C8—C15	−0.2 (5)
S—C1—C2—C3	−1.8 (5)	O2—S—C9—C14	10.3 (2)
C8—C1—C2—C7	0.9 (3)	O3—S—C9—C14	140.3 (2)
S—C1—C2—C7	178.8 (2)	C1—S—C9—C14	−106.4 (2)
C7—C2—C3—C4	−1.0 (4)	O2—S—C9—C10	−169.12 (19)
C1—C2—C3—C4	179.6 (3)	O3—S—C9—C10	−39.1 (2)
C2—C3—C4—C5	1.3 (4)	C1—S—C9—C10	74.3 (2)
C2—C3—C4—C16	−177.6 (3)	C14—C9—C10—C11	0.2 (4)
C3—C4—C5—C6	−0.3 (5)	S—C9—C10—C11	179.6 (2)
C16—C4—C5—C6	178.6 (3)	C9—C10—C11—C12	−0.4 (4)
C4—C5—C6—C7	−0.9 (5)	C10—C11—C12—C13	0.2 (4)
C8—O1—C7—C6	−179.9 (3)	C11—C12—C13—C14	0.1 (4)
C8—O1—C7—C2	−0.2 (3)	C12—C13—C14—C9	−0.3 (4)
C5—C6—C7—O1	−179.2 (3)	C10—C9—C14—C13	0.1 (4)
C5—C6—C7—C2	1.2 (4)	S—C9—C14—C13	−179.2 (2)
C3—C2—C7—O1	−179.9 (2)	C3—C4—C16—C17	5.8 (6)
C1—C2—C7—O1	−0.4 (3)	C5—C4—C16—C17	−173.1 (3)
C3—C2—C7—C6	−0.3 (4)		

Hydrogen-bond geometry (\AA , °)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C15—H15C \cdots Cg2 ⁱ	0.98	2.80	3.592 (4)	139
C16—H16B \cdots Cg2 ⁱⁱ	0.98	3.21	3.903 (4)	128

Symmetry codes: (i) $-x+1, -y+2, -z$; (ii) $-x+3/2, y+1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

