

2,4,6-Tris(1-oxo-2-pyridylsulfanyl-methyl)mesitylene methanol solvate

B. Ravindran Durai Nayagam,^{a*} Samuel Robinson Jebas,^b
C. Ravi Samuelraj^a and Dieter Schollmeyer^c

^aDepartment of Chemistry, Popes College, Sawyerpuram 628 251, Tamil Nadu, India, ^bDepartment of Physics, Karunya University, Karunya Nagar, Coimbatore 641 114, India, and ^cInstitut für Organische Chemie, Universität Mainz, Duesbergweg 10-14, 55099 Mainz, Germany

Correspondence e-mail: b_ravidurai@yahoo.com


Received 1 May 2008; accepted 3 May 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.051; wR factor = 0.151; data-to-parameter ratio = 15.1.

In the title compound, $\text{C}_{27}\text{H}_{27}\text{N}_3\text{O}_3\text{S}_3\cdot\text{CH}_4\text{O}$, the dihedral angles formed by the mesitylene ring with the three oxypyridyl rings are 89.6 (1), 75.5 (1) and 80.69 (1)°, indicating that all three are nearly perpendicular to the mesitylene ring. Intramolecular $\text{C}-\text{H}\cdots\text{S}$ hydrogen bonds generate $S(6)$ ring motifs. The crystal structure is stabilized by intramolecular $\text{C}-\text{H}\cdots\text{S}$ and intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For related literature on the biological activity of N -oxides see: Lobana *et al.*, (1989); Symons & West (1985); Katsuyuki *et al.* (1991); Bovin *et al.* (1992); Leonard *et al.* (1955). For related literature on N -oxides, see: Jebas *et al.* (2005); Ravindran *et al.* (2008). For bond-length data, see: Allen *et al.* (1987); Jebas *et al.* (2005); Ravindran *et al.* (2008). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{27}\text{H}_{27}\text{N}_3\text{O}_3\text{S}_3\cdot\text{CH}_4\text{O}$
 $M_r = 569.74$
Monoclinic, $P2_1/c$
 $a = 11.9644$ (17) Å
 $b = 14.9129$ (8) Å
 $c = 15.467$ (2) Å
 $\beta = 91.733$ (7)°

$V = 2758.4$ (6) Å³
 $Z = 4$
Cu $K\alpha$ radiation
 $\mu = 2.78$ mm⁻¹
 $T = 298$ (2) K
 $0.52 \times 0.42 \times 0.06$ mm

Data collection

Enraf-Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.296$, $T_{\max} = 0.842$
5226 measured reflections

5226 independent reflections
4156 reflections with $I > 2\sigma(I)$
3 standard reflections
frequency: 60 min
intensity decay: 3%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.150$
 $S = 1.08$
5226 reflections
347 parameters

12 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.48$ e Å⁻³
 $\Delta\rho_{\min} = -0.32$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1L—H1L \cdots O27	0.82	2.41	2.804 (7)	110
C16—H16A \cdots O36 ⁱ	0.96	2.47	3.348 (4)	152
C16—H16B \cdots S20	0.96	2.68	3.420 (3)	135
C18—H18B \cdots S29	0.96	2.79	3.515 (3)	133
C25—H25 \cdots O7 ⁱⁱ	0.93	2.44	3.147 (5)	133
C28—H28A \cdots O7 ⁱ	0.97	2.48	3.397 (3)	157
C31—H31 \cdots O27 ⁱⁱⁱ	0.93	2.35	3.107 (4)	139
C2—H2 \cdots Cg1 ^{iv}	0.93	2.91	3.774 (3)	154
C4—H4 \cdots Cg1 ^v	0.93	2.67	3.377 (3)	134

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x + 1, y, z$; (iv) $-x + 1, -y + 1, -z$; (v) $x, -y - \frac{1}{2}, z - \frac{3}{2}$. Cg1 is the centroid of the C10–C15 ring.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *CAD-4 Software*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

BRDN thanks the University Grants Commission, India, for a Teacher Fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2565).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
Bernstein, J., Davis, R. E., Shimon, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bovin, D. H. R., Crepon, E. & Zard, S. Z. (1992). *Bull. Soc. Chim. Fr.* **129**, 145–150.
Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
Jebas, S. R., Balasubramanian, T., Ravidurai, B. & Kumaresan, S. (2005). *Acta Cryst.* **E61**, o2677–o2678.

- Katsuyuki, N., Carter, B. J., Xu, J. & Hetch, S. M. (1991). *J. Am. Chem. Soc.* **113**, 5099–5100.
- Leonard, F., Barklay, F. A., Brown, E. V., Anderson, F. E. & Green, D. M. (1955). *Antibiot. Chemother.* pp. 261–264.
- Lobana, T. S. & Bhatia, P. K. (1989). *J. Sci. Ind. Res.* **48**, 394–401.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Ravindran Durai Nayagam, B., Jebas, S. R., Grace, S. & Schollmeyer, D. (2008). *Acta Cryst.* **E64**, o409.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Symons, M. C. R. & West, D.-X. (1985). *J. Chem. Soc. Dalton Trans.* pp. 379–381.

supplementary materials

Acta Cryst. (2008). E64, o1036-o1037 [doi:10.1107/S1600536808013081]

2,4,6-Tris(1-oxo-2-pyridylsulfanylmethyl)mesitylene methanol solvate

B. Ravindran Durai Nayagam, S. R. Jebas, C. Ravi Samuelraj and D. Schollmeyer

Comment

N-Oxides and their derivatives show a broad spectrum of biological activity, such as antifungal, antibacterial, antimicrobial and antibacterial activities (Lobana & Bhatia, 1989; Symons *et al.*, 1985). These compounds are also found to be involved in DNA strand scission under physiological conditions (Katsuyuki *et al.*, 1991; Bovin *et al.* 1992). Pyridine N-oxides bearing a sulfur group in position 2 display significant antimicrobial activity (Leonard *et al.*, 1955). In view of the importance of N-oxides, we have previously reported the crystal structures of N-oxide derivatives (Jebas *et al.*, 2005; Ravindran *et al.*, 2008). As an extension of our work on these derivatives, we report here the crystal structure of the title compound (Fig. 1).

The bond lengths and angles agree well with the N-oxide derivatives reported earlier (Jebas *et al.*, 2005; Ravindran *et al.*, 2008). The N–O bond length is in good agreement with the mean value of 1.304 (15) Å reported in the literature for pyridine N–oxides (Allen *et al.*, 1987).

The meistyrene ring is planar with the maximum deviation from planarity being -0.036 (1) Å. The dihedral angle formed by the meistyrene ring with the oxopyridinium rings (C1–C5/N6) 89.6 (1) °; (C21–C25/N26) 75.5 (1) ° and (C30–C34/N35) 80.69 (1) ° respectively, indicating that all the three oxopyridinium rings are perpendicular to the meistyrene ring.

Intramolecular C—H···S hydrogen bonds generate S(6)S(6) ring motifs. The crystal structure is stabilized by intramolecular C—H···S and intermolecular C—H···O hydrogen bonds and weak C—H··· π interactions (Table 1, where Cg1 is the centroid of the ring C10-C15).

Experimental

A mixture of tris(bromomethyl)mesitylene (0.399 g, 1 mmol) and 1-hydroxypyridine-2-thione sodium salt (0.448 g, 3 mmol) in water (30 ml) and methanol (30 ml) was heated at 333 K with stirring for 30 min. The compound formed was filtered off, and dried (0.494 g, 92%). The compound was recrystallized from chloroform-methanol (1:2 v/v).

Refinement

H atoms were positioned geometrically [C—H = 0.93 (aromatic), 0.96 Å (methyl) 0.97 Å (methylene), and 0.82 Å O—H] and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2$ or $-1.5U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atomic numbering scheme.

2,4,6-Tris(1-oxo-2-pyridylsulfanylmethyl)mesitylene methanol solvate

Crystal data

$C_{27}H_{27}N_3O_3S_3 \cdot CH_4O$

$M_r = 569.74$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 11.9644$ (17) Å

$b = 14.9129$ (8) Å

$c = 15.467$ (2) Å

$\beta = 91.733$ (7)°

$V = 2758.4$ (6) Å³

$Z = 4$

$F_{000} = 1200$

$D_x = 1.372$ Mg m⁻³

Cu $K\alpha$ radiation

$\lambda = 1.54178$ Å

Cell parameters from 25 reflections

$\theta = 61\text{--}69^\circ$

$\mu = 2.78$ mm⁻¹

$T = 298$ (2) K

Block, colourless

$0.52 \times 0.42 \times 0.06$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer

Monochromator: graphite

$\omega/2\theta$ scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.296$, $T_{\max} = 0.842$

5226 measured reflections

5226 independent reflections

4156 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0$

$\theta_{\max} = 69.9^\circ$

$\theta_{\min} = 3.7^\circ$

$h = 0 \rightarrow 14$

$k = -18 \rightarrow 0$

$l = -18 \rightarrow 18$

3 standard reflections

every 60 min

intensity decay: 3%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.051$

$wR(F^2) = 0.150$

12 restraints

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0834P)^2 + 0.7123P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$S = 1.09$

$$\Delta\rho_{\max} = 0.49 \text{ e } \text{\AA}^{-3}$$

5226 reflections

$$\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$$

347 parameters

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3585 (2)	0.25409 (17)	-0.03619 (16)	0.0411 (6)
C2	0.4031 (2)	0.2831 (2)	-0.11233 (17)	0.0509 (7)
H2	0.4415	0.3372	-0.114	0.061*
C3	0.3906 (3)	0.2318 (2)	-0.18594 (19)	0.0595 (8)
H3	0.4189	0.2516	-0.2379	0.071*
C4	0.3357 (3)	0.1508 (2)	-0.1818 (2)	0.0627 (8)
H4	0.3283	0.1149	-0.2307	0.075*
C5	0.2921 (3)	0.1234 (2)	-0.1054 (2)	0.0593 (8)
H5	0.2552	0.0686	-0.1028	0.071*
N6	0.30179 (19)	0.17480 (15)	-0.03383 (15)	0.0475 (5)
O7	0.2579 (2)	0.15045 (15)	0.03880 (14)	0.0660 (6)
S8	0.36161 (6)	0.30782 (4)	0.06449 (4)	0.04599 (19)
C9	0.4460 (2)	0.40587 (17)	0.04115 (15)	0.0442 (6)
H9A	0.5203	0.3879	0.0245	0.053*
H9B	0.4117	0.4404	-0.0057	0.053*
C10	0.4525 (2)	0.46111 (16)	0.12348 (15)	0.0380 (5)
C11	0.3677 (2)	0.52357 (17)	0.13909 (16)	0.0407 (5)
C12	0.3725 (2)	0.57310 (16)	0.21599 (16)	0.0401 (5)
C13	0.4580 (2)	0.55854 (16)	0.27863 (16)	0.0401 (5)
C14	0.5453 (2)	0.49992 (16)	0.25966 (15)	0.0384 (5)
C15	0.5413 (2)	0.44950 (16)	0.18286 (15)	0.0384 (5)
C16	0.2719 (3)	0.5372 (2)	0.07464 (19)	0.0551 (7)
H16A	0.2881	0.5871	0.0378	0.083*
H16B	0.2047	0.5491	0.1049	0.083*
H16C	0.2621	0.4841	0.0401	0.083*
C17	0.4548 (3)	0.6036 (2)	0.36637 (19)	0.0588 (8)
H17A	0.4966	0.6585	0.3652	0.088*
H17B	0.487	0.5645	0.4096	0.088*
H17C	0.3786	0.6164	0.3799	0.088*
C18	0.6333 (3)	0.3827 (2)	0.16668 (19)	0.0535 (7)
H18A	0.6067	0.3384	0.126	0.08*
H18B	0.6552	0.354	0.2201	0.08*
H18C	0.6964	0.4133	0.1437	0.08*
C19	0.2853 (2)	0.64413 (18)	0.2303 (2)	0.0489 (6)

supplementary materials

H19A	0.2688	0.6758	0.1766	0.059*
H19B	0.313	0.6872	0.2728	0.059*
S20	0.15895 (7)	0.59076 (5)	0.26844 (6)	0.0598 (2)
C21	0.0714 (2)	0.6832 (2)	0.2755 (2)	0.0534 (7)
C22	0.0959 (3)	0.7726 (2)	0.2634 (2)	0.0627 (8)
H22	0.167	0.7891	0.2465	0.075*
C23	0.0164 (3)	0.8378 (3)	0.2760 (3)	0.0754 (10)
H23	0.0333	0.898	0.2674	0.09*
C24	-0.0876 (3)	0.8128 (3)	0.3012 (3)	0.0837 (12)
H24	-0.1412	0.8561	0.3121	0.1*
C25	-0.1127 (3)	0.7240 (3)	0.3104 (3)	0.0831 (12)
H25	-0.1841	0.7072	0.3263	0.1*
N26	-0.0351 (2)	0.6605 (2)	0.29666 (19)	0.0687 (8)
O27	-0.0591 (2)	0.57455 (19)	0.3027 (2)	0.0984 (10)
C28	0.6442 (2)	0.49055 (18)	0.32203 (17)	0.0450 (6)
H28A	0.6531	0.5448	0.3561	0.054*
H28B	0.712	0.4813	0.2903	0.054*
S29	0.62016 (6)	0.39480 (5)	0.39312 (5)	0.0517 (2)
C30	0.7555 (2)	0.37491 (19)	0.43343 (17)	0.0475 (6)
C31	0.8492 (3)	0.4281 (2)	0.4247 (2)	0.0604 (8)
H31	0.8431	0.4844	0.3993	0.073*
C32	0.9529 (3)	0.3962 (3)	0.4545 (3)	0.0771 (10)
H32	1.0171	0.4301	0.4475	0.093*
C33	0.9589 (4)	0.3142 (3)	0.4943 (3)	0.0871 (13)
H33	1.0278	0.2922	0.5142	0.104*
C34	0.8648 (4)	0.2647 (3)	0.5050 (2)	0.0772 (11)
H34	0.8701	0.2094	0.5327	0.093*
N35	0.7633 (2)	0.29458 (17)	0.47588 (16)	0.0572 (6)
O36	0.6727 (2)	0.24782 (16)	0.48565 (16)	0.0756 (7)
O1L	-0.1572 (5)	0.5507 (4)	0.1374 (4)	0.205 (2)
H1L	-0.12	0.5136	0.1649	0.308*
C2L	-0.0867 (7)	0.5996 (6)	0.0829 (5)	0.189 (3)
H2LA	-0.0521	0.6478	0.115	0.284*
H2LB	-0.1301	0.6237	0.0351	0.284*
H2LC	-0.03	0.5606	0.0615	0.284*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0480 (14)	0.0322 (12)	0.0432 (13)	0.0010 (10)	0.0013 (11)	-0.0040 (10)
C2	0.0599 (16)	0.0485 (16)	0.0447 (14)	-0.0036 (13)	0.0087 (12)	-0.0060 (12)
C3	0.0659 (19)	0.066 (2)	0.0471 (15)	0.0032 (16)	0.0085 (14)	-0.0108 (14)
C4	0.0639 (18)	0.065 (2)	0.0594 (18)	0.0085 (16)	-0.0044 (15)	-0.0287 (16)
C5	0.0639 (18)	0.0428 (16)	0.071 (2)	-0.0009 (14)	-0.0044 (15)	-0.0194 (14)
N6	0.0517 (12)	0.0371 (12)	0.0536 (13)	-0.0005 (10)	0.0007 (10)	-0.0030 (10)
O7	0.0869 (16)	0.0497 (12)	0.0620 (13)	-0.0199 (11)	0.0136 (12)	0.0040 (10)
S8	0.0649 (4)	0.0383 (3)	0.0351 (3)	-0.0115 (3)	0.0081 (3)	-0.0010 (2)
C9	0.0598 (16)	0.0381 (14)	0.0352 (12)	-0.0108 (12)	0.0078 (11)	-0.0017 (10)

C10	0.0494 (14)	0.0304 (12)	0.0346 (12)	-0.0079 (10)	0.0060 (10)	0.0004 (9)
C11	0.0465 (13)	0.0332 (12)	0.0425 (13)	-0.0049 (10)	0.0034 (11)	0.0048 (10)
C12	0.0450 (13)	0.0290 (12)	0.0467 (13)	-0.0001 (10)	0.0082 (11)	0.0028 (10)
C13	0.0497 (14)	0.0294 (12)	0.0416 (13)	0.0001 (10)	0.0079 (11)	-0.0017 (10)
C14	0.0464 (13)	0.0305 (12)	0.0384 (12)	-0.0026 (10)	0.0037 (10)	0.0024 (10)
C15	0.0477 (13)	0.0293 (12)	0.0387 (12)	-0.0008 (10)	0.0094 (10)	0.0007 (9)
C16	0.0574 (17)	0.0541 (17)	0.0534 (16)	0.0011 (14)	-0.0053 (13)	0.0049 (13)
C17	0.074 (2)	0.0518 (17)	0.0513 (16)	0.0057 (15)	0.0062 (14)	-0.0164 (13)
C18	0.0585 (17)	0.0464 (16)	0.0557 (16)	0.0127 (13)	0.0063 (13)	-0.0059 (13)
C19	0.0485 (14)	0.0331 (13)	0.0655 (17)	0.0037 (11)	0.0095 (13)	0.0016 (12)
S20	0.0567 (4)	0.0417 (4)	0.0823 (5)	0.0054 (3)	0.0209 (4)	0.0093 (3)
C21	0.0492 (15)	0.0523 (17)	0.0592 (17)	0.0065 (13)	0.0094 (13)	0.0061 (13)
C22	0.0565 (17)	0.0502 (17)	0.082 (2)	0.0055 (14)	0.0066 (16)	0.0005 (16)
C23	0.073 (2)	0.054 (2)	0.100 (3)	0.0166 (18)	0.006 (2)	0.0046 (19)
C24	0.074 (2)	0.078 (3)	0.099 (3)	0.032 (2)	0.014 (2)	0.010 (2)
C25	0.0563 (19)	0.096 (3)	0.099 (3)	0.020 (2)	0.0250 (19)	0.020 (2)
N26	0.0583 (15)	0.0674 (18)	0.0815 (19)	0.0050 (14)	0.0231 (14)	0.0188 (15)
O27	0.0813 (18)	0.0709 (17)	0.146 (3)	-0.0036 (14)	0.0488 (18)	0.0273 (17)
C28	0.0495 (14)	0.0383 (14)	0.0471 (14)	-0.0038 (11)	-0.0007 (12)	0.0030 (11)
S29	0.0522 (4)	0.0503 (4)	0.0524 (4)	-0.0049 (3)	-0.0013 (3)	0.0111 (3)
C30	0.0598 (16)	0.0425 (14)	0.0399 (13)	0.0037 (12)	-0.0041 (12)	-0.0029 (11)
C31	0.0605 (18)	0.0562 (18)	0.0639 (18)	-0.0008 (15)	-0.0104 (15)	-0.0032 (15)
C32	0.058 (2)	0.085 (3)	0.087 (3)	0.0017 (18)	-0.0141 (18)	-0.015 (2)
C33	0.080 (3)	0.092 (3)	0.087 (3)	0.030 (2)	-0.032 (2)	-0.016 (2)
C34	0.101 (3)	0.066 (2)	0.064 (2)	0.025 (2)	-0.022 (2)	0.0010 (17)
N35	0.0801 (18)	0.0444 (14)	0.0467 (13)	0.0080 (13)	-0.0035 (12)	0.0002 (11)
O36	0.0988 (18)	0.0528 (14)	0.0755 (15)	-0.0065 (13)	0.0083 (14)	0.0178 (12)
O1L	0.213 (4)	0.187 (4)	0.214 (4)	-0.065 (4)	-0.012 (3)	-0.012 (3)
C2L	0.178 (5)	0.210 (5)	0.179 (5)	-0.041 (4)	-0.017 (4)	0.006 (4)

Geometric parameters (Å, °)

C1—N6	1.364 (3)	C19—S20	1.822 (3)
C1—C2	1.377 (4)	C19—H19A	0.97
C1—S8	1.751 (2)	C19—H19B	0.97
C2—C3	1.376 (4)	S20—C21	1.737 (3)
C2—H2	0.93	C21—N26	1.368 (4)
C3—C4	1.377 (5)	C21—C22	1.378 (4)
C3—H3	0.93	C22—C23	1.379 (4)
C4—C5	1.368 (5)	C22—H22	0.93
C4—H4	0.93	C23—C24	1.368 (5)
C5—N6	1.349 (4)	C23—H23	0.93
C5—H5	0.93	C24—C25	1.366 (6)
N6—O7	1.306 (3)	C24—H24	0.93
S8—C9	1.820 (3)	C25—N26	1.348 (5)
C9—C10	1.517 (3)	C25—H25	0.93
C9—H9A	0.97	N26—O27	1.318 (4)
C9—H9B	0.97	C28—S29	1.830 (3)
C10—C15	1.394 (4)	C28—H28A	0.97

supplementary materials

C10—C11	1.404 (4)	C28—H28B	0.97
C11—C12	1.400 (4)	S29—C30	1.742 (3)
C11—C16	1.510 (4)	C30—N35	1.368 (4)
C12—C13	1.404 (4)	C30—C31	1.384 (4)
C12—C19	1.508 (3)	C31—C32	1.394 (5)
C13—C14	1.400 (3)	C31—H31	0.93
C13—C17	1.516 (4)	C32—C33	1.370 (6)
C14—C15	1.406 (3)	C32—H32	0.93
C14—C28	1.510 (3)	C33—C34	1.361 (6)
C15—C18	1.511 (4)	C33—H33	0.93
C16—H16A	0.96	C34—N35	1.358 (4)
C16—H16B	0.96	C34—H34	0.93
C16—H16C	0.96	N35—O36	1.302 (4)
C17—H17A	0.96	O1L—C2L	1.414 (9)
C17—H17B	0.96	O1L—H1L	0.82
C17—H17C	0.96	C2L—H2LA	0.96
C18—H18A	0.96	C2L—H2LB	0.96
C18—H18B	0.96	C2L—H2LC	0.96
C18—H18C	0.96		
N6—C1—C2	120.1 (2)	H18A—C18—H18C	109.5
N6—C1—S8	111.72 (18)	H18B—C18—H18C	109.5
C2—C1—S8	128.2 (2)	C12—C19—S20	108.98 (18)
C3—C2—C1	119.8 (3)	C12—C19—H19A	109.9
C3—C2—H2	120.1	S20—C19—H19A	109.9
C1—C2—H2	120.1	C12—C19—H19B	109.9
C2—C3—C4	119.3 (3)	S20—C19—H19B	109.9
C2—C3—H3	120.4	H19A—C19—H19B	108.3
C4—C3—H3	120.4	C21—S20—C19	100.44 (13)
C5—C4—C3	119.8 (3)	N26—C21—C22	118.4 (3)
C5—C4—H4	120.1	N26—C21—S20	112.7 (2)
C3—C4—H4	120.1	C22—C21—S20	128.9 (2)
N6—C5—C4	120.9 (3)	C21—C22—C23	120.8 (3)
N6—C5—H5	119.6	C21—C22—H22	119.6
C4—C5—H5	119.6	C23—C22—H22	119.6
O7—N6—C5	121.4 (2)	C24—C23—C22	119.0 (4)
O7—N6—C1	118.5 (2)	C24—C23—H23	120.5
C5—N6—C1	120.1 (3)	C22—C23—H23	120.5
C1—S8—C9	100.86 (12)	C25—C24—C23	119.9 (3)
C10—C9—S8	106.53 (16)	C25—C24—H24	120
C10—C9—H9A	110.4	C23—C24—H24	120
S8—C9—H9A	110.4	N26—C25—C24	120.7 (3)
C10—C9—H9B	110.4	N26—C25—H25	119.7
S8—C9—H9B	110.4	C24—C25—H25	119.7
H9A—C9—H9B	108.6	O27—N26—C25	121.3 (3)
C15—C10—C11	120.6 (2)	O27—N26—C21	117.7 (3)
C15—C10—C9	120.3 (2)	C25—N26—C21	121.0 (3)
C11—C10—C9	119.1 (2)	C14—C28—S29	108.76 (17)
C12—C11—C10	119.0 (2)	C14—C28—H28A	109.9
C12—C11—C16	120.1 (2)	S29—C28—H28A	109.9

C10—C11—C16	120.9 (2)	C14—C28—H28B	109.9
C11—C12—C13	121.0 (2)	S29—C28—H28B	109.9
C11—C12—C19	119.0 (2)	H28A—C28—H28B	108.3
C13—C12—C19	120.0 (2)	C30—S29—C28	100.76 (13)
C14—C13—C12	119.0 (2)	N35—C30—C31	120.3 (3)
C14—C13—C17	120.3 (2)	N35—C30—S29	111.7 (2)
C12—C13—C17	120.7 (2)	C31—C30—S29	127.9 (2)
C13—C14—C15	120.3 (2)	C30—C31—C32	119.2 (3)
C13—C14—C28	119.9 (2)	C30—C31—H31	120.4
C15—C14—C28	119.8 (2)	C32—C31—H31	120.4
C10—C15—C14	119.8 (2)	C33—C32—C31	119.2 (4)
C10—C15—C18	121.1 (2)	C33—C32—H32	120.4
C14—C15—C18	119.1 (2)	C31—C32—H32	120.4
C11—C16—H16A	109.5	C34—C33—C32	120.4 (4)
C11—C16—H16B	109.5	C34—C33—H33	119.8
H16A—C16—H16B	109.5	C32—C33—H33	119.8
C11—C16—H16C	109.5	N35—C34—C33	121.1 (4)
H16A—C16—H16C	109.5	N35—C34—H34	119.4
H16B—C16—H16C	109.5	C33—C34—H34	119.4
C13—C17—H17A	109.5	O36—N35—C34	121.7 (3)
C13—C17—H17B	109.5	O36—N35—C30	118.7 (3)
H17A—C17—H17B	109.5	C34—N35—C30	119.6 (3)
C13—C17—H17C	109.5	C2L—O1L—H1L	109.5
H17A—C17—H17C	109.5	O1L—C2L—H2LA	109.5
H17B—C17—H17C	109.5	O1L—C2L—H2LB	109.5
C15—C18—H18A	109.5	H2LA—C2L—H2LB	109.5
C15—C18—H18B	109.5	O1L—C2L—H2LC	109.5
H18A—C18—H18B	109.5	H2LA—C2L—H2LC	109.5
C15—C18—H18C	109.5	H2LB—C2L—H2LC	109.5
N6—C1—C2—C3	-0.2 (4)	C13—C14—C15—C10	3.0 (3)
S8—C1—C2—C3	-178.9 (2)	C28—C14—C15—C10	-177.1 (2)
C1—C2—C3—C4	-1.6 (5)	C13—C14—C15—C18	-176.4 (2)
C2—C3—C4—C5	1.6 (5)	C28—C14—C15—C18	3.5 (3)
C3—C4—C5—N6	0.2 (5)	C11—C12—C19—S20	81.2 (3)
C4—C5—N6—O7	178.1 (3)	C13—C12—C19—S20	-99.9 (2)
C4—C5—N6—C1	-1.9 (4)	C12—C19—S20—C21	-177.6 (2)
C2—C1—N6—O7	-178.1 (3)	C19—S20—C21—N26	174.9 (2)
S8—C1—N6—O7	0.8 (3)	C19—S20—C21—C22	-5.5 (4)
C2—C1—N6—C5	1.9 (4)	N26—C21—C22—C23	2.8 (5)
S8—C1—N6—C5	-179.1 (2)	S20—C21—C22—C23	-176.7 (3)
N6—C1—S8—C9	177.65 (19)	C21—C22—C23—C24	0.3 (6)
C2—C1—S8—C9	-3.5 (3)	C22—C23—C24—C25	-2.5 (7)
C1—S8—C9—C10	178.14 (19)	C23—C24—C25—N26	1.5 (7)
S8—C9—C10—C15	92.9 (2)	C24—C25—N26—O27	-177.9 (4)
S8—C9—C10—C11	-86.7 (2)	C24—C25—N26—C21	1.8 (6)
C15—C10—C11—C12	-1.1 (3)	C22—C21—N26—O27	175.8 (3)
C9—C10—C11—C12	178.6 (2)	S20—C21—N26—O27	-4.5 (4)
C15—C10—C11—C16	179.3 (2)	C22—C21—N26—C25	-3.9 (5)
C9—C10—C11—C16	-1.0 (3)	S20—C21—N26—C25	175.7 (3)

supplementary materials

C10—C11—C12—C13	-2.6 (4)	C13—C14—C28—S29	94.8 (2)
C16—C11—C12—C13	177.0 (2)	C15—C14—C28—S29	-85.0 (2)
C10—C11—C12—C19	176.3 (2)	C14—C28—S29—C30	162.19 (18)
C16—C11—C12—C19	-4.1 (4)	C28—S29—C30—N35	-168.0 (2)
C11—C12—C13—C14	6.3 (4)	C28—S29—C30—C31	10.1 (3)
C19—C12—C13—C14	-172.5 (2)	N35—C30—C31—C32	4.3 (5)
C11—C12—C13—C17	-172.2 (2)	S29—C30—C31—C32	-173.7 (3)
C19—C12—C13—C17	9.0 (4)	C30—C31—C32—C33	-2.1 (5)
C12—C13—C14—C15	-6.5 (3)	C31—C32—C33—C34	-0.3 (6)
C17—C13—C14—C15	172.0 (2)	C32—C33—C34—N35	0.7 (6)
C12—C13—C14—C28	173.6 (2)	C33—C34—N35—O36	180.0 (3)
C17—C13—C14—C28	-7.9 (4)	C33—C34—N35—C30	1.4 (5)
C11—C10—C15—C14	0.8 (3)	C31—C30—N35—O36	177.5 (3)
C9—C10—C15—C14	-178.9 (2)	S29—C30—N35—O36	-4.2 (3)
C11—C10—C15—C18	-179.8 (2)	C31—C30—N35—C34	-3.9 (4)
C9—C10—C15—C18	0.5 (3)	S29—C30—N35—C34	174.3 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1L—H1L \cdots O27	0.82	2.41	2.804 (7)	110
C16—H16A \cdots O36 ⁱ	0.96	2.47	3.348 (4)	152
C16—H16B \cdots S20	0.96	2.68	3.420 (3)	135
C18—H18B \cdots S29	0.96	2.79	3.515 (3)	133
C25—H25 \cdots O7 ⁱⁱ	0.93	2.44	3.147 (5)	133
C28—H28A \cdots O7 ⁱ	0.97	2.48	3.397 (3)	157
C31—H31 \cdots O27 ⁱⁱⁱ	0.93	2.35	3.107 (4)	139
C2—H2 \cdots Cg1 ^{iv}	0.93	2.91	3.774 (3)	154
C4—H4 \cdots Cg1 ^v	0.93	2.67	3.377 (3)	134

Symmetry codes: (i) $-x+1, y+1/2, -z+1/2$; (ii) $-x, y+1/2, -z+1/2$; (iii) $x+1, y, z$; (iv) $-x+1, -y+1, -z$; (v) $x, -y-1/2, z-3/2$.

Fig. 1

