

Tetrakis(μ -4-ethylbenzoato- κ^2 O:O')-bis[(4-ethylbenzoic acid- κ O)copper(II)]

Abraham C. Sunil, Barend C. B. Bezuidenhoudt* and J. Marthinus Janse van Rensburg

Department of Chemistry, University of the Free State, PO Box 339, Bloemfontein 9300, South Africa

Correspondence e-mail: bezuidbc.sci@ufs.ac.za


Received 20 May 2008; accepted 27 May 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.041; wR factor = 0.104; data-to-parameter ratio = 18.3.

The molecule of the title compound, $[Cu_2(C_9H_9O_2)_4 \cdot (C_9H_{10}O_2)_2]$, lies on a center of inversion. It consists of four bridging ethylbenzoate ligands, forming a cage around two Cu atoms in a *syn-syn* configuration, and two monodentate ethylbenzoic acid ligands bonded apically to the square-planar Cu atoms. The Cu···Cu distance is 2.6047 (5) Å.

Related literature

For the synthesis of aromatic carboxylic acids, see: Kaeding (1967). For tetrakis(μ_2 -2-methylbenzoato)bis(2-methylbenzoic acid)dicopper(II), see: Sunil *et al.* (2008). For tetrakis(μ_2 -2-fluorobenzoato)bis(2-fluorobenzoic acid)dicopper(II), see: Valach *et al.* (2000). For tetrakis(μ_2 benzoato) bis(2-fluorobenzoic acid)dicopper(II), see: Kawata *et al.* (1992). For tetrakis-[μ -(2-phenoxybenzoato- O,O')]bis[(2-phenoxybenzoic acid)copper(II)], see: Mak & Yip (1990).



Experimental

Crystal data

$[Cu_2(C_9H_9O_2)_4(C_9H_{10}O_2)_2]$	$\gamma = 79.468 (2)^\circ$
$M_r = 1024.07$	$V = 1199.47 (12)$ Å ³
Triclinic, $P\bar{1}$	$Z = 1$
$a = 10.6167 (5)$ Å	Mo $K\alpha$ radiation
$b = 10.7394 (7)$ Å	$\mu = 0.95$ mm ⁻¹
$c = 10.8096 (7)$ Å	$T = 100 (2)$ K
$\alpha = 81.848 (3)^\circ$	$0.54 \times 0.4 \times 0.39$ mm
$\beta = 88.594 (3)^\circ$	

Data collection

Bruker Kappa APEXII	15971 measured reflections
diffractometer	5683 independent reflections
Absorption correction: multi-scan	4721 reflections with $I > 2\sigma(I)$
(SADABS; Bruker, 2004)	$R_{\text{int}} = 0.040$
$T_{\min} = 0.628$, $T_{\max} = 0.708$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	311 parameters
$wR(F^2) = 0.103$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\max} = 0.50$ e Å ⁻³
5683 reflections	$\Delta\rho_{\min} = -0.37$ e Å ⁻³

Table 1
Selected bond lengths (Å).

Cu1—O3	1.9498 (15)	Cu1—O1	2.0040 (16)
Cu1—O4	1.9501 (16)	Cu1—O5	2.1761 (15)
Cu1—O2	1.9593 (16)	Cu1—Cu1 ⁱ	2.6047 (5)

Symmetry code: (i) $-x + 1, -y + 1, -z + 2$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT-Plus* (Bruker, 2004); data reduction: *SAINT-Plus* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2005); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Financial assistance from the University of the Free State and SASOL to ACS is gratefully acknowledged. Opinions, findings, conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of SASOL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FI2063).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2004). *SAINT-Plus* (including *XPREP*) and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Kaeding, W. W. (1967). *J. Org. Chem.* **26**, 3144–3148.
- Kawata, T., Uekusa, H., Ohba, S., Furukawa, T., Tokii, T., Muto, Y. & Kato, M. (1992). *Acta Cryst.* **B48**, 253–261.
- Mak, T. C. W. & Yip, W. H. (1990). *Polyhedron*, **9**, 1667–1670.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sunil, A. C., Bezuidenhoudt, B. C. B. & Janse van Rensburg, J. M. (2008). *Acta Cryst.* **E64**, m553–m554.
- Valach, F., Tokarcik, M., Maris, T., Watkin, D. J. & Prout, C. K. (2000). *Z. Kristallogr.* **215**, 56–60.

supplementary materials

Acta Cryst. (2008). E64, m939 [doi:10.1107/S1600536808015924]

Tetrakis(μ -4-ethylbenzoato- $\kappa^2 O:O'$)bis[(4-ethylbenzoic acid- κO)copper(II)]

A. C. Sunil, B. C. B. Bezuidenhoudt and J. M. Janse van Rensburg

Comment

The title compound forms part of the copper(II) complexes of the type $[Cu_2(RCO_2)_4L_2]$ (R =aryl, L =monodentate ligand). This type of complex forms tetra-(carboxylato-O,O') bridges and four of the carboxylate groups hold together two Cu atoms (Fig. 1). The Cu···Cu distance in the title compound is 2.6047 (5) Å, probably displaying weak orbital interaction considering that the van der Waals radius of copper is 2.32 Å. The axial sites of each copper atom are bonded to a monodentate *p*-ethylbenzoic acid ligand. In turn the acid protons are hydrogen bonded to the cage carboxylate O atoms, O—H···O = 166.79° and O···O = 2.645 Å.

Neighbouring molecules stack with overlap between the axially bonded phenyl rings displaying a centroid to centroid distance of 4.2918 (3) Å and an interplanar distance of 3.6277 Å (Fig. 2 A). This inter-molecular interaction influence the dihedral angle displayed between the phenyl rings from the axially bonded monodentate ligands and the carboxylic oxygen plane, O1, O2, O1ⁱ and O2ⁱ ($i = 1 - x, 1 - y, 2 - z$). Molecular packing in the (0 0 h) plane is in a puckered pseudo-hexagonal close packing fashion. This close packing is stabilized by soft inter-molecular C···H contacts ranging from 2.720–2.813 Å (Fig. 2B).

Experimental

The complex $[Cu_2(C_9H_{10}O_2)_4(C_9H_{11}O_2)_2]$ was prepared by heating 4-ethylbenzoic acid (1.77 g, 11.81 mmol), copper carbonate (0.74 g, 3.34 mmol) and magnesium oxide (0.20 g, 4.98 mmol) under reflux, in toluene (15 ml) for 60 h. The product was extacted and crystallized from diethyl ether to yield a blue crystalline solid. (Yield: 80%)

Refinement

The H atoms were positioned geometrically and refined using a riding model with fixed C—H distances of 0.93 Å (CH) [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$] and 0.96 Å (CH₃) [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}$] respectively. Initial positions of methyl H-atoms were obtained from Fourier difference maps and refined as a fixed rotor.

The highest density peak is 0.50 located 0.65 Å from C14 and the deepest hole is -0.37 located at 0.68 Å from Cu1.

supplementary materials

Figures




Fig. 1. A view of (I) showing the atom-numbering scheme with displacement ellipsoids at the 30% probability level, non labelled atoms are symmetric equivalents. For the phenyl C-atoms, the first digit indicates ring number and the second digit the position of the atom in the ring. Symmetry code: 1 - x , 1 - y , 2 - z .




Fig. 2. (A) Hacked lines indicate overlap between ethylbenzoic groups of neighbouring molecules. (B) Indication of pseudo-hexagonal close packing along the c axis.

Tetrakis(μ -4-ethylbenzoato- κ^2 O:O')bis[(4-ethylbenzoic acid- κ O)copper(II)]

Crystal data

[Cu ₂ (C ₉ H ₉ O ₂) ₄ (C ₉ H ₁₀ O ₂) ₂]	$Z = 1$
$M_r = 1024.07$	$F_{000} = 534$
Triclinic, $P\bar{1}$	$D_x = 1.418 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 10.6167 (5) \text{ \AA}$	$\lambda = 0.71069 \text{ \AA}$
$b = 10.7394 (7) \text{ \AA}$	Cell parameters from 4441 reflections
$c = 10.8096 (7) \text{ \AA}$	$\theta = 2.5\text{--}28.2^\circ$
$\alpha = 81.848 (3)^\circ$	$\mu = 0.95 \text{ mm}^{-1}$
$\beta = 88.594 (3)^\circ$	$T = 100 (2) \text{ K}$
$\gamma = 79.468 (2)^\circ$	Cuboid, blue
$V = 1199.47 (12) \text{ \AA}^3$	$0.54 \times 0.4 \times 0.39 \text{ mm}$

Data collection

Bruker Kappa APEXII diffractometer	4721 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.040$
$T = 100(2) \text{ K}$	$\theta_{\max} = 28^\circ$
ω and φ scans	$\theta_{\min} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2004)	$h = -7 \rightarrow 14$
$T_{\min} = 0.628, T_{\max} = 0.708$	$k = -14 \rightarrow 14$
15971 measured reflections	$l = -14 \rightarrow 14$
5683 independent reflections	

*Refinement*Refinement on F^2

H-atom parameters constrained

Least-squares matrix: full

$$w = 1/[\sigma^2(F_o^2) + (0.0447P)^2 + 0.9309P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$R[F^2 > 2\sigma(F^2)] = 0.040$$

$$(\Delta/\sigma)_{\text{max}} = 0.001$$

$$wR(F^2) = 0.103$$

$$\Delta\rho_{\text{max}} = 0.50 \text{ e \AA}^{-3}$$

$$S = 1.04$$

$$\Delta\rho_{\text{min}} = -0.37 \text{ e \AA}^{-3}$$

$$5683 \text{ reflections}$$

Extinction correction: none

$$311 \text{ parameters}$$

Special details

Experimental. The intensity data was collected on a Bruker X8 Apex II 4 K Kappa CCD diffractometer using an exposure time of 2 s/frame. A total of 1507 frames were collected with a frame width of 0.5° covering up to $\theta = 28.0^\circ$ with 98.3% completeness accomplished.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.45033 (3)	0.46672 (2)	0.90347 (2)	0.01213 (9)
O6	0.58444 (17)	0.40202 (18)	0.62540 (16)	0.0239 (4)
H6	0.599	0.4445	0.679	0.036*
O1	0.60816 (15)	0.51776 (15)	0.82165 (14)	0.0152 (3)
O3	0.36215 (15)	0.64398 (14)	0.86973 (14)	0.0166 (3)
O4	0.54987 (16)	0.29874 (15)	0.96412 (14)	0.0181 (3)
O2	0.30749 (15)	0.42705 (15)	1.01089 (14)	0.0167 (3)
O5	0.40263 (15)	0.39621 (15)	0.73471 (14)	0.0168 (3)
C51	0.4404 (2)	0.2808 (2)	0.5616 (2)	0.0144 (4)
C11	0.8123 (2)	0.58002 (19)	0.80261 (19)	0.0124 (4)
C30	0.3784 (2)	0.7244 (2)	0.94028 (19)	0.0140 (4)
C53	0.3163 (2)	0.1251 (2)	0.5207 (2)	0.0192 (5)
H53	0.2542	0.0759	0.546	0.023*
C55	0.4715 (2)	0.1870 (2)	0.3717 (2)	0.0172 (5)
H55	0.5134	0.1806	0.2957	0.021*
C52	0.3464 (2)	0.2079 (2)	0.5971 (2)	0.0172 (5)
H52	0.3038	0.215	0.6727	0.021*
C54	0.3778 (2)	0.1138 (2)	0.4054 (2)	0.0169 (5)
C36	0.2111 (2)	0.8866 (2)	0.8178 (2)	0.0171 (5)
H36	0.1956	0.823	0.773	0.02*
C16	0.9223 (2)	0.5936 (2)	0.8637 (2)	0.0145 (4)
H16	0.9212	0.5918	0.95	0.017*

supplementary materials

C10	0.6968 (2)	0.55566 (19)	0.87604 (19)	0.0133 (4)
C31	0.3059 (2)	0.8585 (2)	0.9087 (2)	0.0143 (4)
C34	0.1628 (2)	1.1064 (2)	0.8569 (2)	0.0184 (5)
C32	0.3321 (2)	0.9558 (2)	0.9709 (2)	0.0191 (5)
H32	0.3968	0.9384	1.0306	0.023*
C56	0.5043 (2)	0.2694 (2)	0.4481 (2)	0.0162 (5)
H56	0.5682	0.3166	0.424	0.019*
C12	0.8152 (2)	0.5865 (2)	0.6725 (2)	0.0146 (4)
H12	0.743	0.5777	0.63	0.018*
C541	0.3421 (3)	0.0255 (2)	0.3213 (2)	0.0235 (5)
H54A	0.3627	-0.0618	0.3633	0.028*
H54B	0.3936	0.0313	0.2461	0.028*
C141	1.1539 (2)	0.6401 (2)	0.5944 (2)	0.0208 (5)
H14A	1.1803	0.571	0.5452	0.025*
H14B	1.2228	0.6391	0.6521	0.025*
C50	0.4725 (2)	0.3652 (2)	0.6484 (2)	0.0155 (5)
C13	0.9247 (2)	0.6059 (2)	0.6070 (2)	0.0158 (5)
H13	0.9246	0.6114	0.5203	0.019*
C15	1.0327 (2)	0.6096 (2)	0.7979 (2)	0.0164 (5)
H15	1.1059	0.6153	0.8408	0.02*
C14	1.0360 (2)	0.6173 (2)	0.6679 (2)	0.0152 (4)
C642	-0.0475 (3)	1.2359 (3)	0.9030 (3)	0.0385 (7)
H64A	-0.0923	1.1768	0.8717	0.058*
H64B	-0.0989	1.32	0.8901	0.058*
H64C	-0.0311	1.21	0.9907	0.058*
C641	0.0790 (3)	1.2371 (2)	0.8338 (2)	0.0254 (6)
H64D	0.0628	1.2612	0.7449	0.03*
H64E	0.1226	1.2997	0.8623	0.03*
C33	0.2619 (3)	1.0788 (2)	0.9443 (2)	0.0229 (5)
H33	0.2812	1.1435	0.9852	0.028*
C35	0.1394 (2)	1.0083 (2)	0.7935 (2)	0.0186 (5)
H35	0.0746	1.0253	0.7339	0.022*
C142	1.1323 (3)	0.7677 (3)	0.5072 (3)	0.0323 (6)
H14C	1.0651	0.7688	0.449	0.048*
H14D	1.2099	0.7775	0.4623	0.048*
H14E	1.1085	0.8367	0.5556	0.048*
C542	0.2016 (3)	0.0542 (2)	0.2846 (2)	0.0277 (6)
H54C	0.1498	0.0463	0.3583	0.042*
H54D	0.1855	-0.0053	0.231	0.042*
H54E	0.1807	0.1397	0.2412	0.042*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.01076 (16)	0.01499 (14)	0.01135 (13)	-0.00383 (10)	0.00010 (10)	-0.00231 (9)
O6	0.0189 (10)	0.0369 (10)	0.0225 (9)	-0.0156 (8)	0.0057 (7)	-0.0139 (7)
O1	0.0110 (8)	0.0221 (8)	0.0148 (7)	-0.0076 (7)	0.0005 (6)	-0.0041 (6)
O3	0.0161 (9)	0.0154 (7)	0.0184 (8)	-0.0022 (6)	-0.0021 (7)	-0.0032 (6)

O4	0.0212 (9)	0.0164 (8)	0.0168 (8)	-0.0025 (7)	-0.0044 (7)	-0.0032 (6)
O2	0.0130 (8)	0.0256 (8)	0.0138 (7)	-0.0087 (7)	0.0014 (6)	-0.0036 (6)
O5	0.0144 (9)	0.0232 (8)	0.0142 (7)	-0.0042 (7)	0.0001 (6)	-0.0059 (6)
C51	0.0119 (12)	0.0176 (10)	0.0129 (10)	-0.0019 (9)	-0.0023 (8)	0.0001 (8)
C11	0.0097 (11)	0.0118 (9)	0.0152 (10)	-0.0021 (8)	-0.0009 (8)	-0.0005 (8)
C30	0.0114 (11)	0.0179 (10)	0.0130 (10)	-0.0047 (9)	0.0049 (8)	-0.0008 (8)
C53	0.0187 (13)	0.0212 (11)	0.0195 (11)	-0.0096 (10)	0.0022 (10)	-0.0010 (9)
C55	0.0149 (12)	0.0217 (11)	0.0139 (10)	0.0000 (9)	0.0025 (9)	-0.0030 (9)
C52	0.0159 (12)	0.0235 (11)	0.0126 (10)	-0.0057 (10)	0.0022 (9)	-0.0015 (9)
C54	0.0152 (12)	0.0178 (11)	0.0169 (11)	0.0007 (9)	-0.0032 (9)	-0.0041 (9)
C36	0.0165 (12)	0.0166 (10)	0.0184 (11)	-0.0035 (9)	0.0004 (9)	-0.0026 (8)
C16	0.0140 (12)	0.0171 (10)	0.0128 (10)	-0.0039 (9)	-0.0025 (9)	-0.0019 (8)
C10	0.0147 (12)	0.0124 (10)	0.0122 (9)	-0.0027 (9)	-0.0021 (8)	0.0009 (8)
C31	0.0111 (11)	0.0164 (10)	0.0155 (10)	-0.0038 (9)	0.0037 (9)	-0.0014 (8)
C34	0.0192 (13)	0.0159 (11)	0.0191 (11)	-0.0037 (9)	0.0064 (9)	0.0001 (9)
C32	0.0186 (13)	0.0215 (11)	0.0174 (11)	-0.0029 (10)	-0.0034 (9)	-0.0040 (9)
C56	0.0107 (12)	0.0200 (11)	0.0180 (11)	-0.0046 (9)	0.0013 (9)	-0.0015 (9)
C12	0.0112 (12)	0.0175 (10)	0.0161 (10)	-0.0041 (9)	-0.0022 (9)	-0.0029 (8)
C541	0.0256 (14)	0.0240 (12)	0.0239 (12)	-0.0073 (11)	0.0018 (11)	-0.0101 (10)
C141	0.0140 (12)	0.0277 (12)	0.0229 (12)	-0.0086 (10)	0.0037 (10)	-0.0059 (10)
C50	0.0142 (12)	0.0175 (10)	0.0145 (10)	-0.0032 (9)	-0.0027 (9)	-0.0008 (8)
C13	0.0145 (12)	0.0196 (11)	0.0133 (10)	-0.0033 (9)	0.0008 (9)	-0.0025 (8)
C15	0.0118 (12)	0.0185 (11)	0.0199 (11)	-0.0054 (9)	-0.0041 (9)	-0.0016 (9)
C14	0.0114 (12)	0.0131 (10)	0.0210 (11)	-0.0030 (9)	0.0021 (9)	-0.0019 (8)
C642	0.0278 (17)	0.0213 (13)	0.065 (2)	-0.0006 (12)	0.0111 (15)	-0.0078 (13)
C641	0.0290 (15)	0.0182 (11)	0.0273 (13)	-0.0024 (11)	0.0028 (11)	-0.0008 (10)
C33	0.0306 (15)	0.0184 (11)	0.0211 (12)	-0.0052 (11)	0.0008 (11)	-0.0067 (9)
C35	0.0147 (13)	0.0198 (11)	0.0202 (11)	-0.0022 (9)	-0.0024 (9)	-0.0003 (9)
C142	0.0232 (15)	0.0422 (16)	0.0296 (14)	-0.0106 (13)	-0.0002 (12)	0.0071 (12)
C542	0.0304 (16)	0.0261 (13)	0.0292 (13)	-0.0071 (11)	-0.0066 (11)	-0.0086 (10)

Geometric parameters (Å, °)

Cu1—O3	1.9498 (15)	C31—C32	1.392 (3)
Cu1—O4	1.9501 (16)	C34—C33	1.394 (3)
Cu1—O2	1.9593 (16)	C34—C35	1.397 (3)
Cu1—O1	2.0040 (16)	C34—C641	1.509 (3)
Cu1—O5	2.1761 (15)	C32—C33	1.387 (3)
Cu1—Cu1 ⁱ	2.6047 (5)	C32—H32	0.93
O6—C50	1.326 (3)	C56—H56	0.93
O6—H6	0.82	C12—C13	1.380 (3)
O1—C10	1.277 (3)	C12—H12	0.93
O3—C30	1.267 (3)	C541—C542	1.517 (4)
O4—C30 ⁱ	1.267 (3)	C541—H54A	0.97
O2—C10 ⁱ	1.261 (2)	C541—H54B	0.97
O5—C50	1.223 (3)	C141—C14	1.505 (3)
C51—C52	1.392 (3)	C141—C142	1.532 (3)
C51—C56	1.397 (3)	C141—H14A	0.97

supplementary materials

C51—C50	1.479 (3)	C141—H14B	0.97
C11—C16	1.396 (3)	C13—C14	1.400 (3)
C11—C12	1.398 (3)	C13—H13	0.93
C11—C10	1.488 (3)	C15—C14	1.396 (3)
C30—O4 ⁱ	1.267 (3)	C15—H15	0.93
C30—C31	1.501 (3)	C642—C641	1.523 (4)
C53—C52	1.378 (3)	C642—H64A	0.96
C53—C54	1.403 (3)	C642—H64B	0.96
C53—H53	0.93	C642—H64C	0.96
C55—C56	1.386 (3)	C641—H64D	0.97
C55—C54	1.388 (3)	C641—H64E	0.97
C55—H55	0.93	C33—H33	0.93
C52—H52	0.93	C35—H35	0.93
C54—C541	1.505 (3)	C142—H14C	0.96
C36—C35	1.381 (3)	C142—H14D	0.96
C36—C31	1.387 (3)	C142—H14E	0.96
C36—H36	0.93	C542—H54C	0.96
C16—C15	1.382 (3)	C542—H54D	0.96
C16—H16	0.93	C542—H54E	0.96
C10—O2 ⁱ	1.261 (2)		
O3—Cu1—O4	169.67 (6)	C55—C56—C51	119.3 (2)
O3—Cu1—O2	89.21 (7)	C55—C56—H56	120.3
O4—Cu1—O2	89.79 (7)	C51—C56—H56	120.3
O3—Cu1—O1	89.64 (7)	C13—C12—C11	120.3 (2)
O4—Cu1—O1	89.46 (7)	C13—C12—H12	119.8
O2—Cu1—O1	169.42 (6)	C11—C12—H12	119.8
O3—Cu1—O5	100.25 (6)	C54—C541—C542	113.8 (2)
O4—Cu1—O5	90.05 (6)	C54—C541—H54A	108.8
O2—Cu1—O5	99.99 (6)	C542—C541—H54A	108.8
O1—Cu1—O5	90.57 (6)	C54—C541—H54B	108.8
O3—Cu1—Cu1 ⁱ	86.32 (5)	C542—C541—H54B	108.8
O4—Cu1—Cu1 ⁱ	83.36 (5)	H54A—C541—H54B	107.7
O2—Cu1—Cu1 ⁱ	87.95 (5)	C14—C141—C142	112.7 (2)
O1—Cu1—Cu1 ⁱ	81.48 (4)	C14—C141—H14A	109.1
O5—Cu1—Cu1 ⁱ	169.69 (5)	C142—C141—H14A	109.1
C50—O6—H6	109.5	C14—C141—H14B	109.1
C10—O1—Cu1	125.79 (14)	C142—C141—H14B	109.1
C30—O3—Cu1	120.62 (14)	H14A—C141—H14B	107.8
C30 ⁱ —O4—Cu1	124.01 (14)	O5—C50—O6	123.3 (2)
C10 ⁱ —O2—Cu1	120.97 (15)	O5—C50—C51	122.7 (2)
C50—O5—Cu1	128.90 (15)	O6—C50—C51	113.97 (19)
C52—C51—C56	119.6 (2)	C12—C13—C14	121.5 (2)
C52—C51—C50	118.4 (2)	C12—C13—H13	119.2
C56—C51—C50	122.0 (2)	C14—C13—H13	119.2
C16—C11—C12	118.5 (2)	C16—C15—C14	120.9 (2)
C16—C11—C10	120.04 (19)	C16—C15—H15	119.5
C12—C11—C10	121.5 (2)	C14—C15—H15	119.5

O3—C30—O4 ⁱ	125.7 (2)	C15—C14—C13	117.8 (2)
O3—C30—C31	117.29 (19)	C15—C14—C141	121.5 (2)
O4 ⁱ —C30—C31	117.03 (19)	C13—C14—C141	120.7 (2)
C52—C53—C54	121.0 (2)	C641—C642—H64A	109.5
C52—C53—H53	119.5	C641—C642—H64B	109.5
C54—C53—H53	119.5	H64A—C642—H64B	109.5
C56—C55—C54	121.9 (2)	C641—C642—H64C	109.5
C56—C55—H55	119.1	H64A—C642—H64C	109.5
C54—C55—H55	119.1	H64B—C642—H64C	109.5
C53—C52—C51	120.3 (2)	C34—C641—C642	110.1 (2)
C53—C52—H52	119.9	C34—C641—H64D	109.6
C51—C52—H52	119.9	C642—C641—H64D	109.6
C55—C54—C53	117.9 (2)	C34—C641—H64E	109.6
C55—C54—C541	121.5 (2)	C642—C641—H64E	109.6
C53—C54—C541	120.6 (2)	H64D—C641—H64E	108.2
C35—C36—C31	120.3 (2)	C32—C33—C34	120.8 (2)
C35—C36—H36	119.8	C32—C33—H33	119.6
C31—C36—H36	119.8	C34—C33—H33	119.6
C15—C16—C11	120.9 (2)	C36—C35—C34	121.1 (2)
C15—C16—H16	119.5	C36—C35—H35	119.5
C11—C16—H16	119.5	C34—C35—H35	119.5
O2 ⁱ —C10—O1	123.7 (2)	C141—C142—H14C	109.5
O2 ⁱ —C10—C11	117.84 (19)	C141—C142—H14D	109.5
O1—C10—C11	118.46 (18)	H14C—C142—H14D	109.5
C36—C31—C32	119.2 (2)	C141—C142—H14E	109.5
C36—C31—C30	120.3 (2)	H14C—C142—H14E	109.5
C32—C31—C30	120.5 (2)	H14D—C142—H14E	109.5
C33—C34—C35	118.2 (2)	C541—C542—H54C	109.5
C33—C34—C641	121.5 (2)	C541—C542—H54D	109.5
C35—C34—C641	120.2 (2)	H54C—C542—H54D	109.5
C33—C32—C31	120.3 (2)	C541—C542—H54E	109.5
C33—C32—H32	119.9	H54C—C542—H54E	109.5
C31—C32—H32	119.9	H54D—C542—H54E	109.5

Symmetry codes: (i) $-x+1, -y+1, -z+2$.

supplementary materials

Fig. 1




Fig. 2

